

Article

Abstract

Tabacay micro-basin is very important for 27 communes that lives in it, from which they extract resources, develop economic activities and life overall. This micro-basin is also important for Azogues city, which takes water from its rivers for supply the population. This study analyzes land use dynamics and vegetation's quality during 2005-2017 period, using Landsat satellite images along with geographic information systems, for determining critical areas for management and conservation of Tabacay micro-basin. Four land use classes were determined: agricultural, forestry, urban and moor. Analysis results show that agricultural and urban land use has grown in surface terms, as well as forestry and moor surface has decreased. Moor land use has been the most affected by land use dynamics. In terms of surface, this land use shows a 6 ha/year loss during analysis period, which is equivalent to 54% of its initial surface. Analysis also shows that most affected zone by forest and moor fragmentation and surface loss is micro-basin's high part, having as main cause of the problem the agricultural frontier expansion and urbanization.

Keywords: Geographic information systems, Landsat, land use, Tabacay, micro-basin, multitemporal analysis, remote sensing.

Introduction

Tabacay micro-basin has a critical role in Azogues city development. Also for 27 communes which lives in its territory. Although its vital role, there are various social and environmental issues affecting, overall, medium and upper part.

Two of this issues are erosion and landslides, which together are destroying potential farmlands, increasing sediment disposal in many streams. Soil erosion is caused, mainly, for deforestation and inappropriate cropping practices.

Another issue affecting Tabacay micro-basin is urban growth to upper part, being this consequence of building an massive, unplanned and poor thecnical road network. Absense of information about LULC evolution is a constraint for Tabacay micro-basin management. Studies about this variables allows understanding of causes and consequences of deforestation, degradation, desertification and biodiversity loss processes in a specific area. LULC assesments are also crucial for rural growth planning and food safety monitoring.

Materials and methods

Study area

Tabacay micro-basin is located in Cañar province, Ecuador. It also belongs to Paute basin. The area covers a surface of 68.3 Km²

Results and discuss

Conclusions

References