

Question 1(a) Can this be done using the `sample` function? If yes, what table would we call `sample` on?
(1 point)

Yes, with 'top_movies'

Question 1(b) Can this be done using the `np.random.choice` function? If yes, what array would we call `np.random.choice` on? **(1 point)**

Yes, with 'studios__of__all__movies'

Question 1(c) Can this be done using the `sample_proportions` function? If yes, what array would we call `sample_proportions` on? **(1 point)**

Yes, with `'studio__proportions__only'`

Question 2(a) Can this be done using the `sample` function? If yes, what table would we call `sample` on?
(1 point)

Yes, with 'top_movies'

Question 2(b) Can this be done using the `np.random.choice` function? If yes, what array would we call `np.random.choice` on? **(1 point)**

Yes, with 'studios__of__all__movies'

Question 2(c) Can this be done using the `sample_proportions` function? If yes, what array would we call `sample_proportions` on? **(1 point)**

No

Question 5. Given your observed value, do you believe that Jade's model is reasonable, or is our alternative (that our deck is rigged) more likely? Explain your answer using the histogram of statistics simulated using Jade's model (produced above). **(4 Points)**

as we can see, the frequency 8 appears is low, the highest frequency is somewhat between 3-> 5 => alternative model is more likely

