

Chapter 7. Continuous optimization





Chapter 7. Continuous optimization





- 1 7.1 Optimization Using Gradient Descent
- 2 7.2 Constrained Optimization and Lagrange Multipliers
- 3 7.3 Convex Optimization
 - 7.3.1 Linear programming
 - 7.3.2 Quadratic Programming

7.1 Optimization Using Gradient Descent



7.1 Optimization Using Gradient Descent



Consider the problem

$$\min_x f(x),$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable.

Gradient descent is a first-order optimization algorithm. To find a local minimum of a function using gradient descent, one takes steps proportional to the negative of the gradient of the function at the current point.

Consider the problem

$$\min_x f(x),$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable.

Gradient descent is a first-order optimization algorithm. To find a local minimum of a function using gradient descent, one takes steps proportional to the negative of the gradient of the function at the current point.

We can see that, if

$$x_1 = x_0 - \gamma ((\nabla f)(x_0))^T$$

for a small **step size (learning rate)** $\gamma > 0$ then $f(x_1) \leq f(x_0)$.

Consider the problem

$$\min_x f(x),$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable.

Gradient descent is a first-order optimization algorithm. To find a local minimum of a function using gradient descent, one takes steps proportional to the negative of the gradient of the function at the current point.

We can see that, if

$$x_1 = x_0 - \gamma ((\nabla f)(x_0))^T$$

for a small **step size (learning rate)** $\gamma > 0$ then $f(x_1) \leq f(x_0)$.

Iterate

$$x_{i+1} = x_i - \gamma_i ((\nabla f)(x_i))^T.$$

For suitable learning rate γ_i , the sequence $f(x_0) \geq f(x_1) \geq \dots$ converges to a local minimum.

Consider the problem

$$\min_x f(x),$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable.

Gradient descent is a first-order optimization algorithm. To find a local minimum of a function using gradient descent, one takes steps proportional to the negative of the gradient of the function at the current point.

We can see that, if

$$x_1 = x_0 - \gamma ((\nabla f)(x_0))^T$$

for a small **step size (learning rate)** $\gamma > 0$ then $f(x_1) \leq f(x_0)$.

Iterate

$$x_{i+1} = x_i - \gamma_i ((\nabla f)(x_i))^T.$$

For suitable learning rate γ_i , the sequence $f(x_0) \geq f(x_1) \geq \dots$ converges to a local minimum.

Example

Consider the problem of minimizing $f(x, y) = 2x^2 - 2xy + 1.5y^2$ using the gradient descent method.

We have

$$\nabla f(x, y) = \begin{bmatrix} 4x - 2y \\ -2x + 3y \end{bmatrix}$$

Example

Consider the problem of minimizing $f(x, y) = 2x^2 - 2xy + 1.5y^2$ using the gradient descent method.

We have

$$\nabla f(x, y) = \begin{bmatrix} 4x - 2y \\ -2x + 3y \end{bmatrix}$$

We will start from the point $(x_0, y_0) = (1, 2)$ and learning rate $\gamma = 0.05$.

Example

Consider the problem of minimizing $f(x, y) = 2x^2 - 2xy + 1.5y^2$ using the gradient descent method.

We have

$$\nabla f(x, y) = \begin{bmatrix} 4x - 2y \\ -2x + 3y \end{bmatrix}$$

We will start from the point $(x_0, y_0) = (1, 2)$ and learning rate $\gamma = 0.05$. Iterative:

$$\begin{bmatrix} x_{i+1} \\ y_{i+1} \end{bmatrix} = \begin{bmatrix} x_i \\ y_i \end{bmatrix} - \gamma \begin{bmatrix} 4x_i - 2y_i \\ -2x_i + 3y_i \end{bmatrix} = \begin{bmatrix} 0.8x_i + 0.1y_i \\ 0.1x_i + 0.85y_i \end{bmatrix}$$

Example

Consider the problem of minimizing $f(x, y) = 2x^2 - 2xy + 1.5y^2$ using the gradient descent method.

We have

$$\nabla f(x, y) = \begin{bmatrix} 4x - 2y \\ -2x + 3y \end{bmatrix}$$

We will start from the point $(x_0, y_0) = (1, 2)$ and learning rate $\gamma = 0.05$. Iterative:

$$\begin{bmatrix} x_{i+1} \\ y_{i+1} \end{bmatrix} = \begin{bmatrix} x_i \\ y_i \end{bmatrix} - \gamma \begin{bmatrix} 4x_i - 2y_i \\ -2x_i + 3y_i \end{bmatrix} = \begin{bmatrix} 0.8x_i + 0.1y_i \\ 0.1x_i + 0.85y_i \end{bmatrix}$$

We obtain

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1.71 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0.971 \\ 1.5535 \end{bmatrix}, \dots$$

Example

Consider the problem of minimizing $f(x, y) = 2x^2 - 2xy + 1.5y^2$ using the gradient descent method.

We have

$$\nabla f(x, y) = \begin{bmatrix} 4x - 2y \\ -2x + 3y \end{bmatrix}$$

We will start from the point $(x_0, y_0) = (1, 2)$ and learning rate $\gamma = 0.05$. Iterative:

$$\begin{bmatrix} x_{i+1} \\ y_{i+1} \end{bmatrix} = \begin{bmatrix} x_i \\ y_i \end{bmatrix} - \gamma \begin{bmatrix} 4x_i - 2y_i \\ -2x_i + 3y_i \end{bmatrix} = \begin{bmatrix} 0.8x_i + 0.1y_i \\ 0.1x_i + 0.85y_i \end{bmatrix}$$

We obtain

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1.71 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0.971 \\ 1.5535 \end{bmatrix}, \dots$$

Notice that the optimal solution is $(x, y) = (0, 0)$.

Example

Consider the problem of minimizing $f(x, y) = 2x^2 - 2xy + 1.5y^2$ using the gradient descent method.

We have

$$\nabla f(x, y) = \begin{bmatrix} 4x - 2y \\ -2x + 3y \end{bmatrix}$$

We will start from the point $(x_0, y_0) = (1, 2)$ and learning rate $\gamma = 0.05$. Iterative:

$$\begin{bmatrix} x_{i+1} \\ y_{i+1} \end{bmatrix} = \begin{bmatrix} x_i \\ y_i \end{bmatrix} - \gamma \begin{bmatrix} 4x_i - 2y_i \\ -2x_i + 3y_i \end{bmatrix} = \begin{bmatrix} 0.8x_i + 0.1y_i \\ 0.1x_i + 0.85y_i \end{bmatrix}$$

We obtain

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1.71 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0.971 \\ 1.5535 \end{bmatrix}, \dots$$

Notice that the optimal solution is $(x, y) = (0, 0)$.

Exercise. Write a python programme to find the local minima of the function

$$f(x) = x^2 - 2x + 8$$

starting from point $x_0 = 2$ by the gradient descent algorithm. Let us assume the learning rate 0.01 and the difference between x values from two consecutive iterations is less than 0.001.

Choosing a good step-size is important in gradient descent:

- If the step-size is too small, gradient descent can be slow.

Choosing a good step-size is important in gradient descent:

- If the step-size is too small, gradient descent can be slow.
- If the step-size is chosen too large, gradient descent can overshoot, fail to converge, or even diverge.

Choosing a good step-size is important in gradient descent:

- If the step-size is too small, gradient descent can be slow.
- If the step-size is chosen too large, gradient descent can overshoot, fail to converge, or even diverge.

Choosing a step size:

$$\gamma_i^* = \underset{\gamma}{\operatorname{argmin}} f(x_i - \gamma \nabla f(x_i))^T$$

Choosing a good step-size is important in gradient descent:

- If the step-size is too small, gradient descent can be slow.
- If the step-size is chosen too large, gradient descent can overshoot, fail to converge, or even diverge.

Choosing a step size:

$$\gamma_i^* = \underset{\gamma}{\operatorname{argmin}} f(x_i - \gamma \nabla f(x_i))^T$$

Remembers the update $\Delta x_i = x_i - x_{i-1}$ at each iteration i and determines the next update as a linear combination of the current and previous gradients:

$$x_{i+1} = x_i - \gamma_i ((\nabla f)(x_i))^T + \alpha \Delta x_i$$

$$\Delta x_i = x_i - x_{i-1} = \alpha \Delta x_{i-1} - \gamma_{i-1} ((\nabla f)(x_{i-1}))^T.$$

7.2 Constrained Optimization and Lagrange Multi

Consider the **constrained optimization problem**

$$\begin{aligned} & \min f(x) \\ & \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

We can convert the constrained problem into an unconstrained one:

$$J(x) = f(x) + \sum_{i=1}^m \mathbb{1}(g_i(x)),$$

7.2 Constrained Optimization and Lagrange Multi

Consider the **constrained optimization problem**

$$\begin{aligned} \min f(x) \\ \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

We can convert the constrained problem into an unconstrained one:

$$J(x) = f(x) + \sum_{i=1}^m \mathbb{1}(g_i(x)),$$

where $\mathbb{1}(z)$ is an infinite step function

$$\mathbb{1}(z) = \begin{cases} 0 & \text{if } z \leq 0 \\ \infty & \text{otherwise} \end{cases}.$$

7.2 Constrained Optimization and Lagrange Multi

Consider the **constrained optimization problem**

$$\begin{aligned} \min f(x) \\ \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

We can convert the constrained problem into an unconstrained one:

$$J(x) = f(x) + \sum_{i=1}^m \mathbb{1}(g_i(x)),$$

where $\mathbb{1}(z)$ is an infinite step function

$$\mathbb{1}(z) = \begin{cases} 0 & \text{if } z \leq 0 \\ \infty & \text{otherwise} \end{cases}.$$

The associated **the Lagrangian**

$$\begin{aligned} \mathcal{L}(x, \lambda) &= f(x) + \sum_{i=1}^m \lambda_i g_i(x) \\ &= f(x) + \lambda^T g(x), \end{aligned}$$

7.2 Constrained Optimization and Lagrange Multi

Consider the **constrained optimization problem**

$$\begin{aligned} \min f(x) \\ \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

We can convert the constrained problem into an unconstrained one:

$$J(x) = f(x) + \sum_{i=1}^m \mathbb{1}(g_i(x)),$$

where $\mathbb{1}(z)$ is an infinite step function

$$\mathbb{1}(z) = \begin{cases} 0 & \text{if } z \leq 0 \\ \infty & \text{otherwise} \end{cases}.$$

The associated **the Lagrangian**

$$\begin{aligned} \mathcal{L}(x, \lambda) &= f(x) + \sum_{i=1}^m \lambda_i g_i(x) \\ &= f(x) + \lambda^T g(x), \end{aligned}$$

Definition

The problem

$$\begin{aligned} & \min f(x) \\ & \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

is known as the **primal problem**, corresponding to the primal variable $x \in \mathbb{R}^D$.

Definition

The problem

$$\begin{aligned} & \min f(x) \\ & \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

is known as the **primal problem**, corresponding to the primal variable $x \in \mathbb{R}^D$.

The **associated Lagrangian dual problem** is given by

$$\max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{D}(\lambda) = \max_{\lambda \in (\mathbb{R}^+)^m} \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda),$$

Definition

The problem

$$\begin{aligned} & \min f(x) \\ & \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

is known as the **primal problem**, corresponding to the primal variable $x \in \mathbb{R}^D$.

The **associated Lagrangian dual problem** is given by

$$\max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{D}(\lambda) = \max_{\lambda \in (\mathbb{R}^+)^m} \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda),$$

where $\lambda = [\lambda_1, \dots, \lambda_m]$ are dual variables and $\mathcal{D}(\lambda) = \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda)$.

Definition

The problem

$$\begin{aligned} & \min f(x) \\ & \text{subject to } g_i(x) \leq 0, i = 1, \dots, m \end{aligned}$$

is known as the **primal problem**, corresponding to the primal variable $x \in \mathbb{R}^D$.

The **associated Lagrangian dual problem** is given by

$$\max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{D}(\lambda) = \max_{\lambda \in (\mathbb{R}^+)^m} \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda),$$

where $\lambda = [\lambda_1, \dots, \lambda_m]$ are dual variables and $\mathcal{D}(\lambda) = \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda)$.

Theorem (Minimax inequality)

For any function with two arguments $\varphi(x, y)$:

$$\max_y \min_x \varphi(x, y) \leq \min_x \max_y \varphi(x, y)$$

Theorem (Minimax inequality)

For any function with two arguments $\varphi(x, y)$:

$$\max_y \min_x \varphi(x, y) \leq \min_x \max_y \varphi(x, y)$$

We can see that

$$J(x) = \max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{L}(x, \lambda).$$

Theorem (Minimax inequality)

For any function with two arguments $\varphi(x, y)$:

$$\max_y \min_x \varphi(x, y) \leq \min_x \max_y \varphi(x, y)$$

We can see that

$$J(x) = \max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{L}(x, \lambda).$$

Minimax inequality implies that

Theorem (Minimax inequality)

For any function with two arguments $\varphi(x, y)$:

$$\max_y \min_x \varphi(x, y) \leq \min_x \max_y \varphi(x, y)$$

We can see that

$$J(x) = \max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{L}(x, \lambda).$$

Minimax inequality implies that

Corollary (weak duality)

$$\min_{x \in \mathbb{R}^D} \max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{L}(x, \lambda) \geq \max_{\lambda \in (\mathbb{R}^+)^m} \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda).$$

Theorem (Minimax inequality)

For any function with two arguments $\varphi(x, y)$:

$$\max_y \min_x \varphi(x, y) \leq \min_x \max_y \varphi(x, y)$$

We can see that

$$J(x) = \max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{L}(x, \lambda).$$

Minimax inequality implies that

Corollary (weak duality)

$$\min_{x \in \mathbb{R}^D} \max_{\lambda \in (\mathbb{R}^+)^m} \mathcal{L}(x, \lambda) \geq \max_{\lambda \in (\mathbb{R}^+)^m} \min_{x \in \mathbb{R}^D} \mathcal{L}(x, \lambda).$$

Definition

A set \mathcal{C} is **convex** if for any $x, y \in \mathcal{C}$ then the line segment joining x, y lie inside \mathcal{C} , i.e

$$\theta x + (1 - \theta)y \in \mathcal{C}.$$

Example

- The closed interval $[0, 8]$ is convex.

Definition

A set \mathcal{C} is **convex** if for any $x, y \in \mathcal{C}$ then the line segment joining x, y lie inside \mathcal{C} , i.e

$$\theta x + (1 - \theta)y \in \mathcal{C}.$$

Example

- The closed interval $[0, 8]$ is convex.
- The opened ball $B(a, r) := \{x \in \mathbb{R}^D : \|x - a\|_2 < r\}$ is convex.

Definition

A set \mathcal{C} is **convex** if for any $x, y \in \mathcal{C}$ then the line segment joining x, y lie inside \mathcal{C} , i.e

$$\theta x + (1 - \theta)y \in \mathcal{C}.$$

Example

- The closed interval $[0, 8]$ is convex.
- The opened ball $B(a, r) := \{x \in \mathbb{R}^D : \|x - a\|_2 < r\}$ is convex.
- The annulus $A(r, R) := \{x \in \mathbb{R}^2 : 0 < r_1 < \|x\|_2 < r_2\}$ is not convex.

Definition

A set \mathcal{C} is **convex** if for any $x, y \in \mathcal{C}$ then the line segment joining x, y lie inside \mathcal{C} , i.e

$$\theta x + (1 - \theta)y \in \mathcal{C}.$$

Example

- The closed interval $[0, 8]$ is convex.
- The opened ball $B(a, r) := \{x \in \mathbb{R}^D : \|x - a\|_2 < r\}$ is convex.
- The annulus $A(r, R) := \{x \in \mathbb{R}^2 : 0 < r_1 < \|x\|_2 < r_2\}$ is not convex.

Definition

A set \mathcal{C} is **convex** if for any $x, y \in \mathcal{C}$ then the line segment joining x, y lie inside \mathcal{C} , i.e

$$\theta x + (1 - \theta)y \in \mathcal{C}.$$

Example

- The closed interval $[0, 8]$ is convex.
- The opened ball $B(a, r) := \{x \in \mathbb{R}^D : \|x - a\|_2 < r\}$ is convex.
- The annulus $A(r, R) := \{x \in \mathbb{R}^2 : 0 < r_1 < \|x\|_2 < r_2\}$ is not convex.

Definition

Let $f: \mathbb{R}^D \rightarrow \mathbb{R}$ be a function whose domain is a convex set. The function f is a **convex function** if for all x, y in the domain of f , we have

$$f[\theta x + (1 - \theta)y] \leq \theta f(x) + (1 - \theta)f(y), \quad \forall 0 \leq \theta \leq 1.$$

Definition

Let $f: \mathbb{R}^D \rightarrow \mathbb{R}$ be a function whose domain is a convex set. The function f is a **convex function** if for all x, y in the domain of f , we have

$$f[\theta x + (1 - \theta)y] \leq \theta f(x) + (1 - \theta)f(y), \quad \forall 0 \leq \theta \leq 1.$$

Theorem

Suppose that $f: \mathbb{R}^D \rightarrow \mathbb{R}$ is differentiable. Then f is convex if and only if for any two points x, y it holds that

Definition

Let $f: \mathbb{R}^D \rightarrow \mathbb{R}$ be a function whose domain is a convex set. The function f is a **convex function** if for all x, y in the domain of f , we have

$$f[\theta x + (1 - \theta)y] \leq \theta f(x) + (1 - \theta)f(y), \quad \forall 0 \leq \theta \leq 1.$$

Theorem

Suppose that $f: \mathbb{R}^D \rightarrow \mathbb{R}$ is differentiable. Then f is convex if and only if for any two points x, y it holds that

$$f(y) \geq f(x) + \nabla_x f(x)^T (y - x).$$

Definition

Let $f: \mathbb{R}^D \rightarrow \mathbb{R}$ be a function whose domain is a convex set. The function f is a **convex function** if for all x, y in the domain of f , we have

$$f[\theta x + (1 - \theta)y] \leq \theta f(x) + (1 - \theta)f(y), \quad \forall 0 \leq \theta \leq 1.$$

Theorem

Suppose that $f: \mathbb{R}^D \rightarrow \mathbb{R}$ is differentiable. Then f is convex if and only if for any two points x, y it holds that

$$f(y) \geq f(x) + \nabla_x f(x)^T (y - x).$$

*Furthermore, if $f(x)$ is twice differentiable then f is convex if and only if its Hessian $\nabla_x^2 f(x)$ is **positive semi-definite**.*

Definition

Let $f: \mathbb{R}^D \rightarrow \mathbb{R}$ be a function whose domain is a convex set. The function f is a **convex function** if for all x, y in the domain of f , we have

$$f[\theta x + (1 - \theta)y] \leq \theta f(x) + (1 - \theta)f(y), \quad \forall 0 \leq \theta \leq 1.$$

Theorem


Suppose that $f: \mathbb{R}^D \rightarrow \mathbb{R}$ is differentiable. Then f is convex if and only if for any two points x, y it holds that

$$f(y) \geq f(x) + \nabla_x f(x)^T (y - x).$$

*Furthermore, if $f(x)$ is twice differentiable then f is convex if and only if its Hessian $\nabla_x^2 f(x)$ is **positive semi-definite**.*

Example

The function $f(x) = x \ln x$ is convex on $x > 0$.




Example

The function $f(x, y) = x^2 + y^2$ is convex on \mathbb{R}^2

Example

The function $f(x) = x \ln x$ is convex on $x > 0$.




Example

The function $f(x, y) = x^2 + y^2$ is convex on \mathbb{R}^2 since its Hessian

$$\nabla_{x,y}^2 f(x, y) = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \text{ is positive definite.}$$

Example

The function $f(x) = x \ln x$ is convex on $x > 0$.



Example

The function $f(x, y) = x^2 + y^2$ is convex on \mathbb{R}^2 since its Hessian

$$\nabla_{x,y}^2 f(x, y) = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \text{ is positive definite.}$$

Definition

A constrained optimization problem is called a **convex optimization problem** if

$$\begin{aligned} & \min_x f(x) \\ & \text{subject to } g_i(x) \leq 0, \text{ for all } i = 1, \dots, m \\ & \quad h_j(x) = 0, \text{ for all } j = 1, \dots, n \end{aligned}$$

where functions $f(x), g_i(x)$ are convex functions and $\{x : h_j(x) = 0\}$ are convex sets.

Definition

A constrained optimization problem is called a **convex optimization problem** if

$$\begin{aligned} & \min_x f(x) \\ & \text{subject to } g_i(x) \leq 0, \text{ for all } i = 1, \dots, m \\ & \quad h_j(x) = 0, \text{ for all } j = 1, \dots, n \end{aligned}$$

where functions $f(x), g_i(x)$ are convex functions and $\{x : h_j(x) = 0\}$ are convex sets.

Definition

A constrained optimization problem is called a **convex optimization problem** if

$$\begin{aligned} & \min_x f(x) \\ & \text{subject to } g_i(x) \leq 0, \text{ for all } i = 1, \dots, m \\ & \quad h_j(x) = 0, \text{ for all } j = 1, \dots, n \end{aligned}$$

where functions $f(x), g_i(x)$ are convex functions and $\{x : h_j(x) = 0\}$ are convex sets.

7.3.1 Linear programming

We consider a special cases of convex optimization problem called **linear program**

$$\min_{x \in \mathbb{R}^n} c^T x \quad (1)$$

subject to $Ax \leq b$

where $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$. The Lagrangian is given by

$$\begin{aligned} \mathcal{L}(x, \lambda) &= c^T x + \lambda^T (Ax - b) \\ &= (c^T + A^T \lambda)^T x - \lambda^T b. \end{aligned}$$

7.3.1 Linear programming

We consider a special cases of convex optimization problem called **linear program**

$$\min_{x \in \mathbb{R}^n} c^T x \quad (1)$$

subject to $Ax \leq b$

where $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$. The Lagrangian is given by

$$\begin{aligned} \mathcal{L}(x, \lambda) &= c^T x + \lambda^T (Ax - b) \\ &= (c^T + A^T \lambda)^T x - \lambda^T b. \end{aligned}$$

We have

$$\frac{\partial \mathcal{L}(x, \lambda)}{\partial x} = c^T + A^T \lambda = 0.$$

7.3.1 Linear programming

We consider a special cases of convex optimization problem called **linear program**

$$\min_{x \in \mathbb{R}^n} c^T x \quad (1)$$

subject to $Ax \leq b$

where $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$. The Lagrangian is given by

$$\begin{aligned} \mathcal{L}(x, \lambda) &= c^T x + \lambda^T (Ax - b) \\ &= (c^T + A^T \lambda)^T x - \lambda^T b. \end{aligned}$$

We have

$$\frac{\partial \mathcal{L}(x, \lambda)}{\partial x} = c^T + A^T \lambda = 0.$$

Hence the dual Lagrangian

$$\mathcal{D}(\lambda) = -\lambda^T b.$$

7.3.1 Linear programming

We consider a special cases of convex optimization problem called **linear program**

$$\min_{x \in \mathbb{R}^n} c^T x \quad (1)$$

subject to $Ax \leq b$

where $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$. The Lagrangian is given by

$$\begin{aligned} \mathcal{L}(x, \lambda) &= c^T x + \lambda^T (Ax - b) \\ &= (c^T + A^T \lambda)^T x - \lambda^T b. \end{aligned}$$

We have

$$\frac{\partial \mathcal{L}(x, \lambda)}{\partial x} = c^T + A^T \lambda = 0.$$

Hence the dual Lagrangian

$$\mathcal{D}(\lambda) = -\lambda^T b.$$

The dual optimization

$$\begin{aligned} & \max_{\lambda \in (\mathbb{R}^+)^m} -b^T \lambda \\ & \text{subject to } c + A^T \lambda = 0. \end{aligned} \tag{2}$$

This is also a linear program, but with m variables. We can solve the primal (1) or the dual (2) program depending on whether m or n is larger.

The dual optimization

$$\begin{aligned} & \max_{\lambda \in (\mathbb{R}^+)^m} -b^T \lambda \\ & \text{subject to } c + A^T \lambda = 0. \end{aligned} \tag{2}$$

This is also a linear program, but with m variables. We can solve the primal (1) or the dual (2) program depending on whether m or n is larger.

7.3.2 Quadratic Programming

Consider a **quadratic program**:

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & \frac{1}{2} x^T Q x + c^T x \\ \text{subject to} \quad & A x \leq b \end{aligned} \tag{3}$$

where $Q \in \mathbb{R}^{n \times n}$ is positive definite, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ and $c \in \mathbb{R}^n$.

7.3.2 Quadratic Programming

Consider a **quadratic program**:

$$\min_{x \in \mathbb{R}^n} \frac{1}{2} x^T Q x + c^T x \quad (3)$$

subject to $Ax \leq b$

where $Q \in \mathbb{R}^{n \times n}$ is positive definite, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ and $c \in \mathbb{R}^n$.

The Lagrange is given by

$$\begin{aligned} \mathcal{L}(x, \lambda) &= \frac{1}{2} x^T Q x + c^T x + \lambda^T (Ax - b) \\ &= \frac{1}{2} x^T Q x + (c + A^T \lambda)^T x - \lambda^T b. \end{aligned}$$

7.3.2 Quadratic Programming

Consider a **quadratic program**:

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & \frac{1}{2} x^T Q x + c^T x \\ \text{subject to} \quad & Ax \leq b \end{aligned} \tag{3}$$

where $Q \in \mathbb{R}^{n \times n}$ is positive definite, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ and $c \in \mathbb{R}^n$.

The Lagrange is given by

$$\begin{aligned} \mathcal{L}(x, \lambda) &= \frac{1}{2} x^T Q x + c^T x + \lambda^T (Ax - b) \\ &= \frac{1}{2} x^T Q x + (c + A^T \lambda)^T x - \lambda^T b. \end{aligned}$$

The partial derivative

$$\frac{\mathcal{L}(x, \lambda)}{\partial x} = Qx + (c + A^T \lambda) = 0 \Rightarrow x = -Q^{-1}(c + A^T \lambda)$$

We get the dual Lagrangian

$$\mathcal{D}(\lambda) = -\frac{1}{2}(c + A^T \lambda)^T Q^{-1}(c + A^T \lambda) - \lambda^T b.$$

The partial derivative

$$\frac{\mathcal{L}(x, \lambda)}{\partial x} = Qx + (c + A^T \lambda) = 0 \Rightarrow x = -Q^{-1}(c + A^T \lambda)$$

We get the dual Lagrangian

$$\mathcal{D}(\lambda) = -\frac{1}{2}(c + A^T \lambda)^T Q^{-1}(c + A^T \lambda) - \lambda^T b.$$

Therefore, the dual optimization problem is given by

$$\max_{\lambda \in (\mathbb{R}^+)^m} -\frac{1}{2}(c + A^T \lambda)^T Q^{-1}(c + A^T \lambda) - \lambda^T b$$

subject to $\lambda \geq 0$.

The partial derivative

$$\frac{\mathcal{L}(x, \lambda)}{\partial x} = Qx + (c + A^T \lambda) = 0 \Rightarrow x = -Q^{-1}(c + A^T \lambda)$$

We get the dual Lagrangian

$$\mathcal{D}(\lambda) = -\frac{1}{2}(c + A^T \lambda)^T Q^{-1}(c + A^T \lambda) - \lambda^T b.$$

Therefore, the dual optimization problem is given by

$$\max_{\lambda \in (\mathbb{R}^+)^m} -\frac{1}{2}(c + A^T \lambda)^T Q^{-1}(c + A^T \lambda) - \lambda^T b$$

subject to $\lambda \geq 0$.

We have studied:

- optimization using gradient method;
- constrained optimization;
- convex optimization.

Exercises for practice: 7.1-7.8 (page 247, 248).