

Project: Cross-validation for model selection (notes on papers)

Diego Urdapilleta de la Parra

March 20, 2025

Note: I removed the boxes to keep the proposition enumeration consistent, and to make the text easier to read.

Contents

1	Linear Model Selection	2
1.1	Setup and preliminary results	2
1.2	Shao, 1993: Notes on Leave-One-Out CV	3
1.3	Shao, 1997	5
1.3.1	The case of $\lambda_n \equiv 2$	6
2	Nonlinear Model Selection	7
2.1	Yang, 2007	7
2.1.1	Single-split cross-validation (the Hold-out)	7
2.1.2	Voting cross-validation with multiple splits	8

1 Linear Model Selection

1.1 Setup and preliminary results

Let n, p_n be positive integers and $\mathcal{D}_n := \{(y_i, \mathbf{x}_i) : i \in [n]\}$ be a set of independent data points drawn from a distribution $\mathbb{P}_{y, \mathbf{x}}$ for $(y, \mathbf{x}) \in \mathbb{R}^{1+p_n}$. We treat the \mathbf{x}_i as predictors of the outcome y_i , and we assume a linear model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{e}$$

where $\mathbf{X} = [\mathbf{x}_1 \ \mathbf{x}_2 \ \cdots \ \mathbf{x}_n]^\top \in \mathbb{R}^{n \times p_n}$ is the design matrix, $\mathbf{y} = [y_1 \ y_2 \ \cdots \ y_n]^\top$, and \mathbf{e} is a mean-zero random vector with $\text{Cov}(\mathbf{e}) = \sigma_2 \mathbf{I}_n$.

We consider the following setup for model selection. Let $\mathcal{A} \subset 2^{[p_n]}$ be a family of index sets representing candidate models. For $\alpha \in \mathcal{A}$, we denote by $p_n(\alpha)$ the cardinality of α and consider the model

$$m_\alpha(\mathbf{X}) = \mathbf{X}_\alpha \boldsymbol{\beta}_\alpha,$$

where \mathbf{X}_α is the sub-matrix of \mathbf{X} containing only the columns indexed by α , and $\boldsymbol{\beta}_\alpha$ is the coefficient vector containing only the entries indexed by α in $\boldsymbol{\beta}$.

1. We say $\alpha \in \mathcal{A}$ is *correct* if $\mathbb{E}[\mathbf{y} \mid \mathbf{X}] \stackrel{\text{a.s.}}{=} m_\alpha(\mathbf{X})$, and we denote by \mathcal{A}_c the set of correct models in \mathcal{A}
2. We say $\alpha \in \mathcal{A}$ is *wrong* if it is not correct, and we denote by \mathcal{A}_w the set of wrong models in \mathcal{A}
3. We say \mathcal{A} is *embedded* if there exists an enumeration $\alpha_1, \alpha_2, \dots, \alpha_k$ of all elements in \mathcal{A} such that

$$\alpha_1 \subset \alpha_2 \subset \cdots \subset \alpha_k.$$

Definition 1.1. For $\alpha \in \mathcal{A}$, let $\hat{\boldsymbol{\beta}}_\alpha$ be the OLS estimator of $\boldsymbol{\beta}_\alpha$ and $\hat{m}_\alpha(\mathbf{X}) := \mathbf{X}_\alpha \hat{\boldsymbol{\beta}}_\alpha$. We denote the average squared error of \hat{m}_α by

$$L_n(\alpha) := \frac{1}{n} \|\mathbb{E}[\mathbf{y} \mid \mathbf{X}] - \hat{m}_\alpha(\mathbf{X})\|^2.$$

Additionally, we write $R_n(\alpha) := \mathbb{E}[L_n(\alpha) \mid \mathbf{X}]$.

The following conditions will be used throughout this section:

- H1 :** $\liminf_{n \rightarrow \infty} \frac{1}{n} \|\mathbf{X}_\alpha \boldsymbol{\beta}\|^2 > 0$ for all $\alpha \in \mathcal{A}$.
- H2 :** $\mathbf{X}^\top \mathbf{X} = O(n)$ and $(\mathbf{X}^\top \mathbf{X})^{-1} = O(n^{-1})$.
- H3 :** $\lim_{n \rightarrow \infty} \max_{i \leq n} h_{ii, \alpha} = 0$ for all $\alpha \in \mathcal{A}$.
- H4 :** $\sum_{\alpha \in \mathcal{A}_w} \frac{1}{(n R_n(\alpha))^m} \rightarrow_{\mathbb{P}} 0$ for some $m \geq 1$.

Proposition 1.1. *If we assume a linear model $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{e}$, then*

$$L_n(\alpha) = \frac{1}{n} \|H_\alpha \mathbf{e}\|^2 + \frac{1}{n} \|M_\alpha \mathbf{X}\boldsymbol{\beta}\|^2 \quad \text{and} \quad R_n(\alpha) = \frac{1}{n} \sigma^2 p_n(\alpha) + \frac{1}{n} \|M_\alpha \mathbf{X}\boldsymbol{\beta}\|^2,$$

where $H_\alpha = \mathbf{X}_\alpha (\mathbf{X}_\alpha^\top \mathbf{X}_\alpha)^{-1} \mathbf{X}_\alpha^\top$ and $M_\alpha = I_n - H_\alpha$.

Proof. First, we have that

$$\begin{aligned} \|\mathbb{E}[\mathbf{y} \mid \mathbf{X}] - \hat{m}_\alpha(\mathbf{X})\|^2 &= \|\mathbf{X}\boldsymbol{\beta} - \mathbf{X}_\alpha \hat{\boldsymbol{\beta}}_\alpha\|^2 \\ &= \|\mathbf{X}\boldsymbol{\beta} - H_\alpha(\mathbf{X}\boldsymbol{\beta} + \mathbf{e})\|^2 \\ &= \|M_\alpha \mathbf{X}\boldsymbol{\beta} - H_\alpha \mathbf{e}\|^2. \end{aligned}$$

Notice that $M_\alpha \mathbf{X}\boldsymbol{\beta}$ and $H_\alpha \mathbf{e}$ are orthogonal:

$$\mathbf{e}^\top H_\alpha M_\alpha \mathbf{X}\boldsymbol{\beta} = \mathbf{e}^\top H_\alpha (I_n - H_\alpha) \mathbf{X}\boldsymbol{\beta} = \mathbf{e}^\top H_\alpha \mathbf{X}\boldsymbol{\beta} - \mathbf{e}^\top H_\alpha \mathbf{X}\boldsymbol{\beta} = 0.$$

Hence, the first part follows from the Pythagorean theorem.

For the second part, we note that $\mathbb{E}[\|H_\alpha \mathbf{e}\|^2 \mid \mathbf{X}] = \sigma^2 p_n(\alpha)$ by the “trace trick”, where $p_n(\alpha)$ denotes the size of model α . \square

Proposition 1.2. *Suppose that the set of correct candidate models $\mathcal{A}_c \subset \mathcal{A}$ is non-empty, and let α^* be the smallest correct model in \mathcal{A}_c . Then, α^* minimizes $R_n(\alpha)$ over $\alpha \in \mathcal{A}$.*

Proof. Let $\alpha \in \mathcal{A}$ be arbitrary and suppose that $\alpha \in \mathcal{A}_c$. Then, $\mathbf{X}_\alpha \boldsymbol{\beta}_\alpha = \mathbf{X}\boldsymbol{\beta}$ and $p_n(\alpha^*) \leq p_n(\alpha)$. Thus,

$$\begin{aligned} R_n(\alpha) &= \frac{1}{n} \sigma^2 p_n(\alpha) + \frac{1}{n} \|M_\alpha \mathbf{X}\boldsymbol{\beta}\|^2 \\ &= \frac{1}{n} \sigma^2 p_n(\alpha) + \frac{1}{n} \underbrace{\|M_\alpha \mathbf{X}_\alpha \boldsymbol{\beta}_\alpha\|^2}_0 \\ &= \frac{1}{n} \sigma^2 p_n(\alpha) \geq \frac{1}{n} \sigma^2 p_n(\alpha^*) = R_n(\alpha^*). \end{aligned}$$

Now suppose that $\alpha \in \mathcal{A}_w$. If $p_n(\alpha) \geq p_n(\alpha^*)$, the result follows by assumption H1. If $p_n(\alpha) < p_n(\alpha^*)$, then ... **MISSING**. \square

1.2 Shao, 1993: Notes on Leave-One-Out CV

In this section, we assume that $p(\alpha) := p_n(\alpha)$ is constant for each $\alpha \in \mathcal{A}$.

Definition 1.2. *The LOOCV estimator of $R_n(\alpha)$ is defined as*

$$\hat{R}_n^{(1)}(\alpha) := \frac{1}{n} \sum_{i=1}^n \left(\frac{y_i - \mathbf{x}_{i\alpha}^\top \hat{\boldsymbol{\beta}}_\alpha}{1 - h_{ii,\alpha}} \right)^2$$

Lemma 1.3 (Shao, 1993).

$$\hat{R}_n^{(1)} (=) \begin{cases} R_n(\alpha) + \sigma^2 + o_{\mathbb{P}}(1) & \text{if } \alpha \in \mathcal{A}_w \\ \frac{1}{n} \|M_\alpha \mathbf{e}\|^2 + \frac{2}{n} \sigma^2 p(\alpha) + o_{\mathbb{P}}(n^{-1}) & \text{if } \alpha \in \mathcal{A}_c \end{cases}$$

Proof. Using the Taylor expansion of $1/(1-x)^2 = 1 + 2x + O(x^2)$, we have

$$\frac{1}{(1 - h_{ii,\alpha})^2} = 1 + 2h_{ii,\alpha} + O_{\mathbb{P}}(h_{ii,\alpha}^2).$$

Thus,

$$\hat{R}_n^{(1)} (=) \underbrace{\frac{1}{n} \sum_{i=1}^n \left(y_i - \mathbf{x}_{i\alpha}^\top \hat{\boldsymbol{\beta}}_\alpha \right)^2}_{\xi_{\alpha,n}} + \underbrace{\frac{1}{n} \sum_{i=1}^n (2h_{ii,\alpha} + O_{\mathbb{P}}(h_{ii,\alpha}^2)) \left(y_i - \mathbf{x}_{i\alpha}^\top \hat{\boldsymbol{\beta}}_\alpha \right)^2}_{\zeta_{\alpha,n}} \quad (1)$$

Let $\xi_{\alpha,n}$ and $\zeta_{\alpha,n}$ denote the first and second terms in (1), respectively. Note that

$$\begin{aligned} \xi_{\alpha,n} &= \frac{1}{n} \|M_\alpha \mathbf{X} \boldsymbol{\beta} + M_\alpha \mathbf{e}\|^2 \\ &= \frac{1}{n} (\|M_\alpha \mathbf{e}\|^2 + \|M_\alpha \mathbf{X} \boldsymbol{\beta}\|^2 + 2\mathbf{e}^\top M_\alpha \mathbf{X} \boldsymbol{\beta}) \end{aligned} \quad (2)$$

$$\begin{aligned} &= \frac{1}{n} \|\mathbf{e}\|^2 + \frac{1}{n} \|M_\alpha \mathbf{X} \boldsymbol{\beta}\|^2 + \frac{1}{n} \|H_\alpha \mathbf{e}\|^2 + \frac{2}{n} \mathbf{e}^\top M_\alpha \mathbf{X} \boldsymbol{\beta} \\ &= \frac{1}{n} \|\mathbf{e}\|^2 + \frac{1}{n} \|M_\alpha \mathbf{X} \boldsymbol{\beta}\|^2 + o_{\mathbb{P}}(1). \end{aligned} \quad (3)$$

The equality at (3) follows from the fact that $\mathbb{E}[\|H_\alpha \mathbf{e}\|^2 \mid \mathbf{X}] = \sigma^2 p(\alpha)$ and

$$\mathbb{E}[\mathbf{e}^\top M_\alpha \mathbf{X} \boldsymbol{\beta} \mid \mathbf{X}]^2 = \sigma^2 \|M_\alpha \mathbf{X} \boldsymbol{\beta}\|^2 = O_{\mathbb{P}}(n), \quad (?)$$

so that $1/n \|H_\alpha \mathbf{e}\|^2 \rightarrow_{\mathbb{P}} 0$ and $2/n (\mathbf{e}^\top M_\alpha \mathbf{X} \boldsymbol{\beta}) = O_{\mathbb{P}}(1)$. (?)

Since $0 < h_{ii,\alpha} < 1$, $2h_{ii,\alpha} + O_{\mathbb{P}}(h_{ii,\alpha}^2) \leq O_{\mathbb{P}}(\max_i h_{ii,\alpha})$. Thus,

$$\zeta_{\alpha,n} \leq O_{\mathbb{P}}\left(\max_i h_{ii,\alpha}\right) \left(\frac{1}{n} \sum_{i=1}^n \left(y_i - \mathbf{x}_{i\alpha}^\top \hat{\boldsymbol{\beta}}_\alpha \right)^2 \right) = O_{\mathbb{P}}\left(\max_i h_{ii,\alpha}\right) \xi_{\alpha,n}. \quad (4)$$

(3) and (4) imply the first case in the Lemma. DOES IT?

If $\alpha \in \mathcal{A}^c$, it is easy to see from (2) that $\xi_{\alpha,n} = 1/n \|M_\alpha \mathbf{e}\|^2$, Furthermore,

$$\zeta_{\alpha,n} = \frac{2}{n} \sigma^2 p(\alpha) + o_{\mathbb{P}}(1), \quad (?)$$

proving the second case. □

Proposition 1.4 (Shao, 1993). Let $\hat{\alpha}^{(1)}$ be the model minimizing $\hat{R}_n^{(1)}(\alpha)$.

1. Under H1, H2, and H3,

$$\lim_{n \rightarrow \infty} \mathbb{P}(\hat{\alpha}^{(1)} \in \mathcal{A}_w) = 0.$$

2. For $\alpha \in \mathcal{A}_c$ with $\alpha \neq \alpha^*$,

$$\mathbb{P}(\hat{R}_n^{(1)}(\alpha) \leq \hat{R}_n^{(1)}(\alpha^*)) = \mathbb{P}(2(p(\alpha) - p(\alpha^*))\sigma^2 < \mathbf{e}^\top (H_\alpha - H_{\alpha^*})\mathbf{e}) + o_{\mathbb{P}}(1).$$

In particular, if $\mathbf{e} \sim \mathcal{N}(0_n, \sigma^2 I_n)$,

$$\mathbb{P}(\hat{R}_n^{(1)}(\alpha) \leq \hat{R}_n^{(1)}(\alpha^*)) = \mathbb{P}(2k < \chi^2(k)) + o_{\mathbb{P}}(1)$$

for $k = p(\alpha) - p(\alpha^*)$.

3. If $p(\alpha^*) \neq p$,

$$\lim_{n \rightarrow \infty} \mathbb{P}(\hat{\alpha}^{(1)} = \alpha^*) \neq 1.$$

Proof.

1. **MISSING**

2. The first part follows from Lemma 2.1 by algebraic manipulation. The second part follows by noting that, if $\mathbf{e} \sim \mathcal{N}(0_n, \sigma^2 I_n)$, then

$$\frac{\mathbf{e}^\top}{\sigma} (H_\alpha - H_{\alpha^*}) \frac{\mathbf{e}}{\sigma} \sim \chi^2(\text{tr}(H_\alpha - H_{\alpha^*})).$$

3. If $p(\alpha^*) = p$, then $\mathcal{A}_c = \{\alpha^*\}$. It follows from 1. that $\lim_{n \rightarrow \infty} \mathbb{P}(\hat{\alpha}^{(1)} = \alpha^*) = 1$. Conversely, if $\lim_{n \rightarrow \infty} \mathbb{P}(\hat{\alpha}^{(1)} = \alpha^*) = 1$ **MISSING** \square

Corollary 1.5. *LOOCV is not consistent. In particular, overfits with non-vanishing probability.*

A subsequent result states that cross-validation is consistent if $n_v/n \rightarrow 1$ as $n \rightarrow \infty$, where n_v is the number of validation samples.

1.3 Shao, 1997

Definition 1.3. Let $\hat{\alpha}_n$ be the model selected by minimizing some criterion \hat{R}_n over \mathcal{A} , and let α_n^* denote the model minimizing R_n over \mathcal{A} . We say \hat{R}_n is consistent if

$$\mathbb{P}\{\hat{\alpha} = \alpha^*\} \rightarrow 1$$

as $n \rightarrow \infty$. We say that \hat{R}_n is asymptotically loss efficient if

$$\frac{R_n(\hat{\alpha})}{R_n(\alpha_n^*)} \rightarrow 1 \quad \text{a.s.}$$

Proposition 1.6 (Shao, 1997). Suppose H1, $p_n/n \rightarrow 0$, and that \mathcal{A}_c is non-empty for all but finitely many n .

1. If $|\mathcal{A}_c| = 1$ for all but finitely many n , then consistency is equivalent to efficiency in the sense of Definition 3.1
2. If $p_n(\alpha_n^*) \not\rightarrow_{\mathbb{P}} \infty$, then consistency is equivalent to efficiency in the sense of Definition 3.1

Proof. MISSING □

Definition 1.4. We define the GIC loss estimator to be

$$\hat{R}_{n,\lambda_n}(\alpha) := \frac{\|\mathbf{y} - \hat{m}(\mathbf{X})\|^2}{n} + \frac{1}{n} \lambda_n \hat{\sigma}_n^2 p_n(\alpha) \quad \text{for } \alpha \in \mathcal{A},$$

where $\hat{\sigma}_n^2$ is an estimator of σ^2 and λ_n is a sequence of positive real numbers satisfying $\lambda_n \geq 2$ and $\lambda_n/n \rightarrow 0$.

1.3.1 The case of $\lambda_n \equiv 2$

Proposition 1.7 (Shao, 1997). Suppose that $\lambda_n = 2$ for all $n \geq 1$ and that $\hat{\sigma}_n^2$ is a consistent estimator of σ^2 . Then,

$$\hat{R}_{n,2}(\alpha) = \left\{ \text{TBD} \right.$$

Theorem 1.8 (Shao, 1997). Suppose that H_4 holds and that $\hat{\sigma}_n^2$ is consistent for σ^2 . Then, $\hat{\alpha}_n^2$ is consistent and asymptotically loss efficient.

1. If $|\mathcal{A}_c| \leq 1$ for all but finitely many n , then $\hat{\alpha}_n^2$ is asymptotically loss efficient.
2. Suppose that $|\mathcal{A}_c| > 1$ for all but finitely many n . If there exists a positive integer m such that $\mathbb{E}[y_1 - \mathbf{x}_1^\top \boldsymbol{\beta}]^{4m} < \infty$ and

$$\sum_{\alpha \in \mathcal{A}_c} \frac{1}{(p_n(\alpha))^m} \rightarrow 0 \quad \text{or} \quad \sum_{\substack{\alpha \in \mathcal{A}_c, \\ \alpha \neq \alpha^*}} \frac{1}{(p_n(\alpha) - p_n(\alpha^*))^m}, \quad (5)$$

then $\hat{\alpha}_n^2$ is asymptotically loss efficient.

3. Suppose that $|\mathcal{A}_c| > 1$ for all but finitely many n and that for any integer q and constant $c > 2$,

$$\liminf_{n \rightarrow \infty} \inf_{Q_n \in \mathcal{Q}_{n,q}} \mathbb{P}\{\mathbf{e}_n^\top Q_n \mathbf{e}_n > c\sigma^2 q\} > 0, \quad (6)$$

where $\mathcal{Q}_{n,q}$ is the set of all projection matrices of rank q . The condition that

$$p_n(\alpha_n^*) \rightarrow \infty \quad \text{or} \quad \min_{\substack{\alpha \in \mathcal{A}_c, \\ \alpha \neq \alpha^*}} (p_n(\alpha) - p_n(\alpha^*)) \rightarrow \infty \quad (7)$$

is necessary and sufficient for the asymptotic loss efficiency of $\hat{\alpha}_n^2$ whenever $|\mathcal{A}_c|$ is bounded or \mathcal{A} is embedded.

Proof. MISSING □

Note that condition (6) is satisfied if $\mathbf{e} \sim \mathcal{N}(0_n, \sigma^2 I_n)$. Condition (7) is satisfied if \mathcal{A} does not contain two correct models with fixed dimensions for all but finitely many n .

Corollary 1.9 (Shao, 1997). *If \mathcal{A}_c contains exactly one model with fixed dimension for all but finitely many n , then $\hat{\alpha}_n^2$ is consistent.*

Proof. This follows immediately from Theorem 1.8 and Proposition 1.7. \square

INCOMPLETE: Missing $\lambda_n \rightarrow \infty$ and cross-validation discussion.

2 Nonlinear Model Selection

2.1 Yang, 2007

Here we consider two regression procedures, denoted δ_1 and δ_2 , that yield estimators $\hat{f}_{n,1}$ and $\hat{f}_{n,2}$ of the regression function satisfying

$$y_i = f(\mathbf{x}_i) + \epsilon_i \quad i \in [n],$$

for \mathbf{x}_i iid, $\mathbb{E}[\epsilon_i \mid \mathbf{X}] \stackrel{\text{a.s.}}{=} 0$ and $\mathbb{E}[\epsilon_i^2 \mid \mathbf{X}] \stackrel{\text{a.s.}}{<} \infty$.

Definition 2.1. *We say δ_1 is asymptotically better than δ_2 under the loss function L if, for $0 < \varepsilon < 1$, there exists $c_\varepsilon > 0$ such that*

$$\mathbb{P} \left\{ L_n(\hat{f}_{n,2}) \geq (1 + c_\varepsilon L_n(\hat{f}_{n,1})) \right\} \geq 1 - \varepsilon.$$

Given that δ_1 is asymptotically better than δ_2 , we say that a selection procedure is consistent if it selects δ_1 with probability tending to 1 as $n \rightarrow \infty$.

2.1.1 Single-split cross-validation (the Hold-out)

For this section, we assume that the first n_1 elements in \mathcal{D}_n are used as a training/estimation sample and the remaining n_2 elements make up the validation sample. We write p_n and q_n for the rates of convergence of the estimators $\hat{f}_{n,1}$ and $\hat{f}_{n,2}$, respectively. That is,

$$O_{\mathbb{P}}(p_n) = \left\| f - \hat{f}_{n,1} \right\|_2 \quad \text{and} \quad O_{\mathbb{P}}(q_n) = \left\| f - \hat{f}_{n,2} \right\|_2.$$

The hold-out cross-validation method consists in selecting the estimator that minimizes the hold-out loss

$$L_{\text{ho}}(\hat{f}_{n,j}) = \sum_{i=n_1+1}^n \left(y_i - \hat{f}_{n,j}(\mathbf{x}_i) \right)^2 \quad \text{for } j = 1, 2.$$

The propositions in this section rely on the following conditions:

- **C0:** $\mathbb{E}[\epsilon_i^2 \mid \mathbf{x}_i]$ is bounded a.s. for $i \in [n]$.
- **C1:** There exists A_n such that $\left\| f - \hat{f}_{n,j} \right\|_\infty = O_{\mathbb{P}}(A_n)$ for $j = 1, 2$.

- **C2:** One procedure is asymptotically better than the other.
- **C3:** There exists M_n such that $\left\|f - \hat{f}_{n,j}\right\|_4 / \left\|f - \hat{f}_{n,j}\right\|_4 = o_{\mathbb{P}}(M_n)$ for $j = 1, 2$.

Theorem 2.1 (Yang, 2007). *Suppose that **C0–C3** hold. Suppose, furthermore, that*

1. $n_1 \rightarrow \infty$
2. $n_2 \rightarrow \infty$
3. $n_2 M_n^{-4} \rightarrow \infty$
4. $\sqrt{n_2} \max(p_{n_1}, q_{n_1})$

Then, the hold-out CV procedure is consistent.

A very detailed proof of this result is provided in Yang [1], so it will be skipped here.

2.1.2 Voting cross-validation with multiple splits

The (theoretical) majority-vote cross-validation method proceeds as follows: for each permutation $i \mapsto \pi(i)$ of the data, we compute the estimators $\hat{f}_{n_1,1}$ and $\hat{f}_{n_1,2}$ using the first n_1 data points $(y_{\pi(1)}, \mathbf{x}_{\pi(1)}), \dots, (y_{\pi(n_1)}, \mathbf{x}_{\pi(n_1)})$ as the training sample and the remaining $n_2 = n - n_1$ elements as the validation sample. We then find the estimator that minimizes the hold-out loss

$$L_{\pi}(\hat{f}_{n_1,j}) = \sum_{i=n_1+1}^n \left(y_{\pi(i)} - \hat{f}_{n_1,j}(\mathbf{x}_{\pi(i)}) \right)^2 \quad \text{for } j = 1, 2.$$

The chosen estimator is the one favored by the majority of the permutations. More formally, we define

$$\tau_{\pi} = \mathbb{1}_{L_{\pi}(\hat{f}_{n_1,1}) \leq L_{\pi}(\hat{f}_{n_1,2})}$$

We then define our selection criterion as follows:

$$\hat{f}_n = \begin{cases} \hat{f}_{n,1} & \text{if } \sum_{\pi \in \Pi} \tau_{\pi} \geq n!/2, \\ \hat{f}_{n,2} & \text{otherwise,} \end{cases}$$

where Π denotes the set of all permutations of $[n]$.

Theorem 2.2 (Yang, 2007). *Under the conditions of Theorem 4.1 and the condition that the data is iid, the majority-vote cross-validation method is consistent.*

Proof. Suppose that δ_1 is asymptotically better than δ_2 . For $\pi \in \Pi$, we have that

$$\mathbb{P} \left\{ L_{\pi}(\hat{f}_{n_1,1}) \leq L_{\pi}(\hat{f}_{n_1,2}) \right\} = \mathbb{E}[\tau_{\pi}] \stackrel{(*)}{=} \mathbb{E} \left[\frac{1}{n!} \sum_{\pi \in \Pi} \tau_{\pi} \right].$$

The equality at $(*)$ follows from the fact that the data are iid, hence exchangeable, and thus the τ_{π} are identically distributed. By Theorem 2.1, the right-hand side converges to 1 as $n \rightarrow \infty$. Since the average $1/n! \sum_{\pi} \tau_{\pi}$ is almost surely at most 1, it follows that $1/n! \sum_{\pi} \tau_{\pi} \rightarrow 1$ in probability, and the majority-vote cross-validation method is consistent. \square

Remark: The proof of Theorem 2.2 does not rely on using the whole of Π for the majority vote. Indeed, Theorem 2.1 shows that even a single split is enough. What we gain by considering Π (or a subset of it) is more robustness to the choice of the training sample.

References

- [1] Yuhong Yang. “Consistency of cross validation for comparing regression procedures”.
In: *The Annals of Statistics* 35.6 (Dec. 2007).