

## Mathematical Methods II

### Weekly problem set 6

- (1) Consider the 2D wave equation

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left[ \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right]$$

where  $k$  is a real constant. Show that it can be separated into two independent ODEs and form a general solution to the PDE.

**Solution** Sub in the derivatives

$$XYT'' = c^2 [X''YT + XY''T]$$

Separate by dividing by  $c^2XYT$

$$\frac{X''}{X} + \frac{Y''}{Y} = \frac{1}{c^2} \frac{T''}{T} = \mu$$

Let's say

$$l + m = \mu$$

where

$$\frac{X''}{X} = l \qquad \frac{Y''}{Y} = m \qquad \frac{T''}{T} = c^2 \mu$$

Each of these ODEs can be solved separately giving

$$X = c_1 e^{\sqrt{l}x} + c_2 e^{-\sqrt{l}x} \quad Y = c_3 e^{\sqrt{m}y} + c_4 e^{-\sqrt{m}y} \quad T = c_5 e^{c\sqrt{\mu}t} + c_6 e^{-c\sqrt{\mu}t}$$

Hence the general solution is

$$u(x, y, t) = XYT = \left( c_1 e^{\sqrt{l}x} + c_2 e^{-\sqrt{l}x} \right) \left( c_3 e^{\sqrt{m}y} + c_4 e^{-\sqrt{m}y} \right) \left( c_5 e^{c\sqrt{\mu}t} + c_6 e^{-c\sqrt{\mu}t} \right)$$

- (2) Solve the 1D heat equation for a long, thin metal rod of length  $L$

$$\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$$

using the boundary conditions for insulated end points,  $u_x(0, t) = 0$ ,  $u_x(L, t) = 0$ ,  $t > 0$  and the initial condition  $u(x, 0) = x$  for  $0 < x < L$ .

**Solution** Stage 1: Obtain two ODE's. Start by subbing in the  $u=XT$  derivatives

$$XT' = \alpha^2 X''T.$$

Separate the variables

$$\frac{X''}{X} = \frac{1}{\alpha^2} \frac{T'}{T} = \mu.$$

$$\begin{array}{ll} \frac{X''}{X} = \mu & \frac{1}{\alpha^2} \frac{T'}{T} = \mu \\ X'' - \mu X = 0 & T' - \alpha^2 \mu T = 0 \end{array}$$

Stage 2: Consider the boundary conditions on  $u_x$  to restrict  $X'$  and  $T'$

$$u_x(0, t) = 0: X'(0)T(t) = 0 \Rightarrow X'(0) = 0$$

$$u_x(L, t) = 0: X(L)T(t) = 0 \Rightarrow X'(L) = 0$$

Stage 3: Consider the 3 cases of  $\mu$

(1)  $\mu = 0$ : Beginning with the equation for  $X$

$$X'' = 0$$

Examining the roots or by inspection we realise that the solution to this equation is

$$X(x) = ax + b.$$

Differentiate  $X$  to obtain  $X'$

$$X'(x) = a.$$

Now apply the boundary conditions for  $X$

$$X'(0) = 0 \Rightarrow a = 0 \Rightarrow X'(x) = 0,$$

$$X'(L) = 0 \Rightarrow \text{No effect on } b. \Rightarrow X(x) = b.$$

Since we have a non-trivial solution for  $X$ , it is worth looking at  $T$ . Following the same process, applying  $\mu = 0$  the ODE for  $T$  becomes

$$T' = 0$$

Which has a solution

$$T = c$$

So the solution for  $u$  is

$$u(x, t) = bc = d$$

(2)  $\mu > 0$ : Let's say that  $\mu = r^2$  so we can be sure it is a positive constant and avoid fractional powers in our solutions. Beginning with the equation for  $X$

$$X'' - r^2 X = 0$$

Examining the roots we realise that this equation has the solution

$$X(x) = ae^{rx} + be^{-rx}$$

Differentiating

$$X'(x) = rae^{rx} - rbe^{-rx}$$

Now apply the boundary conditions for  $X'$

$$X'(0) = 0 \Rightarrow r(a - b) = 0 \Rightarrow a = b \text{ since } r \neq 0,$$

Using our second BC

$$X'(L) = 0 \Rightarrow ra(e^{rL} - e^{-rL}) = 0 \Rightarrow a = 0 \Rightarrow X(x) = 0.$$

Therefore when  $\mu > 0$ ,  $X(x) = 0$  for any given  $x$  and hence  $u(x, t) = 0$  for any given  $x$ , so this solution is of no interest.

- (3)  $\mu < 0$ : Let's say that  $\mu = -r^2$  so we can be sure it is a negative constant and avoid fractional powers in our solutions. Beginning with the equation for  $X$

$$X'' + r^2X = 0$$

Examining the roots we realise that this equation has the solution

$$X(x) = a \cos rx + b \sin rx$$

Differentiating

$$X'(x) = -ra \sin rx + rb \cos rx$$

Now apply the boundary conditions for  $X'$

$$X'(0) = 0 \Rightarrow rb = 0 \Rightarrow b = 0 \Rightarrow X'(x) = -ra \sin rx,$$

$$X'(L) = 0 \Rightarrow -ra \sin rL = 0 \Rightarrow rL = n\pi.$$

So  $r = n\pi/L$  for  $n = 1, 2, 3, \dots$ , giving

$$X_n(x) = A_n \cos \frac{n\pi x}{L}$$

Since we have a non-trivial solution for  $X$ , it is worth looking at  $T$ . Following the same process, applying  $\mu = 0$  the ODE for  $T$  becomes

$$T' + \alpha r^2 T = 0.$$

This ODE has the solution

$$T(t) = ce^{-\alpha^2 r^2 t}$$

We have already established that  $r = n\pi/L$ , so subbing this in the equation becomes

$$T_n(t) = c_n e^{-\frac{\alpha^2 n^2 \pi^2 t}{L^2}}$$

And so, given that  $u = XT$ , our solution for  $u(x, t)$  is

$$u(x, t) = A_n \cos \frac{n\pi x}{L} e^{-\frac{\alpha^2 n^2 \pi^2 t}{L^2}}$$

where  $A_n = a_n c_n$ .

Stage 4: Sum the 3 cases of  $\mu$

$$u(x, t) = d + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L} e^{-\frac{\alpha^2 n^2 \pi^2 t}{L^2}}$$

Apply the initial condition

$$u(x, 0) = x = d + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L}$$

Use the Fourier formulae to determine the coefficients

$$\begin{aligned} D &= \frac{1}{L} \int_0^L f(x) dx = \frac{1}{L} \int_0^L x dx = \frac{1}{L} \frac{L^2}{2} = \frac{L}{2} \\ A_n &= \frac{2}{L} \int_0^L f(x) \cos \frac{\pi n x}{L} dx = \frac{2}{L} \int_0^L x \cos \frac{\pi n x}{L} dx \\ &= \frac{2L}{\pi^2} \int_0^L \left[ \frac{\cos \pi n}{n^2} - \frac{1}{n^2} \right] = \frac{2L}{\pi^2} \int_0^L \left[ \cos \frac{(-1)^n - 1}{n^2} \right] \end{aligned}$$

Thus the final solution is

$$u(x, t) = \frac{L}{2} + \frac{2L}{\pi^2} \int_0^L \left[ \cos \frac{(-1)^n - 1}{n^2} \right] \cos \frac{n\pi x}{L} e^{-\frac{\alpha^2 n^2 \pi^2 t}{L^2}}$$