



ALICE-India Collaboration



Quarkonium polarization in pp and Pb-Pb collisions with ALICE

Dushmanta Sahu
Indian Institute of Technology Indore, India
April 29, 2023

Introduction:




➤ Why charmonia?

- Charm and anti-charm quarks produced early in the system's evolution : during the pre-equilibrium phase
- Affected by suppression and regeneration at LHC energies
- J/ψ remains largely undiffused in the hadronic phase of a collision which makes it a better probe to study the deconfined phase
- Charmonium studies in hadronic collisions provide powerful tests of quantum chromodynamics (QCD)
- Charmonium production yield in Pb—Pb and p—Pb collisions can also be affected by the cold nuclear matter (CNM) effect (e.g. Shadowing effect)

➤ Polarization in pp collisions:

- Polarization is the measure of how much the spin of a particle is aligned in a given direction
- Gluon's polarization is preserved as the $c\bar{c}$ pair evolves into a bound state of charmonium
- In two-body decays, the spin-alignment will be reflected in the angular distribution of the decay particles

Introduction:



- $h = \frac{\mathbf{S} \cdot \mathbf{p}}{|\mathbf{p}|} \rightarrow$ Helicity operator
- Vector ($J^{PC} = 1^{--}$) quarkonia have the same charge-parity as an electron-positron pair and can be produced in electron-positron annihilation via an intermediate photon
- The states originating from this process are polarized, as a consequence of helicity conservation, a general property of QED (QCD) in the relativistic (massless) limit
- For our case, gluon fragmentation dominates the high p_T region, while Drell-Yan process dominates the low p_T region


Introduction:

J/ψ polarization puzzle ?


- Measurements of polarization parameters from Tevatron, RHIC and LHC show almost no J/ψ polarization in hadronic collisions
- However, theoretical predictions based on the collinear factorized color singlet production channel at leading order (LO) and next-to-leading order (NLO) suggested substantially non-zero polarization at high p_T
- Conflicting theoretical results from non-relativistic quantum chromodynamics (NRQCD) and Color Singlet Model

Importance of $\psi(2S)$ polarization study :

- A small prompt J/ψ polarization can be interpreted as reflecting a mixture of directly produced mesons with those produced in the decays of heavier (P-wave) charmonium states
- $\psi(2S)$ is unaffected by feed-down decays from heavier charmonia



[Phys. Rev. Lett. 108, 172002 (2012)]






[ALICE Collaboration, Phys. Rev. Lett. 108, 082001 (2012)]

Introduction:

Importance of $\Upsilon(nS)$ polarization study :

- $b\bar{b}$ system satisfies the non relativistic calculations at high p_T much better than the $c\bar{c}$
- Better probe for QCD
- Results from Tevatron show almost no (CDF) or longitudinal polarization for $\Upsilon(1S)$ (D0)
- At lower energy and p_T , the E866 experiment has shown yet a different polarization pattern: the $\Upsilon(2S)$ and $\Upsilon(3S)$ states have maximal transverse polarization
- Unexpectedly, the $\Upsilon(1S)$ found to be only weakly polarized



[NuSea Collaboration, Phys. Rev. Lett. 86, 2529 (2001)]

[CDF Collaboration, Phys. Rev. Lett. 88, 161802 (2002)]

[DΦ Collaboration, Phys. Rev. Lett. 101, 182004 (2008)]


Introduction:

- The angular distribution in dilepton decay:

$$\frac{d^2 N}{d\cos\theta d\phi} = \frac{3}{4\pi(3 + \lambda_\theta)} (1 + \lambda_\theta \cos^2\theta + \lambda_\phi \sin^2\theta \cos 2\phi + \lambda_{\theta\phi} \sin 2\theta \cos\phi)$$

[P.Faccioli, et. al., Eur. Phys. J. C 69, 657 (2010)]


- | | | |
|---|---|---------------------------|
| $(\lambda_\theta, \lambda_\phi, \lambda_{\theta\phi}) = (1, 0, 0)$ | → | Transverse polarization |
| $(\lambda_\theta, \lambda_\phi, \lambda_{\theta\phi}) = (-1, 0, 0)$ | → | Longitudinal polarization |
| $(\lambda_\theta, \lambda_\phi, \lambda_{\theta\phi}) = (0, 0, 0)$ | → | Unpolarized state |



Introduction:

Frames of reference


- The helicity frame uses the $\psi(2S)$ momentum as the quantization axis
- In the Collins—Soper frame, the quantization axis is chosen to be the bisector of the angle between the two incoming beams in the rest frame of the $\psi(2S)$ meson
- We can define the frame-invariant variable λ_{inv}



$$\lambda_{inv} = \frac{\lambda_\theta + 3\lambda_\phi}{1 - \lambda_\phi}$$

Quarkonium polarization in Pb–Pb collisions:

- Large non-zero magnetic field in non-central heavy-ion collisions
- Production of vorticity due to large initial angular momentum
- Both the external magnetic field and the initial angular momentum produced in the non-central heavy-ion collisions may influence the quarkonium polarization
- Event Plane (EP) frame: direction of the polarization axis orthogonal to the event plane in the centre-of-mass of the colliding beams
- The studies in Collins–Soper and Helicity frames are also interesting in AA to study quarkonium suppression/regeneration in the QGP



Magnetic field (\vec{B}):

- Huge intensity (10^{14} T)
- Short lived ($\tau = 1 \text{ fm}/c$)


[Kharzeev et al., NPA 803 (2008)]

Angular momentum (\vec{L}):

- Largest in semicentral collisions
- Can affect the system evolution till freeze-out

[Becattini et al., PRC 77 (2008) 024906]


ALICE detector (Run 2):



- Inclusive quarkonium measurements performed at forward rapidity in the dimuon decay channel

Muon spectrometer acceptance: $-4.0 < \eta < -2.5$


[ALICE Muon spectrometer]



New measurements from Run 2 datasets

- $pp : \sqrt{s} = 13 \text{ TeV}$
- $Pb-Pb : \sqrt{s_{NN}} = 5.02 \text{ TeV}$

Quarkonium polarization in Pb-Pb collisions:




- ALICE measurement of J/ψ polarization in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in Helicity (HE) and Collins-Soper (CS) reference frames
- λ_θ shows a 2σ deviation from zero at low p_T
- 3σ deviation from LHCb measurement in pp collisions in the Helicity frame
- Values compatible with ALICE results in pp collisions within uncertainties

ALICE Collaboration, Phys. Lett. B 815, 136146 (2021)


ALICE Collaboration, Eur. Phys. J. C 78, 562 (2018)

LHCb Collaboration, Eur. Phys. J. C 73, 2631 (2013)

Quarkonium polarization in Pb-Pb collisions:



ALI-PUB-521052




ALI-PUB-521057

- ALICE measurement of J/ψ polarization in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV
- First measurement with respect to the Event Plane (EP)
- Small but significant polarisation (3.5σ), particularly in the 40-60% centrality range
- Effect more pronounced at low transverse momentum ($2 < p_T < 4$ GeV/c) in centrality 30-50%
- Qualitatively in agreement with spin alignment observed for light vector mesons [Phys. Rev. Lett. 125, 012301 (2022)]

[ALICE Collaboration, Phys. Rev. Lett. 131, 042303 (2023)]

Quarkonium polarization in pp collisions:




- J/ψ polarization measured in pp collisions in the CS and HE frames
 - Dataset : ALICE $\sqrt{s} = 7$ TeV (2010)
ALICE $\sqrt{s} = 8$ TeV (2012)
LHCb $\sqrt{s} = 7$ TeV (2011)
 - No significant polarisation observed by ALICE and LHCb at forward rapidity
 - Need for studies with higher center of mass energies
- ✓ New ongoing analyses of J/ψ and $\psi(2S)$ in pp collisions at $\sqrt{s} = 13$ TeV

ALICE Collaboration, Phys. Rev. Lett. 108, 082001 (2012)

ALICE Collaboration, Eur. Phys. J. C 78, 562 (2018)

LHCb Collaboration, Eur. Phys. J. C 73, 2631 (2013)

Quarkonium polarization in pp collisions:




- Recent preliminary measurement of $Y(1S)$ polarization at $\sqrt{s} = 13$ TeV from ALICE
- Results compatible with previous LHCb measurements at $\sqrt{s} = 8$ TeV
- Polarization is evaluated down to $p_T \sim 0$
- All values compatible with zero within uncertainties
- Large uncertainties due to limited statistical precision

LHCb Collaboration, JHEP 12, 110 (2017)


ALI-PREL-505590

Quarkonium polarization in pp collisions:



Analysis Note: $\psi(2S)$ polarization measurement in pp collisions at $\sqrt{s} = 13$ TeV, <https://alice-notes.web.cern.ch/node/1472> (ALICE internal)

- Zero polarization within uncertainties for Collins-Soper frame
- Finite polarization in helicity frame for $4 < p_T < 8$ GeV/c



Conclusion and Outlook:

- ALICE has measured the polarization of several quarkonium states both in pp and Pb—Pb collisions
- No significant quarkonium polarization till now in pp collisions
- New J/ψ and $\psi(2S)$ polarization analyses ongoing in pp collision at $\sqrt{s} = 13$ TeV
- Results are compatible with other LHC measurements and recent model predictions
- Hint for non-zero polarization at low p_T in the HE and CS frames in Pb—Pb collisions
- From the results of EP frame analysis, possible correlation with \vec{B} and \vec{L} in the QGP formed in heavy-ion collision
- ALICE Run 3 with high luminosity will provide significant statistics for precision measurements

THANK YOU!