

ADD

Movie stars

Estimating epidemic speed and strength

The role of generation intervals

Jonathan Dushoff, McMaster University

U. Glasgow, Oct 2019

Outline

Overview

Capacity building

Compartmental models

The $r\mathcal{R}$ relationship

Generation intervals


Generations through time

Other kinds of generation interval


Speed and strength

Problem (present)


- ▶ I am fundamentally a math person ...



What is the pattern of Pythagorean triples of integers
 $a^2 + b^2 = c^2$?



Divide a square and a circle each into two complementary subsets
that are pairwise similar



How many at-bats does it take to get a given batting average?

Problem

- ▶ I am fundamentally a math person
 - ▶ but I want to do work that is relevant to people


Solution

- ▶ Dynamical modeling is fun and useful

Dynamical modeling connects scales




Measles reports from England and Wales



- ▶ Start with rules about how things change in short time steps
 - ▶ Usually based on *individuals*
- ▶ Calculate results over longer time periods
 - ▶ Usually about *populations*

Fun and useful!




New problem

- ▶ There is (or was a gulf) between dynamical and statistical modeling
 - ▶ Dynamics are needed to incorporate mechanism
 - ▶ Statistics are needed to incorporate uncertainty

Ebola forecasting

Sierra Leone



Outline

Overview

Capacity building

Compartmental models

The $r\mathcal{R}$ relationship

Generation intervals

Generations through time

Other kinds of generation interval

Speed and strength

- ▶ International Clinics on Infectious Disease Dynamics and Data
- ▶ ici3d.org
- ▶ At least two clinics a year, usually in South Africa
- ▶ MMED every year at the African Institute of Mathematical Sciences in Muizenberg
- ▶ Aimed at engaging dynamical models with infectious disease data

Prior distribution

- Priors are a unique feature of the Bayesian approach
 - May contain subjective information
 - Sometimes without empirical information
- Beta-prior distribution of shape parameters
 - Has prior properties from which it can be derived
 - Sharp prior distributions are very informative

Sasha
Jan = Jan
cold
2
absolute
J

Outline

Overview

Capacity building

Compartmental models

The $r\mathcal{R}$ relationship

Generation intervals


Generations through time

Other kinds of generation interval

Speed and strength


Compartmental models

Divide people into categories:




- ▶ Susceptible → Infectious → Recovered
- ▶ Individuals recover independently
- ▶ Individuals are infected by infectious people

Differential equation implementation



Individual-based implementation




Lessons

- ▶ Tendency to oscillate
- ▶ Thresholds
- ▶ Exponential growth


Big \mathcal{R}

- ▶ \mathcal{R} is the number of people who would be infected by an infectious individual *in a fully susceptible population*.
- ▶ $\mathcal{R} = \beta/\gamma = \beta D = (cp)D$
 - ▶ c : Contact Rate
 - ▶ p : Probability of transmission (infectivity)
 - ▶ D : Average duration of infection
- ▶ A disease can invade a population if and only if $\mathcal{R} > 1$.
- ▶ Often focus on initial period (may also say \mathcal{R}_0)

Big \mathcal{R}




Homogeneous endemic curve



- ▶ Threshold value
- ▶ Sharp response to changes in factors underlying transmission
- ▶ Works – sometimes
 - ▶ Sometimes predicts unrealistic sensitivity

Yellow fever in Panama



Exponential growth


- ▶ Diseases have a tendency to grow exponentially at first
 - ▶ I infect three people, they each infect 3 people ...
 - ▶ How fast does disease grow?
 - ▶ How quickly do we need to respond?

little r

- ▶ We measure epidemic *speed* using little r :
 - ▶ *Units:* [1/time]
 - ▶ Disease increases like e^{rt}
- ▶ Time scale is $C = 1/r$
 - ▶ Ebola, $C \approx 1\text{month}$
 - ▶ HIV in SSA, $C \approx 18\text{month}$
- ▶ Often focus on initial period (may also say r_0)

little *r*

$$R_0 = 5.66$$



Limitations

- ▶ Many conclusions from this framework make strong assumptions:
 - ▶ **Spatial homogeneity:** everywhere is the same
 - ▶ **Individual homogeneity:** everyone is the same
 - ▶ and everyone is everywhere
 - ▶ **Temporal homogeneity:**
 - ▶ It doesn't matter how long I've been infected, I'm either infected or not

Outline

Overview

Capacity building

Compartmental models

The rR relationship

Generation intervals

Generations through time

Other kinds of generation interval

Speed and strength

The $r\mathcal{R}$ relationship


- ▶ We're very interested in the relationship between little r and \mathcal{R} .
- ▶ We might have good estimates of r only
 - ▶ e.g., West African Ebola outbreak, HIV in Africa
- ▶ Or we might have good estimates of \mathcal{R} only
 - ▶ Measles, influenza

Example: Post-death transmission and safe burial


- ▶ How much Ebola spread occurs before vs. after death
- ▶ Highly context dependent
 - ▶ Funeral practices, disease knowledge
- ▶ *Weitz and Dushoff Scientific Reports 5:8751.*



Standard disease model




Add a latent period




- ▶ (i.e., a lag between infection and infectiousness)

Add post-death transmission




Add post-death transmission



What happens if we account for burial transmission?

- ▶ We've made the disease transmitting process slower, so obviously Ebola is *less* dangerous than we thought
- ▶ We've added another source of transmission, so obviously Ebola is *more* dangerous than we thought
- ▶ What we learn depends on what we know!

What do we know?



Outline

Overview

Capacity building

Compartmental models

The rR relationship

Generation intervals

Generations through time


Other kinds of generation interval

Speed and strength


Generation intervals

- ▶ The generation distribution measures the time between generations of the disease
 - ▶ Interval between “index” infection and resulting infection
- ▶ Generation intervals provide the link between \mathcal{R} and r


Approximate generation intervals





Generations and \mathcal{R}



Generations and \mathcal{R}



Generations and \mathcal{R}



Conditional effect of generation time

- ▶ Given the reproductive number \mathcal{R}
 - ▶ faster generation time G means higher r
 - ▶ More danger
- ▶ Given r
 - ▶ faster generation time G means smaller \mathcal{R}
 - ▶ Less danger

Linking framework

- ▶ Epidemic speed (r) is a *product*:
 - ▶ (something to do with) generation speed
 - ▶ \times (something to do with) epidemic strength
- ▶ Epidemic strength is therefore (approximately) a *quotient*
 - ▶ Epidemic speed
 - ▶ \div (something to do with) generation speed

Filtered means

- ▶ There is a sensible way to define an “effective” generation time
- ▶ Preserve the exponential growth equation

$$\mathcal{R} = \exp(r\hat{G})$$

- ▶ \hat{G} is a “filtered mean” of the distribution g :





$$\exp(-r\hat{G}) = \langle \exp(-r\tau) \rangle_g.$$

,

- ▶ This is cool, but not easy to interpret (our estimates about the generation time change when r changes)



Approximations

Approximate generation intervals





Moment approximation

Approximate generation intervals





Moment approximation



Moment approximation

Approximate generation intervals



Compound-interest interpretation

- ▶ Define $\mathcal{R} \approx (1 + r\kappa\bar{G})^{1/\kappa} \equiv X(r\bar{G}; 1/\kappa)$
- ▶ X is the compound-interest approximation to the exponential
 - ▶ Linear when $\kappa = 1$ (i.e., when g is exponential)
 - ▶ Approaches exponential as $\kappa \rightarrow 0$
- ▶ Key quantity is $r\bar{G}$: the relative length of the generation interval compared to the characteristic time scale of spread

Qualitative response

- ▶ For a given value of \bar{G} , smaller values of κ mean:
 - ▶ less variation in generation interval
 - ▶ less compounding of growth
 - ▶ greater \mathcal{R} required for a given r



Intuition

- ▶ Longer generation times mean less speed
 - ▶ \implies more strength, when speed is fixed
- ▶ What about more variation?
 - ▶ More action (both before and after the mean time)
 - ▶ But what happens early is more important in a growing system
- ▶ More variation means more speed
 - ▶ \implies less strength, when speed is fixed


How well do approximations work

- ▶ Simulate realistic generation intervals for various diseases
- ▶ Compare approximate $r\mathcal{R}$ relationship with known exact relationship
 - ▶ Known because we are testing ourselves with simulated data


Ebola distribution




Ebola curve



Measles curve



Rabies curve



Generation intervals



"I cannot imagine how they will spend half of it."

- ▶ Sort of the poor relations of disease-modeling world
- ▶ Ad hoc methods
- ▶ Error often not propagated

Summary

- ▶ Generation intervals are the missing link between r and \mathcal{R}
- ▶ We need better methods for estimating them, and propagating uncertainty to other parts of the model
- ▶ Filtered means may help with intuition
- ▶ For many practical applications:
 - ▶ Estimating the mean generation interval is not enough
 - ▶ But estimating the mean and CV may be enough
 - ▶ A good basis for understanding and propagating uncertainty

Outline

Overview

Capacity building

Compartmental models

The $r\mathcal{R}$ relationship

Generation intervals

Generations through time

Other kinds of generation interval

Speed and strength

Generations through time

- ▶ Generation intervals can be estimated by:
 - ▶ Observing patients:
 - ▶ How long does it take to become infectious?
 - ▶ How long does it take to recover?
 - ▶ What is the time profile of infectiousness/activity?
 - ▶ Contact tracing
 - ▶ Who (probably) infected whom?
 - ▶ When did each become infected?
 - ▶ — or ill (serial interval)?

Which is the real interval?



- ▶ Contact-tracing intervals look systematically different, depending on when you observe them.
- ▶ Observed in:
 - ▶ Real data, detailed simulations, simple model
- ▶ Also differ from intrinsic (infector centered) estimates

Types of interval

- ▶ Define:
 - ▶ *Intrinsic interval*: How infectious is a patient at time τ after infection?
 - ▶ *Forward interval*: When will the people infected today infect others?
 - ▶ *Backward interval*: When did the people who infected people today themselves become infected?
 - ▶ *Censored interval*: What do all the intervals observed up until a particular time look like?
 - ▶ Like backward intervals, if it's early in the epidemic

Growing epidemics


- ▶ Generation intervals look *shorter* at the beginning of an epidemic
 - ▶ A disproportionate number of people are infectious right now
 - ▶ They haven't finished all of their transmitting
 - ▶ We are biased towards observing faster events



What changes backward intervals?

- ▶ Who is likely to infect me depends on:
 - ▶ How infectious they are (intrinsic GI)
 - ▶ How many of them there are (changes in disease incidence)

Backward intervals




Champredon and Dushoff, 2015. DOI:10.1098/rspb.2015.2026

What changes forward intervals?

- ▶ Who I am likely to infect depends on:
 - ▶ How infectious I am (intrinsic GI)
 - ▶ How many of them there are (changes in numbers of susceptibles)

Forward intervals



Champredon and Dushoff, 2015. DOI:10.1098/rspb.2015.2026

Conclusion

- ▶ Backward intervals change if the number of infectious individuals is changing as you look back
- ▶ Forward intervals change if the number of *susceptible* individuals is changing as you look forward
- ▶ Lack of care in defining generation intervals can lead to bias
 - ▶ In particular, censored intervals look too short, lead to underestimates of \mathcal{R} .

Outline

Overview

Capacity building

Compartmental models

The $r\mathcal{R}$ relationship

Generation intervals

Generations through time

Other kinds of generation interval

Speed and strength

Other kinds of generation interval




- ▶ Once you think carefully about generation intervals, they're everywhere
- ▶ Spatial heterogeneity
- ▶ Individual heterogeneity

Generations in space

- ▶ How do local interactions affect realized generation intervals?




Individual




Surprising results

- ▶ \mathcal{R} on networks generally *smaller* than values estimated using r .
 - ▶ Trapman et al., 2016. *JRS Interface*
DOI:10.1098/rsif.2016.0288
- ▶ Because people don't question the intrinsic generation interval
 - ▶ Local interactions
 - ▶ \implies wasted contacts
 - ▶ \implies shorter generation intervals
 - ▶ \implies smaller estimates of \mathcal{R} .

Observed and estimated intervals



Outbreak estimation




Serial intervals

- ▶ Do serial intervals and generation intervals have the same distribution?
- ▶ It seems that they should: they describe generations of the same process
- ▶ In fact, they don't
 - ▶ Serial intervals can even be negative!
 - ▶ You might report to the clinic with flu before me, even though I infected you

Single Transmission



Multiple Transmissions



○ Getting Bitten

● Become Infectious

★ Clinical Signs

● Contact/Biting

··· Incubation Period

— Infectious Period

— Waiting Time

X Removed


Rabies

- ▶ If symptoms always start *before* infectiousness happens, then serial interval should equal generation interval:
 - ▶ incubation time + extra latent time + waiting time
 - ▶ extra latent time + waiting time + incubation time

Single Transmission



Multiple Transmissions



○ Getting Bitten

● Become Infectious

★ Clinical Signs

● Contact/Biting


··· Incubation Period

— Infectious Period

— Waiting Time

X Removed

Repeated biter incubation period



Outline

Overview

Capacity building

Compartmental models

The $r\mathcal{R}$ relationship

Generation intervals

Generations through time

Other kinds of generation interval

Speed and strength



Can treatment stop the HIV epidemic?

- ▶ Modern treatments are well tolerated and highly effective
- ▶ Virus is undetectable, and transmission is negligible
- ▶ Can active testing and treatment stop the epidemic?




Are HIV generations fast or slow?

- ▶ Fast generations mean:
 - ▶ Testing and treating will help less
 - ▶ *but* lower epidemic strength




- ▶ Predicted effectiveness of test and treat intervention minimally sensitive to proportion of early transmission
- ▶ Fast transmission:
 - ▶ low proportion prevented, but low \mathcal{R} estimate
- ▶ Slow transmission:
 - ▶ high proportion prevented, but high \mathcal{R} estimate
- ▶ *Eaton JW, Hallett TB. Proc Natl Acad Sci U S A. 2014 Nov 11;111(45):16202-7.*


Epidemic strength (present)




Strength of intervention




Different interventions (present)




Different interventions (present)




Epidemic speed




Epidemic speed



Speed of intervention



Different interventions (present)



The strength paradigm

- ▶ $\mathcal{R} > 1$ is a threshold
- ▶ If we can reduce transmission by a constant *factor* of $\theta > \mathcal{R}$, disease can be controlled
- ▶ In general, we can define θ as a (harmonic) mean of the reduction factor over the course of an infection
 - ▶ weighted by the *intrinsic* generation interval
- ▶ Epidemic is controlled if $\theta > \mathcal{R}$

The speed paradigm

- ▶ $r > 0$ is a threshold
- ▶ If we can reduce transmission at a constant *hazard rate* of $\phi > r$, disease can be controlled
- ▶ In general, we can define ϕ as a (very weird) mean of the reduction factor over the course of an infection
 - ▶ weighted by the *backward* generation interval
- ▶ Epidemic is controlled if $\phi > r$

Measuring the intervention



HIV

- ▶ The importance of transmission speed to HIV control is easier to understand using the speed paradigm
 - ▶ We know the speed of invasion
 - ▶ $\approx 0.7/\text{yr}$
 - ▶ Characteristic scale $\approx 1.4\text{yr}$
 - ▶ And can hypothesize the speed of intervention
 - ▶ Or aim to go fast enough

Paradigms are complementary

- ▶ HIV
 - ▶ Information and current intervention are both “speed-like”
- ▶ Measles
 - ▶ Information (long-term) is strength-like
 - ▶ Intervention (vaccine) also strength-like
- ▶ Ebola vaccination
 - ▶ Information is speed-like
 - ▶ Pre-emptive vaccination is strength-like



Thanks

- ▶ Department
- ▶ Collaborators
- ▶ Funders: NSERC, CIHR

Linking framework

- ▶ Epidemic speed (r) is a *product*:
 - ▶ (something to do with) generation speed ×
 - ▶ (something to do with) epidemic strength
- ▶ In particular:
 - ▶ $r \approx (1/\bar{G}) \times \ell(\mathcal{R}; \kappa_g)$
 - ▶ ℓ is the inverse of X