The SIR Model Family

Jonathan Dushoff

MMED 2017

Goals

- This lecture will:
  - introduce the idea of dynamical modeling
  - explain why dynamical modeling is a key tool for understanding infectious disease
  - discuss and demonstrate simple dynamical models from the SIR model family
  - investigate some insights that can be gained from these models

Dynamic modeling connects scales

- Start with rules about how things change in short time steps
  - Usually based on *individuals*
- Calculate results over longer time periods
  - Usually about *populations*

Compartmental models

Divide people into categories:



- Susceptible $\rightarrow$ Infectious $\rightarrow$ Recovered

What determines transition rates?

- People get better independently
- People get infected by infectious people

Conceptual modeling

- What is the final result?
- When does disease increase, decrease?

Dynamic implementation

- Requires assumptions about recovery and transmission
- The *conceptually simplest* implementation uses Ordinary Differential Equations (ODEs)
  - Other options may be more realistic
  - Or simpler in practice

Recovery

- Infectious people recover at *per capita* rate γ
  - Total recovery rate is γI
  - Mean time infectious is $D = 1/\gamma$

Transmission

- Susceptible people get infected by:
  - Going around and contacting people (rate c)
  - Some of these people are infectious (proportion I/N)
  - Some of these contacts are effective (proportion p)
- Per capita rate of becoming infected is $cpI/N \equiv \beta I/N$
- Population-level transmission rate is $\mathcal{T} = \beta SI/N$

Another perspective on transmission

- Infectious people infect others by:
  - Going around and contacting people (rate c)
  - Some of these people are susceptible (proportion S/N)
  - Some of these contacts are effective (proportion p)
- Per capita rate of infecting others is $cpS/N \equiv \beta S/N$
- Population-level transmission rate is $\mathcal{T} = \beta SI/N$

ODE implementation

$$
\begin{aligned}
\frac{dS}{dt} &= -\beta\frac{SI}{N} \\
\frac{dI}{dt} &= \beta\frac{SI}{N} - \gamma I \\
\frac{dR}{dt} &= \gamma I
\end{aligned}
$$

Spreadsheet implementation

`http://tinyurl.com/SIR-MMED-2017`

ODE assumptions

- Lots and lots of people
- Perfectly mixed

ODE assumptions



- Waiting times are exponentially distributed
- Rarely realistic

Scripts vs. spreadsheets

- Scripts are more transparent, less redundant
- Spreadsheets are more intuitive for simple problems

More about transmission

- $\beta = pc$
  - What is a contact?
  - What is the probability of transmission?
- Sometimes this decomposition is clear
- But usually it's not

Population sizes

$$
\begin{aligned}
\frac{dS}{dt} &= -\beta(N)\frac{SI}{N} \\
\frac{dI}{dt} &= \beta(N)\frac{SI}{N} - \gamma I \\
\frac{dR}{dt} &= \gamma I
\end{aligned}
$$



- $\beta(N) = \beta_0$

- $\mathcal{T} = \frac{\beta_0 SI}{N}$
- Also known as *frequency-dependent* transmission



- $\beta(N) = \beta_1 N$
- $\mathcal{T} = \beta_1 SI$
- Also known as *density-dependent* transmission



- May not go to zero when N does
- May not go to ∞ when N does

Digression – units

- $\mathcal{T} = \beta SI/N$: [ppl/time]
- β : [1/time]
  - $\beta/\gamma = \beta D$: [1]
  - Standard incidence, β_0 : [1/time]
  - Mass-action incidence, β_1 : [1/(people $\cdot$ time)]

Closing the circle





Tendency to oscillate



With individuality



Summary

- Dynamic models are an essential tool because they allow us to link between scales
- There are many ways to construct and implement dynamic models
- Very simple models can provide useful insights
  - Reproductive numbers and thresholds
  - Tendency for oscillation (and tendency for damping)
- More complex models can provide more detail, but also require more assumptions, and more choices
- Understanding simple models can help guide our understanding of more complicated models