

Method	Required information	Assumptions/Notes	References
Incidence based	<ul style="list-style-type: none"> • Generation interval CDF • Incidence reports 	<ul style="list-style-type: none"> • Uses cumulative incidence directly without estimating the initial growth rate • Entire generation interval distribution is required to obtain CDF 	[1]
Delta approximation	<ul style="list-style-type: none"> • Mean generation interval 	<ul style="list-style-type: none"> • Assumed fixed generation interval 	[2, 3]
Empirical (histogram-based)	<ul style="list-style-type: none"> • Generation interval samples 	<ul style="list-style-type: none"> • Relies on binning generation intervals into discrete histograms 	[2]
Empirical (sample-based)	<ul style="list-style-type: none"> • Generation interval samples 	<ul style="list-style-type: none"> • Requires more samples than the histogram-based method 	[4]
Euler-Lotka	<ul style="list-style-type: none"> • Entire generation interval distribution 	<ul style="list-style-type: none"> • Uses moment generating function 	[5]
Normal approximation	<ul style="list-style-type: none"> • Mean/SD generation interval 	<ul style="list-style-type: none"> • Assumes normally distributed generation intervals • Predicts decreasing r-\mathcal{R} relationship for high r 	[2]
SIR model	<ul style="list-style-type: none"> • Mean generation interval 	<ul style="list-style-type: none"> • Assumes exponentially distributed generation interval • Predicts linear relationship between r and \mathcal{R} 	[6, 7, 8, 2]
SEIR (gamma)	<ul style="list-style-type: none"> • Mean/SD latent period • Mean/SD infectious period 	<ul style="list-style-type: none"> • Assumes gamma distributed latent and infectious periods 	[9, 10, 2, 3]
SEIR (exponential)	<ul style="list-style-type: none"> • Mean generation interval • Ratio of the infectious period to the generation interval 	<ul style="list-style-type: none"> • Assumes exponentially distributed latent and infectious periods • Predicts quadratic relationship between r and \mathcal{R} 	[11, 12, 3]
Subexponential	<ul style="list-style-type: none"> • Generation interval shape • Corresponding generation interval parameters • Sub-exponential growth parameter • Disease generation 	<ul style="list-style-type: none"> • Assumes sub-exponential growth rate • Provides examples for uniform, gamma, exponential and delta distributed generation intervals 	[13]
Trapezoid approximation	<ul style="list-style-type: none"> • Period of no infection after exposure • Time until maximum infectivity • Duration of maximum infectivity • Time until recovery 	<ul style="list-style-type: none"> • Assumes trapezoid shaped infection kernel 	[3]

References

- [1] Nishiura H. Correcting the actual reproduction number: a simple method to estimate R_0 from early epidemic growth data. *International journal of environmental research and public health*. 2010;7(1):291–302.
- [2] Wallinga J, Lipsitch M. How generation intervals shape the relationship between growth rates and reproductive numbers. *Proceedings of the Royal Society of London B: Biological Sciences*. 2007;274(1609):599–604.
- [3] Roberts M, Heesterbeek J. Model-consistent estimation of the basic reproduction number from the incidence of an emerging infection. *Journal of mathematical biology*. 2007;55(5-6):803.
- [4] Hampson K, Dushoff J, Cleaveland S, Haydon DT, Kaare M, Packer C, et al. Transmission dynamics and prospects for the elimination of canine rabies. *PLoS biology*. 2009;7(3):e1000053.
- [5] Lotka AJ. Relation Between Birth Rates and Death Rates. *Science*. 1907 Jul;26(653):21–22.
- [6] Anderson RM, May RM. *Infectious diseases of humans: dynamics and control*. Oxford university press; 1992.
- [7] Pybus OG, Charleston MA, Gupta S, Rambaut A, Holmes EC, Harvey PH. The epidemic behavior of the hepatitis C virus. *Science*. 2001;292(5525):2323–2325.
- [8] Ferguson NM, Cummings DA, Cauchemez S, Fraser C, Riley S, Meeyai A, et al. Strategies for containing an emerging influenza pandemic in Southeast Asia. *Nature*. 2005;437(7056):209.
- [9] Anderson D, Watson R. On the spread of a disease with gamma distributed latent and infectious periods. *Biometrika*. 1980;67(1):191–198.
- [10] Wearing HJ, Rohani P, Keeling MJ. Appropriate models for the management of infectious diseases. *PLoS Med*. 2005;2(7):e174.
- [11] Lipsitch M, Cohen T, Cooper B, Robins JM, Ma S, James L, et al. Transmission dynamics and control of severe acute respiratory syndrome. *Science*. 2003;300(5627):1966–1970.
- [12] Chowell G, Nishiura H, Bettencourt LM. Comparative estimation of the reproduction number for pandemic influenza from daily case notification data. *Journal of the Royal Society Interface*. 2007;4(12):155–166.
- [13] Chowell G, Viboud C, Simonsen L, Moghadas SM. Characterizing the reproduction number of epidemics with early subexponential growth dynamics. *Journal of The Royal Society Interface*. 2016;13(123):20160659.