

Diagrammatic Categories in Representation Theory

Honours Thesis

Victor Zhang
Supervisor: Dr Anna Romanov

UNSW Australia

October 14, 2022

Contents

1	Introduction	1
2	Background	2
2.1	Coxeter Groups	2
2.2	Hecke Algebra	2

Chapter 1

Introduction

This page was empty.

Chapter 2

Background

Notation: we write 1 for the neutral element of a group.

2.1 Coxeter Groups

Definition 2.1.1. A *Coxeter system* (W, S) is a group W and a finite subset $S = \{s_1, \dots, s_n\} \subset W$ under the following conditions. For any $s, t \in S$ there exists $m_{st} \in \mathbb{Z}_{>0} \cup \{\infty\}$ such that $(st)^{m_{st}} = 1$, where $m_{st} = 1$ if $s = t$, and $m_{st} \in \{2, 3, \dots\} \cup \{\infty\}$ if $s \neq t$. In other words, $W = \langle s \in S \mid (st)^{m_{st}} = 1 \rangle$ with generator S . We call W a *Coxeter group*.

Note that $m_{st} = \infty$ indicates there are no relations of the form $(st)^m = 1$ for any $m \in \mathbb{Z}_{>0}$.

The first

2.2 Hecke Algebra