

17. Suppose that we observe  $m$  independent and identically distributed (IID) Bernoulli( $\theta$ ) random variables  $Y_1, Y_2, \dots, Y_m$ . Consider testing

$$H_0 : \theta \leq \theta_0 \text{ versus } H_1 : \theta > \theta_0$$

- (a) Find the maximum likelihood estimator (MLE)  $\hat{\theta}$  of  $\theta$ .
- (b) Under  $H_0 : \theta \leq \theta_0$ , find the restricted MLE  $\hat{\theta}_0$  of  $\theta$ .
- (c) Use R to draw the likelihood function  $L(\theta)$ , and illustrate both  $\hat{\theta}$  and  $\hat{\theta}_0$  on the graph.
- (d) Find the likelihood ratio test (LRT) of

$$H_0 : \theta \leq \theta_0 \text{ versus } H_1 : \theta > \theta_0$$

- (e) Use R to graph the likelihood ratio (LR) test statistic as a function of  $\sum_{i=1}^n Y_i$ . On the graph, illustrate the rejection region
- (f) Prove the likelihood ratio test statistic  $\lambda(\mathbf{y}) \leq c$  if and only if  $\bar{y} \geq k$  (or equivalently  $\sum_{i=1}^m y_i > mk$ )