

# 2072U Computational Science I

## Winter 2022

Week	Topic
1	Introduction
1–2	Solving nonlinear equations in one variable
3–4	Solving systems of (non)linear equations
5–6	Computational complexity
6–8	Interpolation and least squares
8–10	Integration & differentiation
10–12	Additional Topics

1. What conditions, how fast, how accurate?
2. Interpolation error
3. Computational complexity
4. Taylor polynomials

Key questions:

- ▶ Polynomial interpolation: what conditions, how fast, how accurate?
- ▶ What are Taylor polynomials?
- ▶ What are the differences between Taylor and interpolating polynomials?

The three questions:

1. When does it work?
2. How fast does it work?
3. How accurate is the result?

## Theorem (Polynomial interpolation error)

*Let  $\{x_k\}_{k=0}^n$  be distinct interpolation nodes in  $I \subset \mathbb{R}$  and let  $f$  be  $n + 1$  times continuously differentiable in  $I$ . Then,  $\forall x \in I, \exists \xi \in I$  such that*

$$E_n(x) := f(x) - \Pi_n(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{k=0}^n (x - x_k),$$

*where  $\Pi_n$  is the unique polynomial interpolant of degree at most  $n$  that interpolates the data  $\{(x_k, f(x_k))\}_{k=0}^n$ .*

## Theorem (Polynomial interpolation error)

Let  $\{x_k\}_{k=0}^n$  be distinct interpolation nodes in  $I \subset \mathbb{R}$  and let  $f$  be  $n + 1$  times continuously differentiable in  $I$ . Then,  $\forall x \in I, \exists \xi \in I$  such that

$$E_n(x) := f(x) - \Pi_n(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{k=0}^n (x - x_k),$$

where  $\Pi_n$  is the unique polynomial interpolant of degree at most  $n$  that interpolates the data  $\{(x_k, f(x_k))\}_{k=0}^n$ .

- $|E_n(x)| = |f(x) - \Pi_n(x)|$  is the **error of polynomial interpolation**

## Theorem (Polynomial interpolation error)

Let  $\{x_k\}_{k=0}^n$  be distinct interpolation nodes in  $I \subset \mathbb{R}$  and let  $f$  be  $n + 1$  times continuously differentiable in  $I$ . Then,  $\forall x \in I, \exists \xi \in I$  such that

$$E_n(x) := f(x) - \Pi_n(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{k=0}^n (x - x_k),$$

where  $\Pi_n$  is the unique polynomial interpolant of degree at most  $n$  that interpolates the data  $\{(x_k, f(x_k))\}_{k=0}^n$ .

- ▶  $|E_n(x)| = |f(x) - \Pi_n(x)|$  is the **error of polynomial interpolation**
- ▶ What theorem says: size of error controlled by
  - ▶ size of  $f^{(n+1)}(\xi)$
  - ▶ size of  $\prod_{k=0}^n (x - x_k)$

## Theorem (Polynomial interpolation error)

Let  $\{x_k\}_{k=0}^n$  be distinct interpolation nodes in  $I \subset \mathbb{R}$  and let  $f$  be  $n + 1$  times continuously differentiable in  $I$ . Then,  $\forall x \in I, \exists \xi \in I$  such that

$$E_n(x) := f(x) - \Pi_n(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{k=0}^n (x - x_k),$$

where  $\Pi_n$  is the unique polynomial interpolant of degree at most  $n$  that interpolates the data  $\{(x_k, f(x_k))\}_{k=0}^n$ .

- ▶  $|E_n(x)| = |f(x) - \Pi_n(x)|$  is the **error of polynomial interpolation**
- ▶ What theorem says: size of error controlled by
  - ▶ size of  $f^{(n+1)}(\xi)$
  - ▶ size of  $\prod_{k=0}^n (x - x_k)$
- ▶ This error comes on top of the error of linear solving.



Because we can't know  $\xi$  unless we know the error, for practical application we use the theorem to find an upper bound on the error. In particular, the theorem implies:


$$|E_n(x)| := |f(x) - \Pi_n(x)| \leq \max_{x \in I} \frac{|f^{(n+1)}(x)|}{(n+1)!} \prod_{k=0}^n |x - x_k|,$$

Back to the last example:

$x$	0	$\pi/6$	$\pi/4$	$\pi/3$	$\pi/2$
$\sin(x)$	0	$1/2$	$1/\sqrt{2}$	$\sqrt{3}/2$	1

these exact values can be used to approximate  $\sin(x)$  by a polynomial:


$$\sin(x) \approx 0.9956261 x + 0.021372984 x^2 - 0.2043407 x^3 + 0.02879711 x^4 \quad (\text{for } 0 < x < \pi/2)$$



## Maximum Error

$$|f(x) - \Pi_n(x)| \leq \max_{x \in I} \frac{|f^{(n+1)}(x)|}{(n+1)!} \prod_{k=0}^n |x - x_k|,$$

$$\begin{aligned} |\sin(x) - \Pi_4(x)| &\leq \max_{x \in I} \frac{|f^{(5)}(x)|}{5!} |x| \left| x - \frac{\pi}{6} \right| \left| x - \frac{\pi}{4} \right| \left| x - \frac{\pi}{3} \right| \left| x - \frac{\pi}{2} \right| \\ &\leq \max_{x \in I} \frac{|\cos(x)|}{5!} |x| \left| x - \frac{\pi}{6} \right| \left| x - \frac{\pi}{4} \right| \left| x - \frac{\pi}{3} \right| \left| x - \frac{\pi}{2} \right| \\ &\leq \end{aligned}$$



The difference between  $\sin(x)$  and  $\Pi_4(x)$  is less than  $3 \times 10^{-4}$  on  $[0, \pi/2]$ .

Three conclusions we can draw from the error formula:

1. Functions with *small higher derivatives* are well-approximated by interpolating polynomials.  
Such functions are *smooth*. The smoothest functions are the polynomials.
2. We can choose the location of the interpolation nodes to minimize the error (see CSII).  
In fact, it turns out that *equidistant* nodes are bad. Good sets of interpolation nodes have more nodes near the boundaries.
3. *Extrapolation* is far more dangerous than *interpolation*.  
The upper bound for the error of extrapolation grows without bound – as  $x^{n+1}$ .

Recall: To compute the polynomial interpolant, we can solve  $\forall \mathbf{a} = \mathbf{y}$  :

$$\underbrace{\begin{bmatrix} 1 & x_0^1 & \cdots & x_0^{n-1} & x_0^n \\ 1 & x_1^1 & \cdots & x_1^{n-1} & x_1^n \\ 1 & x_2^1 & \cdots & x_2^{n-1} & x_2^n \\ \vdots & \vdots & & \vdots & \vdots \\ 1 & x_n^1 & \cdots & x_n^{n-1} & x_n^n \end{bmatrix}}_V \underbrace{\begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}}_a = \underbrace{\begin{bmatrix} y_0 \\ y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}}_y$$

To compute FLOPS, we need to consider

- ▶ cost of building the matrix (see Assignment 3)
- ▶ cost of solving the system

Is there a more efficient approach?

Is there another approach that doesn't lead to badly conditioned systems?

Recall: The interpolant is written as:

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$



Recall: The interpolant is written as:

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Instead of using monomial basis  $\phi_k(x) = x^k$ , use:

*Newton polynomial basis*

$$\phi_0(x) = 1 \quad \phi_1(x) = x - x_0 \quad \phi_2(x) = (x - x_0)(x - x_1) \quad \dots$$

$$\phi_n(x) = (x - x_0)(x - x_1) \dots (x - x_{n-1})$$

Recall: The interpolant is written as:

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Instead of using monomial basis  $\phi_k(x) = x^k$ , use:

*Newton polynomial basis*

$$\phi_0(x) = 1 \quad \phi_1(x) = x - x_0 \quad \phi_2(x) = (x - x_0)(x - x_1) \quad \dots$$

$$\phi_n(x) = (x - x_0)(x - x_1) \dots (x - x_{n-1})$$

With this basis, the resulting system of linear equations for the coefficients  $\{a_k\}_{k=0}^n$  is  $M\mathbf{a} = \mathbf{y}$ , where

- ▶ the matrix  $M$  is now triangular
- ▶ the matrix  $M$  depends on the interpolation points  $x_k$

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Write out interpolation conditions  $\Pi_n(x_k) = y_k$  for new basis

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Write out interpolation conditions  $\Pi_n(x_k) = y_k$  for new basis

►  $\Pi_n(x_0) = y_0 \quad \rightarrow \quad a_0 = y_0$

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Write out interpolation conditions  $\Pi_n(x_k) = y_k$  for new basis

- ▶  $\Pi_n(x_0) = y_0 \quad \rightarrow \quad a_0 = y_0$
- ▶  $\Pi_n(x_1) = y_1 \quad \rightarrow \quad a_0 + a_1(x_1 - x_0) = y_1$

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Write out interpolation conditions  $\Pi_n(x_k) = y_k$  for new basis

- ▶  $\Pi_n(x_0) = y_0 \quad \rightarrow \quad a_0 = y_0$
- ▶  $\Pi_n(x_1) = y_1 \quad \rightarrow \quad a_0 + a_1(x_1 - x_0) = y_1$
- ▶  $\Pi_n(x_2) = y_2 \quad \rightarrow \quad a_0 + a_1(x_2 - x_0) + a_2(x_2 - x_0)(x_2 - x_1) = y_2$

$$\Pi_n(x) = \sum_{k=0}^n a_k \phi_k(x) = a_0 \phi_0(x) + a_1 \phi_1(x) + \cdots + a_n \phi_n(x)$$

Write out interpolation conditions  $\Pi_n(x_k) = y_k$  for new basis

- ▶  $\Pi_n(x_0) = y_0 \rightarrow a_0 = y_0$
- ▶  $\Pi_n(x_1) = y_1 \rightarrow a_0 + a_1(x_1 - x_0) = y_1$
- ▶  $\Pi_n(x_2) = y_2 \rightarrow a_0 + a_1(x_2 - x_0) + a_2(x_2 - x_0)(x_2 - x_1) = y_2$
- ▶  $\vdots$

$$\underbrace{\begin{bmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & (x_1 - x_0) & 0 & 0 & \cdots \\ 1 & (x_2 - x_0) & (x_2 - x_0)(x_2 - x_1) & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}}_M \underbrace{\begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}}_a = \underbrace{\begin{bmatrix} y_0 \\ y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}}_y$$

## Theorem (Taylor's Theorem with Lagrange Remainder)

If  $f^{(n)}$  is continuous in the interval  $[a, b]$  and if  $f^{(n+1)}$  exists on the open interval  $(a, b)$ , then for any points  $c$  and  $x$  in  $[a, b]$  there exists a  $\xi$  between  $c$  and  $x$  such that

$$f(x) = T_n(x) + R_n(x), \quad \text{with}$$

$$T_n(x) = \text{nth Taylor polynomial}$$

$$= \sum_{k=0}^n \frac{f^{(k)}(c)}{k!} (x - c)^k \quad \text{and}$$

$$R_n(x) = \text{remainder or truncation error}$$

$$= \frac{f^{(n+1)}(\xi)}{(n+1)!} (x - c)^{n+1}$$



Example of a Taylor polynomial: e.g.,  $f(x) = e^x$ ,  $c = 0$ ,  $n = 3$

Example of a Taylor polynomial: e.g.,  $f(x) = e^x$ ,  $c = 0$ ,  $n = 3$

►  $f(0) = f'(0) = f''(0) = f^{(3)}(0) = e^0 = 1$ ,  $f^{(4)}(\xi) = e^\xi$

Example of a Taylor polynomial: e.g.,  $f(x) = e^x$ ,  $c = 0$ ,  $n = 3$

$$\blacktriangleright f(0) = f'(0) = f''(0) = f^{(3)}(0) = e^0 = 1, f^{(4)}(\xi) = e^\xi$$

$$\blacktriangleright T_3(x) = \sum_{k=0}^3 \frac{f^{(k)}(0)}{k!} (x - 0)^k = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$

Example of a Taylor polynomial: e.g.,  $f(x) = e^x$ ,  $c = 0$ ,  $n = 3$

$$\blacktriangleright f(0) = f'(0) = f''(0) = f^{(3)}(0) = e^0 = 1, f^{(4)}(\xi) = e^\xi$$

$$\blacktriangleright T_3(x) = \sum_{k=0}^3 \frac{f^{(k)}(0)}{k!} (x - 0)^k = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$

$$\blacktriangleright R_3(x) = \frac{f^{(4)}(\xi)}{4!} x^4 = \frac{e^\xi}{24} x^4$$

Example of a Taylor polynomial: e.g.,  $f(x) = e^x$ ,  $c = 0$ ,  $n = 3$

$$\blacktriangleright f(0) = f'(0) = f''(0) = f^{(3)}(0) = e^0 = 1, f^{(4)}(\xi) = e^\xi$$

$$\blacktriangleright T_3(x) = \sum_{k=0}^3 \frac{f^{(k)}(0)}{k!} (x - 0)^k = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$

$$\blacktriangleright R_3(x) = \frac{f^{(4)}(\xi)}{4!} x^4 = \frac{e^\xi}{24} x^4$$

For this function, we can consider the *series*

$$\exp(x) = \sum_{k=0}^{\infty} \frac{1}{k!} x^k$$

This series converges for any  $x \in \mathbb{R}$ .

We say it has an *infinite radius of convergence*.

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$

$$S_3(x) = x - \frac{x^3}{3!}$$

Nested form of polynomials simplifies evaluation.



Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$

$$S_3(x) = x - \frac{x^3}{3!} = x \left( 1 - \frac{x^2}{2 \cdot 3} \right)$$

Nested form of polynomials simplifies evaluation.

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$

$$S_3(x) = x - \frac{x^3}{3!} = x \left( 1 - \frac{x^2}{2 \cdot 3} \right)$$

$$S_5(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!}$$

Nested form of polynomials simplifies evaluation.

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$

$$S_3(x) = x - \frac{x^3}{3!} = x \left( 1 - \frac{x^2}{2 \cdot 3} \right)$$

$$S_5(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} = x \left( 1 + \frac{x^2}{2 \cdot 3} \left( -1 + \frac{x^2}{4 \cdot 5} \right) \right)$$

Nested form of polynomials simplifies evaluation.

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$

$$S_3(x) = x - \frac{x^3}{3!} = x \left( 1 - \frac{x^2}{2 \cdot 3} \right)$$

$$S_5(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} = x \left( 1 + \frac{x^2}{2 \cdot 3} \left( -1 + \frac{x^2}{4 \cdot 5} \right) \right)$$

$$S_7(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!}$$

Nested form of polynomials simplifies evaluation.

Second example: Taylor polynomials for  $\sin x$ :

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1} \quad (x \in \mathbb{R})$$

again with an infinite radius of convergence.

$$S_1(x) = x$$


$$S_3(x) = x - \frac{x^3}{3!} = x \left( 1 - \frac{x^2}{2 \cdot 3} \right)$$

$$S_5(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} = x \left( 1 + \frac{x^2}{2 \cdot 3} \left( -1 + \frac{x^2}{4 \cdot 5} \right) \right)$$


$$S_7(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} = x \left( 1 + \frac{x^2}{2 \cdot 3} \left( -1 + \frac{x^2}{4 \cdot 5} \left( 1 - \frac{x^2}{6 \cdot 7} \right) \right) \right)$$

Nested form of polynomials simplifies evaluation.

# sin x and Taylor polynomials



# Taylor polynomial errors for $\sin x$



Third example:  $f(x) = \ln(1 + x)$ .

$$\begin{aligned} f^{(1)}(x) &= \frac{1}{1+x} & f^{(3)}(x) &= \frac{2}{(1+x)^3} \\ f^{(2)}(x) &= \frac{-1}{(1+x)^2} & f^{(k)}(x) &= \frac{(-1)^{k+1}(k-1)!}{(1+x)^k} \end{aligned}$$

so  $f^{(1)}(0) = 1$ ,  $f^{(2)}(0) = -1$ ,  $f^{(3)}(0) = 2, \dots$ ,  
 $f^{(k)}(0) = (-1)^{k+1}(k-1)!$  and

$$\ln(1+x) = \sum_{k=1}^n \frac{(-1)^{k+1}}{k} x^k + R_n(x), \quad R_n(x) = \frac{(-1)^n x^{n+1}}{(n+1)(1+\xi)^{n+1}}$$

This series converges for  $-1 < x < 1$ , we say the radius of convergence is 1.



## Summary for Taylor polynomials:

- ▶ The Taylor polynomial exists if  $f$  is sufficiently smooth.
- ▶ We have an expression for the remainder.
- ▶ The computation of Taylor polynomials requires the computation of derivatives. **Not many functions have such simple derivatives as in our examples!**
- ▶ Properties of  $f$  at  $x = c$  completely determine  $T_n(x)$ , so all information comes from  $x = c$ . **The Taylor polynomial is useful as a local approximation only.**
- ▶ For  $|x - c|$  large, many terms needed for convergence.

In contrast, polynomial interpolation is useful on a whole domain.