

Maintenance olaparib plus bevacizumab in patients with newly diagnosed, advanced high-grade ovarian carcinoma: final analysis of second progression-free survival (PFS2) in the Phase III PAOLA-1/ENGOT-ov25 trial

Antonio González Martín,¹ Youssef Tazi,² Florian Heitz,³ Laure Montane,⁴ Piera Gargiulo,⁵ Regina Berger,⁶ Kan Yonemori,⁷ Ignace Vergote,⁸ Alessandra Bologna,⁹ Johanna Mäenpää,¹⁰ Cristina Costan,¹¹ Ulrich Canzler,¹² Claudio Zamagni,¹³ Eva Maria Guerra-Alia,¹⁴ Charles Briac Levaché,¹⁵ Frederik Marmé,¹⁶ Elsa Kalbacher,¹⁷ Nikolaus de Gregorio,¹⁸ Nadine Dohollou,¹⁹ Isabelle Ray-Coquard²⁰

¹Clinica Universidad de Navarra, Madrid, and GEICO, Spain; ²Strasbourg Oncologie Libérale, Strasbourg, and GINECO, France; ³Kliniken Essen-Mitte, Essen, and AGO, Germany; ⁴Centre Léon Bérard, Lyon, France; ⁵Istituto Nazionale Tumori IRCCS Fondazione G. Pascale, Naples, and MITO, Italy; ⁶Medical University of Innsbruck, Univ. Clinic for Gynaecology and Obstetrics, Innsbruck, and AGO-Austria, Austria; ⁷National Cancer Center, Tokyo, and GOTIC, Japan; ⁸University Hospital Leuven, Leuven Cancer Institute, Leuven, and BGOG, Belgium; ⁹Azienda Unità Sanitaria Locale di Reggio Emilia, IRCCS, Reggio Emilia, and MANGO, Italy; ¹⁰Tampere University and University Hospital, Tampere, and NSGO, Finland; ¹¹Hôpital Michallon, Centre Hospitalier Universitaire de Grenoble, and GINECO, France; ¹²Universitätsklinikum Carl Gustav Carus, Technische Universität Dresden, Dresden, and AGO, Germany; ¹³Policlinico S.Orsola-Malpighi, Bologna, and MITO, Italy; ¹⁴Hospital Universitario Ramón y Cajal, Madrid, and GEICO, Spain; ¹⁵Polyclinique Francheville, Périgueux, and GINECO, France; ¹⁶Universitätsklinikum Heidelberg, Heidelberg, and AGO, Germany; ¹⁷CHRU de Besançon, Hôpital Jean Minjoz, Besançon, and GINECO, France; ¹⁸Universitätsklinikum Ulm, Ulm, and AGO, Germany; ¹⁹Centre Léon Bérard, Lyon, and GINECO, France; ²⁰Centre Léon Bérard and University Claude Bernard Lyon 1, Lyon, and GINECO, France

UNCONTROLLED COPY



Downloadable
slides

Copies of this presentation obtained through QR (Quick Response) codes are for personal use only and may not be reproduced without written permission of the authors

I Ray-Coquard: Honoraria (self) from Abbvie, Agenus, Advaxis, BMS, PharmaMar, Genmab, Pfizer, AstraZeneca, Roche, GSK, MSD, Deciphera, Mersena, Merck Sereno, Novartis, Amgen, Tesaro and Clovis; honoraria (institution) from GSK, MSD, Roche and BMS; Novartis, AstraZeneca, Roche, GSK, Advaxis, BMS, PharmaMar, Genmab, Pfizer, AstraZeneca, Roche/Genentech, GSK, MSD, Deciphera, Mersena, Merck Sereno, Novartis, Amgen, Tesaro and Clovis; research grant/funding (self) from MSD, Roche and BMS; research grant/funding (institution) from MSD, Roche, BMS, Novartis, Astra Zeneca and Merck Sereno; and travel support from Roche, AstraZeneca and GSK


PAOLA-1/ENGOT-ov25 trial design

Patients:

- Newly diagnosed, FIGO stage III–IV high-grade serous or endometrioid ovarian, fallopian tube and/or primary peritoneal cancer*

First-line treatment:

- Upfront or interval surgery
- Platinum-taxane based chemotherapy plus ≥ 2 cycles of bevacizumab[†]



- Primary endpoint: investigator-assessed PFS (RECIST v1.1)
- In the primary analysis, a statistically significant PFS benefit was observed¹

Primary PFS analysis (DCO 22 March 2019)

	Olaparib + bev (N=537)	Placebo + bev (N=269)
Median PFS, months	22.1	16.6
HR (95% CI); P value	0.59 (0.49–0.72) P<0.001	


*Patients with other epithelial non-mucinous ovarian cancer were eligible if they had a germline *BRCA1* and/or *BRCA2* mutation; [†]Patients must have received ≥ 3 cycles of bevacizumab with the last 3 cycles of chemotherapy, apart from patients undergoing interval surgery who were permitted to receive only 2 cycles of bevacizumab with the last 3 cycles of chemotherapy; [‡]Bevacizumab 15 mg/kg, every 3 weeks for a total of 15 months, including when administered with chemotherapy; [§]According to timing of surgery and NED/CR/PR

bid, twice daily; BRCAm, BRCA mutation; CI, confidence interval; CR, complete response; DCO, data cut-off; FIGO, International Federation of Gynecology and Obstetrics; HR, hazard ratio; NED, no evidence of disease; PFS; time from randomization to progression or death; PR, partial response; RECIST, Response Evaluation Criteria in Solid Tumours;

1. Ray-Coquard I *et al.* *N Engl J Med* 2019;381:2416–28

PAOLA-1/ENGOT-ov25 PFS2 analysis

PFS2 is measured from the time of randomization to second progression or death and evaluates the effect of maintenance therapy with olaparib plus bevacizumab beyond first progression



Primary endpoint

- Investigator-assessed PFS

Secondary endpoints

- TFST

- PFS2**

- TSST

- OS


- HRQoL

- Safety and tolerability

- PFS2 was immature at the time of primary PFS analysis (DCO 22 March 2019)
- We present the **prespecified final PFS2 analysis** planned for ≈53% data maturity or 1 year after primary analysis (DCO 22 March 2020)
- We also present post hoc analyses of PFS2 by biomarker status

UNCONTROLLED COPY

A statistically significant PFS2 benefit was observed with olaparib plus bevacizumab in the ITT population



Olaparib + bev (N=537)	Placebo + bev (N=269)
260 (48)	164 (61)
36.5	32.6
HR 0.78 (95% CI 0.64–0.95) P=0.0125	

Patients receiving a PARP inhibitor during first subsequent treatment:
 Olaparib plus bevacizumab: **9.1%** (49/537)
 Placebo plus bevacizumab: **26.8%** (72/269)


Median PFS2 follow-up of 35.5 months for olaparib plus bevacizumab and 36.5 months for placebo plus bevacizumab

Number of patients at risk:

Olaparib + bev	537	527	515	491	459	434	408	376	339	309	263	177	99	71	27	14	6	0
Placebo + bev	269	266	258	245	226	215	206	190	171	149	131	106	72	40	27	9	4	1

PFS2 subgroup analysis by HRD status

HRD positive,* including tumour BRCAm




Events, n (%)

Median PFS2, months

	Olaparib + bev (n=255)	Placebo + bev (n=132)
Events, n (%)	85 (33)	70 (53)
Median PFS2, months	50.3 [†]	35.3
HR	HR 0.56 (95% CI 0.41–0.77)	

HRD positive,* excluding tumour BRCAm




Events, n (%)

Median PFS2, months

	Olaparib + bev (n=97)	Placebo + bev (n=55)
Events, n (%)	41 (42)	33 (60)
Median PFS2, months	50.3 [†]	30.1
HR	HR 0.60 (95% CI 0.38–0.96)	

HRD negative/unknown




Events, n (%)

Median PFS2, months

	Olaparib + bev (n=282)	Placebo + bev (n=137)
Events, n (%)	175 (62)	94 (69)
Median PFS2, months	26.3	28.1
HR	HR 0.98 (95% CI 0.77–1.27)	

The PFS2 benefit was supported by a statistically significant* TSST improvement in the ITT population



Olaparib + bev (N=537)	Placebo + bev (N=269)
266 (50)	164 (61)
38.2	31.5
HR 0.78 (95% CI 0.64–0.95) P=0.0115*	

Biomarker subgroup	TSST HR (95% CI)
Tumour BRCAm	0.48 (0.31–0.75)
HRD positive	0.48 (0.35–0.66)
HRD negative/unknown	1.05 (0.82–1.36)

Number of patients at risk:


Olaparib + bev	537	528	520	505	469	442	421	394	364	329	297	245	184	127	86	39	19	6	1
Placebo + bev	269	267	260	247	232	218	207	191	172	149	135	113	75	49	31	17	5	1	0

Months since Randomization

UNCONTROLLED COPY

*TSST analysis not adjusted for multiplicity

Interim OS analysis



Number of patients at risk:

Olaparib + bev	537	528	526	515	499	475	458	436	416	392	366	306	245	167	110	53	25	8	2
Placebo + bev	269	267	264	261	250	240	227	218	206	197	185	158	119	81	58	33	11	3	1


Events, n (%)

Median OS, months

Olaparib + bev (N=537)	Placebo + bev (N=269)
195 (36)	108 (40)
NR	45.8
HR 0.93 (95% CI 0.74–1.18) P=0.5631	

- Immature OS analysis:
 - The low event rate (38%) means that definitive conclusions cannot be drawn
- Updated OS data will be presented at greater data maturity:
 - Prespecified final OS analysis planned at ≈60% data maturity or 3 years after primary PFS analysis (March 2022)

Safety analyses



- Olaparib + bev: All grades (frequency ≥10%)
- Olaparib + bev: Grade ≥3
- Placebo + bev: All grades (frequency ≥10%)
- Placebo + bev: Grade ≥3

AEs of special interest

	Primary PFS analysis (DCO 22 March 2019)		Final PFS2 analysis (DCO 22 March 2020)	
	Olaparib + bev (n=535)	Placebo + bev (n=267)	Olaparib + bev (n=535)	Placebo + bev (n=267)
MDS/AML/AA, n (%)	6 (1.1)	1 (0.4)	6 (1.1)	4 (1.5)‡
New primary malignancies,§ n (%)	7 (1.3)	3 (1.1)	14 (2.6)	5 (1.9)
Pneumonitis/ILD/bronchiolitis, n (%)	6 (1.1)	0	6 (1.1)	0
Median duration of olaparib/placebo treatment, months			17.3	15.6
Median duration of bev treatment, months			11.0	10.6
Discontinuation due to AEs, n (%)			112 (21)	15 (6)

*Grouped-term AEs; †All-grade grouped-term thrombocytopenia occurred in 8% of olaparib plus bevacizumab patients and 3% of placebo plus bevacizumab patients; grade ≥3 grouped-term thrombocytopenia occurred in 2% of olaparib plus bevacizumab patients and <1% of placebo plus bevacizumab patients; ‡3 of the 4 patients in the placebo plus bevacizumab group who developed MDS/AML/AA received a PARP inhibitor as first subsequent treatment before onset of AML.

§At primary PFS analysis, new primary malignancies in the olaparib plus bevacizumab group were acute lymphocytic leukaemia (n=1), breast cancer (n=2), lung cancer (n=1), myeloma (n=1), squamous skin cancer (n=1), and pancreatic cancer (n=1), and in the placebo group were breast cancer (n=2) and thyroid cancer (n=1). Additional new primary malignancies reported at final PFS2 analysis in the olaparib plus bevacizumab group were breast cancer (n=5), squamous skin cancer (n=1), and colon cancer (n=1), and in the placebo group were breast cancer (n=1) and malignant neoplasm (n=1).

AA, aplastic anaemia; AE, adverse event; AML, acute myeloid leukaemia; ILD, interstitial lung disease; MDS, myelodysplastic syndrome; UTI, urinary tract infection

Conclusions

- In PAOLA-1/ENGOT-ov25, the addition of maintenance olaparib to bevacizumab provided continued benefit beyond first progression, with a statistically significant improvement in PFS2:
 - A substantial PFS2 benefit was seen in patients who were HRD positive, regardless of tumour BRCAm status
- The significant PFS2 improvement was supported by a significant delay in TSST
- No new safety signals were observed with longer-term follow-up
- OS data are still immature



Copies of this presentation obtained through QR (Quick Response) codes are for personal use only and may not be reproduced without written permission of the authors

Acknowledgements

We thank the patients, their families, the investigators, and the staff

GINECO – France I Ray-Coquard

S Abadie-Lacourtoisie
J Alexandre
D Berton-Rigaud
E Boissier
H Bourgeois
A Chevalier-Place
P Combe
C Costan
J Dauba
L De Cock
C Desauw
R Despax
N Dohollou
C Dubot
M Fabbro
L Favier
A Floquet
P Follana
G Garnier
C Garnier Tixidre
L Gladieff
J Grenier
C Guillemet
AC Hardy-Bessard
F Joly
E Kalbacher
MC Kaminsky

JE Kurtz
R Largillier
CB Levache
C Lefeuve-Plesse
A Lesoin
T L'Haridon
A Lortholary
JP Lotz
J Meunier
M Mousseau
MA Mouret-Reynier
P Pautier
T Petit
M Provansal
E Pujade-Lauraine
N Raban
M Rodrigues
F Selle
R Sverdlin
Y Tazi
B You
AGO DE – Germany
P Harter
B Aktas
DO Bauerschlag
T Beck
A Belau
H Bronger
S Buchholz

P Buderath
A Burges
U Canzler
N de Gregorio
D Denschlag
M Dieterich
M Eichbaum
A El-Balat
G Emons
P Fasching
G Feisel-Schwickardi
M Frank
M Friedrich
EM Grischke
M Gropp-Meier
L Hanker
C Hannig
A Hasenburger
M Hellriegel
U Herwig
M Heubner
J Hulde
C Jackisch
M Kögel
P Krieger
T Kühn
C Liebrich
HJ Lück
P Mallmann

F Marmé
W Meier
V Möbus
OF Mohamed
C Nestle-Krämling
T Neunhöffer
G Oskay-Özcelik
TW Park-Simon
B Rautenberg
D Rein
W Ruhwedel
I Runnebaum
J Sagasser
B Schmalfeldt
A Schnelzer
A Schneeweiss
H Scholz
J Sehouli
A Sperfeld
A Steckkönig
HG Strauß
O Tomé
J Treustedt
H Voß
A Wischnik
R Witteler
A Wöckel
HH Woeltjen
A Zorr

MITO – Italy
S Pignata
S Cinieri
D Lorusso
AM Mosconi
A Savarese
G Scambia
R Sorio
C Zamagni
GEICO – Spain
A González-Martín
R Bratos
C Caballero
Y Garica
EM Guerra Alia
S Hernando
A Herrero
N Lainez
L Manso
C Martin
E Murata
E Ortega
I Palacio
A Poveda
I Romero
MJ Rubio
AGO AU – Austria
U Pöschl
G Bogner

E Petru
A Reinthaller
C Schauer
P Sevela
GOTIC – Japan
K Fujiwara
H Fujiwara
H Kobayashi
T Matsumoto
S Nagao
T Satoh
K Yonemori
H Yoshida
BGG – Belgium
I Vergote
L D'Hondt
P Vuylsteke
MANGO – Italy
N Colombo
A Bologna
G Tognon
**NSGO – Denmark,
Sweden and Finland**
J Mäenpää
S Hietanen
G Lindahl
TJ Nøttrup
U Pöschl

Sponsor ARCAGY

S Armanet, S Mijonnet,
C Montoto-Grillot,
A Morvan, K Thiam-Kieffer
B Votan

Study group coordinators

AGO De: G Elser, N Cron
AGO Au: R Berger, A Riha
MITO and MANGO: J Bryce
GEICO: C Larios
GOTIC: H Naito
NSGO: N Buchner Vinum

IDMC

J Vermorken
S Kaye
G Pond

Ascopharm Gr Novasco

S Perrin Brutto
A Lasfargues

Statisticians

D Perol
S Chabaud
C Cropet
L Montané
QOL: A Anota

Screening platforms

Institut Curie
Gustave Roussy
APH
Institut Bergonié
Centre François Baclesse

Centre de Ressources Biologiques of ARCAGY- GINECO (Institut Curie)

L Fuhrmann
A Degnieau

French National Cancer Institute (INCa)

Funding

ARCAGY Research,
AstraZeneca, Merck & Co.,
Inc. and F. Hoffmann-La
Roche

Medical writing support

Gillian Keating, MBChB,
funded by AstraZeneca and
Merck & Co., Inc.

UNCONTROLLED COPY

