



Macular Corneal Dystrophy: An Updated Review

Shalini Singh , Sujata Das , Chitra Kannabiran , Saumya Jakati & Sunita Chaurasia

To cite this article: Shalini Singh , Sujata Das , Chitra Kannabiran , Saumya Jakati & Sunita Chaurasia (2020): Macular Corneal Dystrophy: An Updated Review, Current Eye Research, DOI: [10.1080/02713683.2020.1849727](https://doi.org/10.1080/02713683.2020.1849727)

To link to this article: <https://doi.org/10.1080/02713683.2020.1849727>



Accepted author version posted online: 10 Nov 2020.



Submit your article to this journal [↗](#)



Article views: 3



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis

Journal: *Current Eye Research*

DOI: 10.1080/02713683.2020.1849727

Macular Corneal Dystrophy: An Updated Review

Authors: ¹Shalini Singh, MS; ²Sujata Das, FRCS; ³Chitra Kannabiran, PhD; ⁴Saumya Jakati, MD; ¹Sunita Chaurasia, MD

¹Cornea and Anterior Segment services, LVPEI, Hyderabad

²Cornea and Anterior segment services, LVPEI, Bhubneshwar

³Kallam Anji Reddy Molecular Genetics Laboratory, Prof Brien Holden Eye Research Centre

⁴Ophthalmic Pathology Laboratory, LVPEI, Hyderabad

Corresponding Author

Sunita Chaurasia, MD

Cornea and Anterior Segment services,

LV Prasad Eye Institute,

Banjara Hills, Hyderabad-560036

Email: sunita@lvpei.org

Word Count: 2473

Keywords- macular corneal dystrophy; genetics; keratoplasty; CHST6; glycosaminoglycans

ABSTRACT

Macular Corneal Dystrophy is an autosomal recessive form of corneal dystrophy due to a mutation in *CHST6* gene, which results in abnormal proteoglycan synthesis. There is accumulation of abnormal glycosaminoglycans in the corneal stroma and endothelium. The deposition results in progressive loss of corneal transparency and visual acuity. The histopathology shows characteristic alcian blue positive deposits. Management in the cases with visual loss requires keratoplasty either full thickness or lamellar. The decision about the ideal type of keratoplasty depends on age and pre-operative clinical features. Although prognosis after keratoplasty is good, recurrences can occur. Future research should be targeted towards gene therapy in this condition.

INTRODUCTION

Corneal dystrophies are typically defined as rare, inherited disorders of the cornea, that are bilateral, often symmetric, slowly progressive and not related to environmental and systemic factors.^{1,2} Abnormal accumulation of deposits in the different layers of the cornea determine the primary characteristics.³ However, there are deviations to the typical description of dystrophies as not all dystrophies are inherited, bilateral, symmetrically bilateral or associated with deposition of abnormal substances in the different layers of the cornea. The traditional system of classification of corneal dystrophies was based on the layer of cornea primarily affected. The newer classification of corneal dystrophies by the International Committee for Classification of Corneal Dystrophy (IC3D) are based on the clinical and genetic documentation of the dystrophies.^{4,5} As per the IC3D system, macular corneal dystrophy

(MCD)[MIM#217800] falls in the Category 1, i.e., a well-defined corneal dystrophy in which the gene has been mapped and identified and the specific mutations are known.

MCD was first described by Groenouw in 1890. It is an autosomal recessive corneal stromal dystrophy, characterized by deposition of glycosaminoglycans within keratocytes and corneal endothelium with extracellular deposition of the similar material in corneal stroma and Descemet membrane.³ Although classified under stromal dystrophies, the condition can have variable involvement of the endothelium- Descemet membrane complex.^{3,6}

EPIDEMIOLOGY

The prevalence of MCD is higher in those communities where consanguinity is common. Although it has been recognised throughout the world, its prevalence is notably more common in India, Saudi Arabia and Iceland.⁷⁻¹⁰ According to the registry maintained by Klintworth in the United states, prevalence of MCD is 0.3 individuals per 2,50,000 inhabitants.^{7,11} Estimates from Claims data in US, found that the prevalence of stromal dystrophies was low (~1%), but amongst the three dystrophies namely lattice, granular and macular, MCD was commonest.¹² A report from Iceland indicated that MCD accounted for one third of corneal transplants.⁸ Similarly, in the studies from Saudi Arabia, it was suggested that MCD was the most common corneal stromal dystrophy that required keratoplasty.^{9,13} A recent retrospective electronic medical record based study from Southern India, reported MCD to be the second commonest corneal dystrophy after Fuchs endothelial corneal dystrophy.¹⁴

CLINICAL FEATURES

Patient's symptoms commonly include photophobia, tearing and gradually deteriorating vision, which commonly manifests in the second and third decade of life. The lesions of MCD are mostly symmetrical and grayish white in color. As per the classical description of

MCD, the deposits are seen in the superficial cornea as fleck like opacities in the initial stages, that later coalesce to reach the limbus, deeper stroma and the endothelium.^{1-3,7} However, the diversity in the clinical presentation of MCD is increasingly recognized in the recent years.¹⁵⁻¹⁷ Chaurasia et al¹⁵ described patients of MCD who at an early age had co-existing anterior stromal lesions in the central cornea and non-contiguous deep stromal pre-Descemet lesions in the peripheral cornea. An unusual case of a 67-year-old female with MCD who had predominantly deep stromal lesions in the peripheral cornea with sparse anterior stromal deposits is recognised.¹⁶ Another case report of MCD with isolated peripheral Descemet membrane deposits has been reported by Zhang et al.¹⁷ The anterior stromal and subepithelial deposits affect the visual acuity due to an altered corneal surface and irregular astigmatism. Recurrent corneal erosions can occur.¹⁸ One of the important features characteristic of MCD is progressive corneal thinning which is because of the decreased inter-fibrillar spacing.^{3,19} However, progression of deposits in the endothelial cells can eventually lead to thickening of the Descemet membrane, endothelial decompensation and increased pachymetry.^{6,15,16}

PATHOGENESIS

MCD appears to be the result of a metabolic abnormality in keratan sulfate.²⁰ Keratan sulfate is found in the corneal epithelium, Bowman's membrane, keratocytes, Descemet membrane and endothelium. Carbohydrate sulfotransferase (*CHST6*) gene on chromosome 16(16q22) gene is explained to be important in producing sulfated keratan sulfate which is important glycosaminoglycan in the adult cornea. *CHST6* gene encodes N-acetylglucosamine 6-O-sulfotransferase, an enzyme which transfers sulfate to the unsulfated keratin chains on lumican.^{21,22} Lumican preserves the crucial size, ordered structure, impacts corneal hydration and hence the corneal transparency. This explains the stromal haze and the loss of corneal transparency due to mutation in *CHST6* gene in MCD.

Initially, it was believed that the metabolic abnormality of keratan sulfate was restricted to cornea alone. As keratan sulfate is derived from the normal turnover of cartilage, studies have looked at the keratan sulfate levels in cornea, serum and other cartilage samples of the body.²³ Klintworth et al in one their studies reported that serum assays of patients with MCD showed absence of sulphated keratan sulfate.²⁰ This pointed towards the defects in the synthesis of keratan sulfate in MCD. Thonar et al²⁴ showed that the levels of keratan sulfate were higher in children of age group 13-15 years and patients with osteoarthritis. They also showed decreased levels of keratan sulfate in serum of patients with MCD, deriving that there is a possibility of systemic manifestation in patients with MCD.

Depending on the reactivity of keratan sulfate in the serum and cornea to anti-keratan sulfate antibodies, MCD is categorised into 3 variants⁵:

1. MCD type I-Keratan sulfate is absent in both the serum and the cornea. Corneal epithelium, stroma and endothelium all of them contain unsulfated keratan
2. MCD type IA-Keratan sulfate is absent in the serum but stroma shows immunoreactivity to Keratan sulfate antibodies
3. MCD type 2-Keratan sulfate levels are decreased in serum and corneal stroma. But epithelium and endothelium show unsulfated keratan

Clinically and histopathologically, the three variants are indistinguishable.^{25,26}

HISTOLOGY AND ULTRASTRUCTURAL CHARACTERSTICS

Histopathologic features of MCD is characteristic (Figure 1a-c). The stromal keratocytes, Descemet membrane and endothelial cells show intra and extracellular deposition of granular material.^{3,6} Bowman's membrane can be absent, disrupted and thinned out. Epithelium invariably remains spared of the deposits. The depositions which are due to abnormally sulphated keratin sulphate appear as basophilic granular material. Granular material is

chemically acid mucopolysaccharide (glycosaminoglycans) and stains positively with periodic acid Schiff, alcian blue, metachromatic dyes and colloidal iron.²⁷ Electron microscopy substantiates this finding with intracellular deposition of glycosaminoglycans within the endoplasmic reticulum of stromal keratocytes especially superficial keratocytes.²⁸ Numerous electron- transparent lacunae are randomly distributed throughout MCD corneas. The abnormal keratan sulphate interferes with the collagen fibrillar rearrangement. The collagen fibrils may have a normal diameter but the interfibrillar spacing of collagen fibrils in affected corneas is less than that in the normal cornea.³ Some studies have reported that the collagen fibrils are smaller with a decrease in the interfibrillar spacing.²⁹ This close packing of collagen fibrils is responsible for the reduced corneal thickness in MCD.

The anterior banded portion of Descemet membrane is of normal thickness and has an unremarkable ultrastructure, whereas the posterior layer usually contains numerous corneal guttae with granular deposits within the corneal endothelium.^{3,6} This indicates that the abnormal glycosaminoglycan is not produced by the fetal endothelial cells. Snip et al⁶ reported that Descemet membrane acts like a physical barrier and it is unlikely for the fibrogranular material in the vacuoles of endothelium to be a phagocytosed product of granular material deposited in the stroma and suggested that the material is derived from the endothelium suggesting that the involvement of corneal endothelium may be primary feature of the disease.

One of the close differential diagnosis of MCD is mucopolysaccharidoses (MPS).³⁰ Unlike the sparing of the epithelium that is seen in MCD, systemic mucopolysaccharidoses shows pan-corneal involvement with glycosaminoglycans. Another differentiating feature is that the site of deposition in MCD is rough endoplasmic reticulum whereas in MPS, it is in Golgi apparatus.

GENETICS

Genetics of a dystrophy is important in understanding the underlying basis of the disease and may also help in the development of gene therapy or other targeted therapies for the disease. The involvement of keratan sulphate in the pathology of MCD was evident from biochemical studies of MCD corneas using organ culture of corneas of patients affected with MCD and those of unaffected controls.³¹ The defect in MCD appeared due to two disturbances. There is an absence of sulfate residues in the carbohydrate side chains from keratan sulfate proteoglycan and secondly the oligosaccharide side chains are much smaller in MCD-affected corneas than keratan sulfate side chains present in normal corneas.

MCD was linked to human chromosome number 16 in families of American and Icelandic origin.³² Akama et al^{33,34} identified mutations in the gene for *CHST6* among patients with both the types of MCD. MCD type I is the predominant type and involves mutations of the coding regions of *CHST6* whereas in MCD type II, deletions and rearrangements have been found in the upstream region of the gene. Various studies show that MCD patients from different regions of the world have mutations in the same gene.^{22,35-37} There are many mutant alleles identified so far indicating allelic heterogeneity in MCD. Data from about 200 families from across the world analysed for mutations in *CHST6* show that approximately half of all MCD patients reported so far are from India. There is a predominance of missense mutations, detected in about 50% or more of patients with MCD from different populations while nonsense mutations, deletions, insertions or indels are found in about one-third of cases.³⁸ Being an autosomal recessive disorder, MCD is mostly seen in consanguineous and inbred families, in which the majority of patients are homozygous for the mutations, and rest are compound heterozygotes for two different alleles.³⁹ There are no obvious correlations

between genotypes and phenotypes of patients with MCD, either the immunophenotypes or clinical features.^{40,41}

INVESTIGATIONS

Anterior segment optical coherence tomography (OCT)

High Resolution OCT shows hyperreflectivity at the site of deposits associated with thinning of the epithelium over the deposits.⁴² Line scans passing through the peripheral cornea characterise the pre-Descemet nature of the peripheral deposits.¹⁵ In some patients, thickening of the Descemet membrane may be noted. In advanced stages of the dystrophy, the dense stromal deposits cause an optical shadow in the posterior part of cornea.

In vivo confocal microscopy

In vivo confocal microscopy showed altered reflectivity of basal epithelial cells, hyperreflectivity in the anterior stroma, granular appearance of keratocytes and extracellular matrix, and dark striae in the middle and posterior stroma. The endothelial cells showed bright granules and polymegathism.⁴³ In one study, normal endothelium was reported.⁴⁴

Scheimpflug imaging

There are few reports of coexistence of keratoconus and MCD.⁴⁵ Several studies have shown thinning of the central cornea in MCD. Dudakova et al⁴⁶ used pentacam scheimpflug imaging system in patients with MCD and concluded that all eyes showed diffuse corneal thinning with paracentral steepening of the anterior corneal surface that was graded as keratoconus by the integrated software, but without associated ectasia of the posterior corneal surface or regional thinning.

Ultrasound bio microscopy (UBM)

Rubinstein et al⁴² used UBM in the evaluation of patients with MCD that could help to examine important posterior lamellar changes that could help to decide the choice of keratoplasty- full thickness versus lamellar keratoplasty. UBM helped to characterise posterior corneal surface interruptions, loss of continuity and focal protrusions. Penetrating keratoplasty (PK) could be the surgery of choice compared to deep anterior lamellar keratoplasty (DALK) in patients with involvement of the posterior corneal layers.

MANAGEMENT

Phototherapeutic keratectomy (PTK)

Excimer laser has been used for removal of superficial opacities. Corneal dystrophies are the common indications of PTK. PTK helps to improve vision as well as those who suffer from recurrent erosion. Hafner et al⁴⁷ had described long-term result of PTK in 10 eyes of six patients with MCD. They concluded that PTK can improve visual acuity moderately for limited period. They noticed high recurrence rate (90%) during follow-up. Mitomycin (0.02%) has been used for 30 seconds on stromal bed after PTK for granular and macular dystrophies.⁴⁸ There was no difference in recurrence rate and recurrence free interval between both. Many patients need keratoplasty despite PTK due to recurrence leading to reduction of vision. Preceding PTK does not impair outcome of subsequent keratoplasties.⁴⁹ PTK can be performed in recurrence of primary pathology in graft.⁵⁰ The advantages of PTK are repeatability of the procedure and precise ablation. However, it induces hyperopic shift and stromal haze.

Femtosecond assisted lamellar keratectomy

Femtosecond LASER has been successfully used for anterior lamellar keratectomy to remove corneal opacities secondary to anterior corneal dystrophies.^{51,52} It may delay more invasive surgery like keratoplasty. Corneal flap of 9.5 mm diameter was created. Corneal flap

thickness was around 110 to 140 microns.⁵² Best-spectacle-corrected-visual acuity improved after the procedure. There was no hyperopic shift.

Keratoplasty

MCD constitutes a major portion of transplants performed in many places.^{8,10,53} Penetrating keratoplasty was standard of care in the earlier days. Al-Swailemet al have described penetrating keratoplasty in 229 eyes of 141 patients.⁵⁴ Probability of graft survival was 98.1% at 1 year, 89.8% at 5 years, 82.1% at 10 years and 74.1% at 15 years. They found the risk of graft failure is more if patient was < 40 years during surgery. There was clinically significant recurrence in 5.2% cases after a mean interval of 84 ± 48.2 months. Karimian et al have described visually insignificant recurrence in one out of 62 eyes.⁵⁵ Though 19.4% eyes had immunological rejection, none of them failed.

Few studies compare outcome of DALK with PK in MCD (**Table-1**). In a prospective randomised controlled trial by Sari et al⁵⁶, there was no significant difference in final visual outcome between both groups. There is significant more endothelial cell loss in PK compared to DALK. Cheng et al had reported better visual outcome in PK group compared to DALK group.⁵⁷ However, endothelial cell loss was more in PK group. Reddy et al compared clinical outcome and risk factors of graft failure in both groups in a retrospective analysis.⁵⁸ There was no significant difference in visual outcome, astigmatism and spherical equivalent. Graft rejection was the main cause of graft failure in PK group. Intraoperative micro perforation and double anterior chamber was common complications of DALK. While most complications of DALK happened intraoperatively and early postoperative period, late phase complications such as rejection and secondary glaucoma are more in PK group.⁵⁹ Late endothelial decompensation after DALK was reported in 2 patients in the same study.⁵⁹

Endothelial cell loss and recurrence in MCD after DALK has been more compared to Lattice dystrophy who had undergone DALK.^{59,60} Kawashima et al⁵ reported increased endothelial cell loss after DALK in macular corneal dystrophy in comparison to lattice corneal dystrophy. Recurrence of primary pathology is less compared to lattice and granular dystrophy after DALK.⁶⁰ Primary graft failure has been reported after DALK.⁶¹ Descemet stripping endothelial keratoplasty is described for an atypical case of MCD, wherein glycosaminoglycans deposits were predominantly at the endothelial level.¹⁶

FUTURE PROSPECTS

Over last several years, there is some evidence of successful gene transfer to the cornea in experimental models and in human corneas ex vivo.⁶²⁻⁶⁴ MCD has an identified and mapped genetic defect and hence, fulfils the important prerequisites that are required to attempt a clinical gene therapy. The conventional gene therapy involves transfer of a new/ replacement gene to the cells to drive its expression or to silence the expression of the abnormal gene.⁶⁵ Genome editing is a therapy where the defective gene is repaired or deleted, i.e. the host genome is 'edited'.

The cornea is a suitable organ for attempting gene replacement or correction of affected cells, since it is easily accessible, its immune privilege and limited size. Correction of specific mutations in the *CHST6* gene may be potentially developed by means of patient-specific gene editing approaches. An alternative that may be applicable to MCD, is that since it is an autosomal recessive disease, it is amenable to replacement of the entire gene by gene therapy with suitable viral vectors such as adeno-associated virus (AAV) that have been tried and tested in other gene therapy trials in humans. Several methods of gene editing exist such as using zinc finger nucleases (ZFNs) or artificial restriction enzymes [(transcription activator- like effector nucleases (TALENs)]. Currently, Clustered regularly interspaced

short palindromic DNA repeat (or CRISPR) and CRISPR- associated genes (cas) system of gene editing have seen increased interest and these may emerge as future therapies as it is capable of correcting gene effects in the somatic cells of a whole organism.⁶⁶ Such an approach was employed in an experimental model for correction of the keratin 12 gene defect in a mouse model of Meesmann corneal dystrophy and also in limbal epithelial cells of patients.⁶⁷

CONCLUSION

MCD is an autosomal recessive disorder which affects young individuals. The deposition of abnormal protein lead to corneal opacification. Imaging techniques help to understand ultrastructural changes in cornea. Various modalities of treatment such as PK and DALK have been described for visual rehabilitation. The success of DALK depends upon the endothelial health. Recurrence can occur after keratoplasty. Newer therapies like gene transfer and gene editing may be a potential treatment in the future.

METHOD OF LITERATURE SEARCH

PubMed search was performed using search terms: macular corneal dystrophy, genetics, histopathology, investigations, and management of MCD. All articles published in English language and relevant to the topic were reviewed and included in this review.

Contributors: All the authors fulfil the criteria of authorship.

Funding: Hyderabad Eye Research Foundation, Hyderabad

Competing interests: None

Ethics approval: The Institutional Review board of LV Prasad Eye Institute, KAR campus, Hyderabad, Telangana approved the study.

REFERENCES

1. Duke Elder S. System of Ophthalmology. Vol 8. 1965:921.
2. Born AJ. Genetics of the corneal dystrophies: what we have learned in the past twenty-five years. *Cornea* 2000; 19:699-711.
3. Klintworth GK. Corneal dystrophies. *Orphanet J Rare Dis.* 2009; 4:7
4. Weiss JS, Møller HU, Lisch W, Kinoshita S, Aldave AJ, Belin MW, Kivelä T, Busin M, Munier FL, Seitz B, et al. The IC3D classification of the corneal dystrophies. *Cornea.* 2008; 27 Suppl 2:S1-83.
5. Weiss JS, Moller HU, Aldave AJ, Seitz B, Bredrup C, Kivelä T, Munier FL, Rapuano CJ, Nischal KK, Kim EK, et al. IC3D classification of corneal dystrophies edition 2. *Cornea.* 2015; 34(2):117-59.
6. Snip RC, Kenyon KR, Green WR. Macular corneal dystrophy: Ultrastructural pathology of corneal endothelium and Descemet's membrane. *Investigative Ophthalmology* 1973; Vol 12(No 2):88-97.
7. Agrawal S, Peck T, Golen J, Karcioğlu ZA. Macular Corneal Dystrophy: A Review. *Survey Ophthalmology*; 2018; 63:609-617.
8. Jonasson F, Johannsson JH, Garner A, Rice NS. Macular Corneal dystrophy in Iceland. *Eye (Lond).* 1989; 3(Pt 4):446-54.
9. Al Faran MF, Tabbara KF. Corneal dystrophies among patients undergoing keratoplasty in Saudi Arabia. *Cornea* 1991; 10(1):13-6.
10. Pandrowala H, Bansal A, Vemuganti GK, Rao GN. Frequency, distribution, and outcome of keratoplasty for corneal dystrophies at a tertiary eye care center in South India. *Cornea.* 2004; 23(6):541-6.

11. Klintworth GK. Macular corneal dystrophy-a localised disorder of mucopolysaccharide metabolism. Clinical, structural and biochemical advances in hereditary eye disorders. New York, Alan R, Liss Inc; 1982.pp69-101.
12. Møller HU, Sunde L. Prevalence of corneal dystrophies in the United States: estimates from claims data. Invest Ophthalmol Vis Sci. 2013; 14:54(1).
13. Alzuhairy S, Alkatan HM, Al-Rajhi AA. Prevalence and histopathological characteristics of corneal stromal dystrophies in Saudi Arabia. Middle East Afr J Ophthalmol. 2015;22(2):179-85
14. Das AV, Chaurasia S. Clinical Profile and Demographic Distribution of Corneal Dystrophies in India: A Study of 4198 Patients. Cornea 2020 Jul 30. (ahead of print)
15. Chaurasia S, Ramappa M, Mishra DK. Clinical diversity in macular corneal dystrophy: an optical coherence tomography study. Int Ophthalmol. 2019;39(12):2883-2888.
16. Chaurasia S, Mishra DK. Atypical presentation of macular corneal dystrophy managed by Descemet stripping endothelial keratoplasty. Indian J Ophthalmol. 2019; 67(1):118-119.
17. Zhang W, Kassels AC, Barrington A, Khan S, Tomatsu S, Alkadi T, Aldave A. Macular corneal dystrophy with isolated peripheral Descemet membrane deposits. Am J of Ophthalmology Case Reports 2019; 16:1- 4.
18. Ewald M, Hammersmith KM. Review of diagnosis and management of recurrent erosion syndrome. Curr Opin Ophthalmol. 2009; 20(4):287-291.
19. Donnenfeld ED, Cohen EJ, Ingraham HJ, Poleski SA, Goldsmith E, Laibson PR. Corneal thinning in macular corneal dystrophy. Am J Ophthalmol 1986; 101: 112-113.

20. Klintworth GK. Research into the pathogenesis of macular corneal dystrophy. *Trans Ophthalmol Soc UK* 1980, 100: 186-94.
21. Klintworth GK, Smith CF, Bowling BL. CHST6 mutations in North American subjects with macular corneal dystrophy: a comprehensive molecular genetic review. *Mol Vis* 2006;12:159-76.
22. Liu NP, Smith CF, Bowling BL, Jonasson F, Klintworth GK. Macular corneal dystrophy types I and II are caused by distinct mutations in the CHST6 gene in Iceland. *Mol Vis* 2006; 12:1148-52.
23. Lewis D, Davies Y, Nie duszynski IA, Lawrence F, Quantock AJ, Bonshek R, Fullwood NJ. Ultrastructural localization of sulphated and unsulfated keratan sulfate in normal and macular corneal dystrophy type I. *Glycobiology* 2000; 10(3):305-12.
24. Thonar EJ, Lenz ME, Klintworth GK, Caterson B, Pachman LM, Glickman P, Katz R, Huff J, Kuettner KE. Quantification of keratan sulfate in blood as a marker of cartilage catabolism. *Arthritis Rheuma* 1985; 28 (12):1367-76.
25. Edward DP, Yue BYJT, Sugar J, Thonar EJ, SunderRaj N, Stock EL, Tso MO. Heterogeneity in Macular Corneal Dystrophy. *Arch Ophthalmol*. 1988; 106 (11):1579–1583.
26. Klintworth GK, Smith CF. Macular corneal dystrophy. Studies of sulphated glycosaminoglycans in corneal explant and confluent stromal cell cultures. *Am J Pathol*. 1977;89(1):167-82.
27. Klintworth GK, Vogel FS. Macular corneal dystrophy: An inherited acid mucopolysaccharide storage disease of the corneal fibroblast. *Am J Pathol*. 1964; 45:565–586.
28. Meek KM, Quantock AJ, Elliott GF, Ridgway AE, Tullo AB, Bron AJ, Thonar EJ. Macular corneal dystrophy: the macromolecular structure of the stroma observed

using electron microscopy and synchrotron X-ray diffraction. *Exp Eye Res.* 1989; 49(6):941–958.

29. Akhtar S, Alkatan HM, Kirat O, Khan AA, Almubrad T. Collagen Fibrils and Proteoglycans of Macular Dystrophy Cornea: Ultrastructure and 3D Transmission Electron Tomography. *Microsc Microanal.* 2015; 21(3):666–679.
30. Ashworth JL, Biswas S, Wraith E, Lloyd IC. Mucopolysaccharidoses and the eye. *Surv Ophthalmol* 2006; 51:1–17.
31. Hassell JR, Newsome DA, Krachmer JH, Rodrigues MM. Macular corneal dystrophy: failure to synthesize a mature keratan sulfate proteoglycan. *Proc Natl Acad Sci U S A.* 1980;77(6):3705-9.
32. Vance JM, Johansson F, Lennon F, Sarrica J, Damji KF, Stauffer J, Pericak-Vance MA, Klintworth GK. Linkage of a gene for macular corneal dystrophy to chromosome 16. *Am J Hum Genet* 1996;58(4):757- 62.
33. Akama TO, Nakayama J, Nishida K, Hiraoka N, Suzuki M, McAuliffe J, Hindsgaul O, Fukuda M, Fukuda MN. Human corneal GlcNac 6-O-sulfotransferase and mouse intestinal GlcNac 6- O-sulfotransferase both produce keratan sulfate. *J Biol Chem.* 2001;276(19):16271-8.
34. Akama TO, Nishida K, Nakayama J, Watanabe H, Ozaki K, Nakamura T, Dota A, Kawasaki S, Inoue Y, Maeda N, Yamamoto S, Fujiwara T, Thonar EJ, Shimomura Y, Kinoshita S, Tanigami A, Fukuda MN. Macular corneal dystrophy type I and type II are caused by distinct mutations in a new sulphotransferase gene. *Nat Genet.* 2000;26(2):237-41.
35. Sultana A, Sridhar MS, Jagannathan A, Balasubramanian D, Kannabiran C, Klintworth GK. Novel mutations of the carbohydrate sulfotransferase-6 (CHST6) gene causing macular corneal dystrophy in India. *Mol Vis.* 2003; 9:730-734.

36. El-Ashry MF, Abd El-Aziz MM, Wilkins S, Cheetham ME, Wilkie SE, Hardcastle AJ, Halford S, Bayoumi AY, Ficker LA, Tuft S, et al. Identification of novel mutations in the carbohydrate sulfotransferase gene (CHST6) causing macular corneal dystrophy. *Invest Ophthalmol Vis Sci*. 2002; 43(2):377-382.
37. Aldave A, Yellore VS, Thonar EJ, Udar N, Warren JF, Yoon MK, Cohen EJ, Rapuano CJ, Laibson PR, Margolis TP. Novel mutations in the carbohydrate sulfotransferase gene (CHST6) in American patients with macular corneal dystrophy. *Am J Ophthalmol*. 2004; 137(3):465-473.
38. Sultana A, Sridhar MS, Klintworth GK, Balasubramanian D, Kannabiran C. Allelic heterogeneity of the carbohydrate sulfotransferase-6 gene in patients with macular corneal dystrophy. *Clin Genet*. 2005; 68(5):454-460.
39. C. Kannabiran. Genetics of Eye Diseases. Springer Nature Singapore Pte Ltd. 2019. doi-https://doi.org/10.1007/978-981-13-7146-2_1
40. Sultana A, Klintworth GK, Thonar EJ, Vemuganti GK, Kannabiran C. Immunophenotypes of macular corneal dystrophy in India and correlation with mutations in CHST6. *Mol Vis*. 2009; 15:319-325.
41. Nowinska AK, Wylegala E, Teper S, Wróblewska-Czajka E, Aragona P, Roszkowska AM, Micali A, Pisani A, Puzzolo D. Phenotype and genotype analysis in patients with macular corneal dystrophy. *Br Journal of Ophthalmology* 2014; 98:1514-1521.
42. Rubinstein Y, Weiner C, Einan-Lifshitz A, Chetrit N, Shoshany N, Zadok D, Avni I, Pras E. Macular Corneal Dystrophy and Posterior Corneal Abnormalities. *Cornea*. 2016; 35(12):1605-1610.
43. Micali A, Pisani A, Puzzolo D, Nowińska A, Wylegala E, Teper S, Czajka E, Roszkowska AM, Orzechowska-Wylegala B, Aragona P. Macular corneal dystrophy: in vivo confocal and structural data. *Ophthalmology*. 2014; 121(6):1164-1173.

44. Kobayashi A, Fujiki K, Fujimaki T, Murakami A, Sugiyama K. In vivo laser confocal microscopic findings of corneal stromal dystrophies. *Arch Ophthalmol.* 2007; 125(9):1168-1173.
45. Javadi MA, Rafee AB, Kamalian N, Karimian F, Ja'farinasab MR, Yazdani S. Concomitant keratoconus and macular corneal dystrophy. *Cornea* 2004; 23(5):508-12.
46. Dudakova L, Palos M, Svobodova M, Bydzovsky J, Huna L, Jirsova K, Hardcastle AJ, Tuft SJ, Liskova P. Macular corneal dystrophy and associated corneal thinning. *Eye (Lond).* 2014; 28(10):1201-1205.
47. Hafner A, Langenbucher A, Seitz B. Long-term results of phototherapeutic keratectomy with 193-nm excimer laser for macular corneal dystrophy. *Am J Ophthalmol.* 2005; 140(3):392-396.
48. Yuksel E, Cubuk MO, Eroglu HY, Bilgihan K. Excimer laser phototherapeutic keratectomy in conjunction with mitomycin C in corneal macular and granular dystrophies. *Arq Bras Oftalmol.* 2016; 79(2):69-72.
49. Szentmáry N, Langenbucher A, Hafner A, Seitz B. Impact of phototherapeutic keratectomy on the outcome of subsequent penetrating keratoplasty in patients with stromal corneal dystrophies. *Am J Ophthalmol.* 2004; 137(2):301-307.
50. Reddy JC, Rapuano CJ, Nagra PK, Hammersmith KM. Excimer laser phototherapeutic keratectomy in eyes with corneal stromal dystrophies with and without a corneal graft. *Am J Ophthalmol.* 2013; 155(6):1111-1118.
51. Femtosecond Laser-assisted lamellar keratectomy for corneal opacities secondary to anterior corneal dystrophies: An interventional case series. *Cornea* 2016; 35:6-13.
52. Tsagkataki M, Tey A, Ramasamy B, Anijeet D, Kaye SB. Femtosecond laser-assisted lamellar keratectomy for corneal opacities due to macular dystrophy: an interventional case report. *Clin Exp Ophthalmol.* 2014; 42(3):288-289.

53. Klintworth GK, Oshima E, al-Rajhi A, al-Saif A, Thonar EJ, Karcioğlu ZA. Macular corneal dystrophy in Saudi Arabia: a study of 56 cases and recognition of a new immunophenotype. *Am J Ophthalmol*. 1997; 124(1):9–18.
54. Al-Swailem SA, Al-Rajhi AA, Wagoner MD. Penetrating keratoplasty for macular corneal dystrophy. *Ophthalmology*. 2005; 112(2):220–224.
55. Karimian F, Baradaran-Rafii AR, Feizi S, Zare M, Jafarinasab MR, Javadi MA, Mirdehghan SA, Einollahi B. Outcomes of penetrating keratoplasty for macular corneal dystrophy. *J Ophthalmic Vis Res*. 2009; 4(1):14–18.
56. Sogutlu Sari E, Kubaloglu A, Unal M, Pinero D, Bulut N, Erol MK, Özürtürk Y. Deep anterior lamellar keratoplasty versus penetrating keratoplasty for macular corneal dystrophy: a randomized trial. *Am J Ophthalmol*. 2013;156(2):267–274.
57. Cheng J, Qi X, Zhao J, Zhai H, Xie L. Comparison of penetrating keratoplasty and deep lamellar keratoplasty for macular corneal dystrophy and risk factors of recurrence. *Ophthalmology*. 2013; 120(1):34–39.
58. Reddy JC, Murthy SI, Vaddavalli PK, Garg P, Ramappa M, Chaurasia S, Rath V, Sangwan VS. Clinical outcomes and risk factors for graft failure after deep anterior lamellar keratoplasty and penetrating keratoplasty for macular corneal dystrophy. *Cornea*. 2015; 34(2):171–176.
59. Kawashima M, Kawakita T, Den S, Shimmura S, Tsubota K, Shimazaki J. Comparison of deep lamellar keratoplasty and penetrating keratoplasty for lattice and macular corneal dystrophies. *Am J Ophthalmol*. 2006; 142(2):304–309.
60. Unal M, Arslan OS, Atalay E, Mangan MS, Bilgin AB. Deep anterior lamellar keratoplasty for the treatment of stromal corneal dystrophies. *Cornea*. 2013;32(3):301–305.

61. Chaurasia S, Ramappa M, Murthy S, Garg P. Primary graft failure after big bubble deep anterior lamellar keratoplasty in macular corneal dystrophy. *Indian J Ophthalmol.* 2018; 66(8):1196-1197.
62. Klauser EA, Peer D, Chapman RL, Multack R, Andurkar SV. Corneal gene therapy. *J Control Release.* 2007; 124:107-133.
63. Kampik D, Ali RR, Larkin DF. Experimental gene transfer to the corneal endothelium. *Exp Eye Res.* 2012; 23:276-279.
64. Williams KA, Klebe S. Gene therapy for corneal dystrophies and disease, where are we? *Curr Opin Ophthalmol.* 2012; 23:276-279.
65. Williams KA, Irani YD. Gene therapy and gene editing for the corneal dystrophies. *Asia Pacific Journal of Ophthalmology.* 2016.
66. Byrne SM, Mali P, Church GM. Genomic editing in human stem cells. *Methods Enzymol.* 2014; 546:119-138.
67. Courtney DG, Moore JE, Atkinson SD, Maurizi E, Allen EHA, Pedrioli DML, McLean WHI, Nesbit MA, Moore CBT. CRISPR/Cas9 DNA cleavage at SNP-derived PAM enables both in vitro and in vivo KRT12 mutation-specific targeting. *Gene Ther.* 2016; 23(1):108-12.

Table -1: Modality of treatment in MCD

Study	Number of eyes (patients)	Intervention	Mean follow up	Recurrence	Recurrence free period
Hafner A et al ⁴⁷	10	PTK	4.5±3.1 years	90%	3.4±0.4 years
Yuksel et al ⁴⁸	9	PTK+MMC	-----	55.5%	5.6±1.4months
Al Swalim et al ⁵⁴	229(141)	PK	5.9±3.8years	5.2%	84± 48.2 months
Karimian et al ⁵⁵	62(39)	PK	52 ± 47.3months	1 case (visually insignificant)	-----
Sari SE et al ⁵⁶	82 (54)	PK vs. DALK	-----	4.8% PK 5.7% DALK	-----
Cheng et al ⁵⁷	78(51)	PK vs. DALK	5.1±4.1 years	17.5%PK 42.9%DALK	6.3±3.1 years in PK 2.3±2.1 years in DALK

Figure 1(a-c): a.) Fleck like deposits seen in macular dystrophy. b.) Histology of MCD showing thickened Descemet membrane and excrescences (guttae). c.) Alcian blue positive glycosaminoglycan deposits in the stroma.

ACCEPTED MANUSCRIPT




Fig 1