[CLS] Transmaternal Helicobacter pylori exposure reduces allergic airway inflammation in offspring through regulatory T cells BACKGROUND Transmaternal exposure to tobacco, microbes, nutrients, and other environmental factors shapes the fetal immune system through epigenetic processes. The gastric microbe Helicobacter pylori represents an ancestral constituent of the human

microbiota that causes gastric disorders on the one hand and is inversely associated with allergies and chronic inflammatory conditions on the other. OBJECTIVE Here we investigate the consequences of transmaternal exposure to H pylori in utero and / or during lactation for susceptibility to viral and

bacterial infection, predisposition to allergic airway inflammation, and development of immune cell populations in the lungs and lymphoid organs. METHODS We use experimental models of house dust mite - or ovalbumin - induced airway inflammation and influenza A virus or Citrobacter rodentium infection along with metagenomics analyses, multicolor flow cytometry, and bisulfite pyrosequencing, to study the effects of H pylori on allergy severity and immunologic and microbiome correlates thereof. RESULTS Perinatal exposure to

H pylori extract or its immunomodulator vacuolating cytotoxin confers robust protective effects against allergic airway inflammation not but does not increase susceptibility to viral or bacterial infection. Immune correlates of allergy protection include skewing of regulatory over effector T cells, expansion of regulatory T - cell orphan receptor t, and demethylation of the

only in first - but also second - generation offspring

subsets expressing CXCR3 or retinoic acid - related forkhead box P3 (FOXP3) locus. The composition and diversity of the gastrointestinal microbiota is measurably affected by perinatal H pylori exposure. CONCLUSION We conclude that exposure to H pylori has consequences not only for the carrier but

also for subsequent generations that can be exploited for interventional purposes. [SEP]