

Inference for multivariate time series models: a measure transportation approach

Davide La Vecchia
(joint work with M. Hallin & H. Liu)

Cyprus, Nov-2021

I am going to explain how **measure transportation theory** can be applied to derive **novel semiparametric inference procedures for multivariate time series**.

I am going to explain how **measure transportation theory** can be applied to derive novel **semiparametric inference procedures** for multivariate time series.

Features

The resulting techniques:

- *are useful to conduct inference on semiparametric models, involving an Euclidean parameter θ and an infinite nuisance dimensional parameter f*

I am going to explain how **measure transportation theory** can be applied to derive **novel semiparametric inference procedures** for multivariate time series.

Features

The resulting techniques:

- *are useful to conduct inference on semiparametric models, involving an Euclidean parameter θ and an infinite nuisance dimensional parameter f*
- *are based on novel concepts of multivariate ranks, quantiles and signs and on a reference density (user selected and possibly misspecified)*

I am going to explain how **measure transportation theory** can be applied to derive **novel semiparametric inference procedures** for multivariate time series.

Features

The resulting techniques:

- *are useful to conduct inference on semiparametric models, involving an Euclidean parameter θ and an infinite nuisance dimensional parameter f*
- *are based on novel concepts of multivariate ranks, quantiles and signs and on a reference density (user selected and possibly misspecified)*
- *remain valid (root- n consistent estimators and well-sized tests) for a large class of actual densities and improve (efficiency/power) on routinely-applied Gaussian procedures*

I am going to explain how **measure transportation theory** can be applied to derive **novel semiparametric inference procedures for multivariate time series**.

Features


The resulting techniques:

- *are useful to conduct inference on semiparametric models, involving an Euclidean parameter θ and an infinite nuisance dimensional parameter f*
- *are based on novel concepts of multivariate ranks, quantiles and signs and on a reference density (user selected and possibly misspecified)*
- *remain valid (root-n consistent estimators and well-sized tests) for a large class of actual densities and improve (efficiency/power) on routinely-applied Gaussian procedures*

Talk based on

- (HLLa) Hallin, La Vecchia & Liu, Center-outward R-estimation for semiparametric VARMA models, JASA, accepted in 2020.
- (HLLb) Hallin, La Vecchia & Liu, Rank-based testing for semiparametric VAR models: a measure transportation approach, (2021), working paper.

For R-estimation, <https://github.com/HangLiu10/RestVARMA>. Other related codes available on my [GitHub](#) that you may join via my website:



Outline

- Monge-Kantorovich problem: measure transportation
- Motivation
 - ▶ Empirical and theoretical issues
 - ▶ Overview of our solution: center-outward ranks and signs estimators and tests
- The method at a glance:
 - ▶ Mapping the residuals to the unit ball: the measure transportation approach
 - ▶ Center-outward ranks and signs
 - ▶ Sketch of the construction for R-estimators and tests
- A real-data example
- Take home message

Looking at the issue of finding the best way to move given piles of sand to fill up given holes of the same total volume, **Gaspard Monge** (1746-1818) formulated a **mathematical problem** that in modern jargon reads as:

Looking at the issue of finding the best way to move given piles of sand to fill up given holes of the same total volume, **Gaspard Monge** (1746-1818) formulated a **mathematical problem** that in modern jargon reads as:

Let α and β denote two probability measures over (for simplicity) $(\mathbb{R}^d, \mathcal{B}^d)$, for $d \geq 1$. Let $c : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ be a Borel-measurable cost function such that $c(\mathbf{x}, \mathbf{y})$ represents the cost of transporting \mathbf{x} to \mathbf{y} . Then, find a measurable transport map $\mathcal{T} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ that achieves

$$\inf_{\mathcal{T} \in M} \int_{\mathbb{R}^d} c[\mathbf{x}, \mathcal{T}(\mathbf{x})] d\alpha \quad (1)$$

where

$$M := \{\mathcal{T} : \mathbf{X} \rightarrow \mathbf{Y}\},$$

with $\mathbf{X} \sim \alpha$, $\mathbf{Y} \sim \beta$. The map $\mathcal{T} \# \alpha = \beta$ does the “push forward of α to β ”.

Looking at the issue of finding the best way to move given piles of sand to fill up given holes of the same total volume, **Gaspard Monge** (1746-1818) formulated a **mathematical problem** that in modern jargon reads as:

Let α and β denote two probability measures over (for simplicity) $(\mathbb{R}^d, \mathcal{B}^d)$, for $d \geq 1$. Let $c : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ be a Borel-measurable cost function such that $c(\mathbf{x}, \mathbf{y})$ represents the cost of transporting \mathbf{x} to \mathbf{y} . Then, find a measurable transport map $\mathcal{T} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ that achieves

$$\inf_{\mathcal{T} \in M} \int_{\mathbb{R}^d} c[\mathbf{x}, \mathcal{T}(\mathbf{x})] d\alpha \quad (1)$$

where

$$M := \{\mathcal{T} : \mathbf{X} \rightarrow \mathbf{Y}\},$$

with $\mathbf{X} \sim \alpha$, $\mathbf{Y} \sim \beta$. The map $\mathcal{T} \# \alpha = \beta$ does the “push forward of α to β ”.

⇒ The map solution to (1) is called the optimal transportation map.

Monge's problem remained open until the 1940s, when it was revisited by **Leonid Vitaliyevitch Kantorovich** (1912-1986; Nobel Prize in Economics in 1975) for the economic problem of optimal allocation of resources; see e.g. **Villani (2008)**, **Santambrogio (2015)**, **Galichon (2016)**.

Monge's problem remained open until the 1940s, when it was revisited by **Leonid Vitaliyevitch Kantorovich** (1912-1986; Nobel Prize in Economics in 1975) for the economic problem of optimal allocation of resources; see e.g. **Villani (2008)**, **Santambrogio (2015)**, **Galichon (2016)**.

In the **Kantorovich primal problem**, the objective is to find the **optimal transportation plan** γ , which solves

$$\inf_{\gamma \in \Gamma(\alpha, \beta)} \int_{\mathbb{R}^d \times \mathbb{R}^d} c(\mathbf{x}, \mathbf{y}) d\gamma(\mathbf{x}, \mathbf{y}), \quad (2)$$

where the infimum is over all coupling (\mathbf{X}, \mathbf{Y}) of (α, β) , belonging to $\Gamma(\alpha, \beta)$, the set of probability measures γ on $\mathbb{R}^d \times \mathbb{R}^d$, satisfying

$$\gamma(A \times \mathbb{R}^d) = \alpha(A) \text{ and } \gamma(\mathbb{R}^d \times B) = \beta(B),$$

for measurable sets $A, B \subset \mathbb{R}^d$.

Monge's problem remained open until the 1940s, when it was revisited by **Leonid Vitaliyevitch Kantorovich** (1912-1986; Nobel Prize in Economics in 1975) for the economic problem of optimal allocation of resources; see e.g. **Villani (2008)**, **Santambrogio (2015)**, **Galichon (2016)**.

In the **Kantorovich primal problem**, the objective is to find the **optimal transportation plan** γ , which solves


$$\inf_{\gamma \in \Gamma(\alpha, \beta)} \int_{\mathbb{R}^d \times \mathbb{R}^d} c(\mathbf{x}, \mathbf{y}) d\gamma(\mathbf{x}, \mathbf{y}), \quad (2)$$

where the infimum is over all coupling (\mathbf{X}, \mathbf{Y}) of (α, β) , belonging to $\Gamma(\alpha, \beta)$, the set of probability measures γ on $\mathbb{R}^d \times \mathbb{R}^d$, satisfying


Remark

*Solving Kantorovich problem with the quadratic cost yields **L_2 -optimal assignment**. The solution is an optimal transportation plan induced by the optimal map \mathcal{T} , which can be expressed as the gradient of a convex function.*

We make use of this theory to transport different types of measures, as depicted in [Peyré & Cuturi \(2019\)](#)




We make use of this theory to transport different types of measures, as depicted in [Peyré & Cuturi \(2019\)](#)



Some examples:

- PDEs: Jacobi equation, Monge-Ampère equation
- Differential geometry: geodesic, curvature, exponential mapping
- Machine learning: image processing, adversarial learning
- Statistics: Wasserstein distance based procedures

- Kids entertainment (actually, image processing in ML)




from Peyre and Cuturi (2019).

- Kids entertainment (actually, shape analysis and morphing in ML)



- Kids entertainment (actually, shape analysis and morphing in ML)





Q: How is this related to semiparametric inference for multivariate time series?

Q: How is this related to semiparametric inference for multivariate time series?

A: In short: measure transportation theory provides us with the tools needed to define ranks and signs in a multivariate setting. Those tools are pivotal for our novel theory.


Example (Ex 1: US Hstarts & Mortg)

We consider the seasonally adjusted series of **monthly** housing starts (Hstarts, left panel) and the 30-year conventional mortgage rate (Mortg, right panel—no need for seasonal adjustment) **from January 1989 to January 2016**, with a sample size $n = 325$.



Example (Ex 1: US Hstarts & Mortg)

We consider the seasonally adjusted series of **monthly** housing starts (Hstarts, left panel) and the 30-year conventional mortgage rate (Mortg, right panel—no need for seasonal adjustment) **from January 1989 to January 2016**, with a sample size $n = 325$.



Comment

- Visual inspection suggests the presence of significant auto- and cross-correlations, as expected from macroeconomic theory
- Hstarts and Mortg series seems to be driven by skew innovations (with large positive values more likely than the negative ones) \Rightarrow **No Gaussian or elliptical innovation density!!**

Issue (Empirical)

Despite the overwhelming empirical evidence of non-Gaussian and non-symmetric (non-elliptical) distributions, the statistical analysis of univariate and multivariate time series remains very deeply marked by explicit or implicit Gaussian assumptions:

- *Correlogram-based estimation methods*
- *Pseudo-Gaussian tests*
- *Gaussian quasi-likelihood and its possible robustification,*
- *Spectral methods (Whittle)*
- *...*

are ubiquitous in methodological developments as well as in daily practice.

Issue (Theoretical)

*In support of the Gaussian procedures, there is a **theoretical justification**: Gaussian QMLE is asymptotically valid (viz., pseudo-Gaussian tests have correct asymptotic nominal size and Gaussian quasi-maximum-likelihood estimators (QMLEs) are root- n consistent) under a broad range \mathcal{F} of non-Gaussian innovation densities f (typically, under finite fourth-order moments); see, e.g., **Gourieroux, Monfort & Trognon (Econometrica, 1984)** in the econometric literature...*

Issue (Theoretical)

*In support of the Gaussian procedures, there is a **theoretical justification**: Gaussian QMLE is asymptotically valid (viz., pseudo-Gaussian tests have correct asymptotic nominal size and Gaussian quasi-maximum-likelihood estimators (QMLEs) are root- n consistent) under a broad range \mathcal{F} of non-Gaussian innovation densities f (typically, under finite fourth-order moments); see, e.g., **Gourieroux, Monfort & Trognon (Econometrica, 1984)** in the econometric literature...*

....but there exist theoretical aspects which are often overlooked...

Issue (Theoretical (cont'd))

- Estimation (HLLa)

- (i) while achieving efficiency under Gaussian innovations, their asymptotic performance can be quite poor under non-Gaussian ones;
- (ii) due to technical reasons (the Fisher consistency requirement), the choice of a quasi-likelihood is always the most pessimistic one: quasi-likelihoods automatically are based on the least favourable innovation density;
- (iii) actual fourth-order moments may be infinite.

- Testing (HLLb)

- (iv) Asymptotic validity of Gaussian test holds *pointwise*, for any given $f \in \mathcal{F}$, but it fails to hold *uniformly* over \mathcal{F} .
- (v) Lack of power under alternative hypothesis

Rank-based procedures represent valid alternatives:

Rank-based procedures represent valid alternatives:

- In the **univariate setting**, rank-based procedures (estimation and testing) offer a solution to all these issues. R-estimation and testing allows for efficiency/power improvements in the context of location (Hodges and Lehmann 1956) and regression models with independent observations (Jurečková 1971, Koul 1971, van Eeden 1972, Jaeckel 1972). Later on, it was extended to autoregressive time series (Koul and Saleh 1993, Koul and Ossiander 1994, Terpstra et al. 2001, Hettmansperger and McKean 2008, Mukherjee and Bai 2002, Andrews 2008, 2012) and non-linear time series (Mukherjee 2007, Andreou and Werker 2015, Hallin and La Vecchia 2017, 2019).

Rank-based procedures represent valid alternatives:

- In the **univariate setting**, rank-based procedures (estimation and testing) offer a solution to all these issues. R-estimation and testing allows for efficiency/power improvements in the context of location (Hodges and Lehmann 1956) and regression models with independent observations (Jurečková 1971, Koul 1971, van Eeden 1972, Jaeckel 1972). Later on, it was extended to autoregressive time series (Koul and Saleh 1993, Koul and Ossander 1994, Terpstra et al. 2001, Hettmansperger and McKean 2008, Mukherjee and Bai 2002, Andrews 2008, 2012) and non-linear time series (Mukherjee 2007, Andreou and Werker 2015, Hallin and La Vecchia 2017, 2019).
- In the **multivariate setting**, notions of multivariate ranks and signs have been proposed in the statistical literature. Among them, the componentwise ranks (Puri and Sen 1971), the spatial ranks (Oja 2010), the depth-based ranks (Liu 1992; Liu and Singh 1993). Those ranks and signs all have their own merits but also some drawbacks, which make them unsuitable for our needs. A notable exception are the Mahalanobis ranks and signs in Hallin and Paindaveine (2002, 2004a), under the restrictive assumption of elliptical innovation density.

Rank-based procedures represent valid alternatives:

- In the **univariate setting**, rank-based procedures (estimation and testing) offer a solution to all these issues. R-estimation and testing allows for efficiency/power improvements **in the context of location** (Hodges and Lehmann 1956) and regression models with independent observations (Jurečková 1971, Koul 1971, van Eeden 1972, Jaeckel 1972). Later on, it was extended to autoregressive time series (Koul and Saleh 1993, Koul and Ossander 1994, Terpstra et al. 2001, Hettmansperger and McKean 2008, Mukherjee and Bai 2002, Andrews 2008, 2012) and non-linear time series (Mukherjee 2007, Andreou and Werker 2015, Hallin and La Vecchia 2017, 2019).

- **Comment**

The problem with other notions of ranks is that they are not distribution-free (spatial ranks) and/or they do not feature essentially maximal ancillarity (depth-based ranks): this implies that semiparametric efficiency cannot be obtained—do not exploit completely spatial info; see **Hallin, del Barrio, Cuesta Albertos, and Matran (AoS, 2020)** and **Hallin, Hlubinka and Hudecova (AoS, 2020)**.

General problem: R-estimation involves the ranking (an ordering) of some quantities. Unlike the real line, the space \mathbb{R}^d , $d \geq 2$ is not canonically ordered. As a consequence basic univariate concepts as

- (in population) quantile and distribution functions,
- (in samples) empirical quantiles (quantile contours), ranks, signs, etc.,

do not canonically/straightforwardly/satisfactorily extend to the multivariate context. *How to define rank-based procedures in the multivariate setting?*

General problem: R-estimation involves the ranking (an ordering) of some quantities. Unlike the real line, the space \mathbb{R}^d , $d \geq 2$ is not canonically ordered. As a consequence basic univariate concepts as

- (in population) quantile and distribution functions,
- (in samples) empirical quantiles (quantile contours), ranks, signs, etc.,

do not canonically/straightforwardly/satisfactorily extend to the multivariate context. **How to define rank-based procedures in the multivariate setting?**

Contribution

We tackle the general problem. We derive a novel class of semiparametric inference procedures, a family of estimators and tests (select the rank-based estimating function/test statistic 'you like') working for multivariate time series: the procedures are based on center-outward ranks and signs (related to measure transportation).

General problem: R-estimation involves the ranking (an ordering) of some quantities. Unlike the real line, the space \mathbb{R}^d , $d \geq 2$ is not canonically ordered. As a consequence basic univariate concepts as

- (in population) quantile and distribution functions,
- (in samples) empirical quantiles (quantile contours), ranks, signs, etc.,

do not canonically/straightforwardly/satisfactorily extend to the multivariate context. **How to define rank-based procedures in the multivariate setting?**

Contribution

We tackle the general problem. We derive a novel class of semiparametric inference procedures, a family of estimators and tests (select the rank-based estimating function/test statistic 'you like') working for multivariate time series: the procedures are based on center-outward ranks and signs (related to measure transportation).

Q: *Are our R-estimators and tests really better than Gaussian procedures?*

Example (Ex. 3 : semiparametric 2D time series (HLLa))

Let's consider the bivariate VAR(1) model

$$(I_2 - \mathbf{A}L) \mathbf{X}_t = \epsilon_t \quad t \in \mathbb{Z}, \quad \text{where } \mathbf{A} = \begin{bmatrix} 0.2 & 0.3 \\ 0.6 & 1.1 \end{bmatrix}$$

with $\mathbf{X}_t \in \mathbb{R}^2$ and innovation densities:

Example (Ex. 3 : semiparametric 2D time series (HLLa))

Let's consider the bivariate VAR(1) model

$$(I_2 - \mathbf{A}L) \mathbf{X}_t = \epsilon_t \quad t \in \mathbb{Z}, \quad \text{where } \mathbf{A} = \begin{bmatrix} 0.2 & 0.3 \\ 0.6 & 1.1 \end{bmatrix}$$

with $\mathbf{X}_t \in \mathbb{R}^2$ and innovation densities:

- a bivariate $\mathcal{N}(\mathbf{0}, \mathbf{I}_2)$ density

Example (Ex. 3 : semiparametric 2D time series (HLLa))

Let's consider the bivariate VAR(1) model

$$(I_2 - \mathbf{A}L) \mathbf{X}_t = \epsilon_t \quad t \in \mathbb{Z}, \quad \text{where } \mathbf{A} = \begin{bmatrix} 0.2 & 0.3 \\ 0.6 & 1.1 \end{bmatrix}$$

with $\mathbf{X}_t \in \mathbb{R}^2$ and innovation densities:

- a bivariate $\mathcal{N}(\mathbf{0}, \mathbf{I}_2)$ density
- a mixture of Gaussians

Example (Ex. 3 : semiparametric 2D time series (HLLa))

Let's consider the bivariate VAR(1) model

$$(I_2 - \mathbf{A}L) \mathbf{X}_t = \epsilon_t \quad t \in \mathbb{Z}, \quad \text{where } \mathbf{A} = \begin{bmatrix} 0.2 & 0.3 \\ 0.6 & 1.1 \end{bmatrix}$$


with $\mathbf{X}_t \in \mathbb{R}^2$ and innovation densities:

- a bivariate $\mathcal{N}(\mathbf{0}, \mathbf{I}_2)$ density
- a mixture of Gaussians
- a contaminated model, with additive outliers (AO)

Those densities were used to generate independent innovations ϵ_t in $N = 300$ replications of length $n = 1000$ of the stationary solution of this VAR(1).


Example (cont'd, Gaussian innovations)

For each replication, \mathbf{A} was estimated via Gaussian QMLE and **R-estimators** (defined using MK results and selecting a reference density)




Example (cont'd, Gaussian innovations)


For each replication, \mathbf{A} was estimated via Gaussian QMLE and **R-estimators** (defined using MK results and selecting a reference density)




Example (MSE ratio: Mixture innovations (top) & Gaussian (bottom))



Example (cont'd, Outliers)



Example (cont'd, Outliers)







Example (cont'd, graphical intuition)

In these examples the innovation density is not spherical and/or elliptical.

Example (cont'd, graphical intuition)

In these examples the innovation density is not spherical and/or elliptical. Empirical center-outward quantile contours computed from $n = 1000$ points drawn from (clockwise) the Gaussian mixture, the skew-normal and skew- t_3 and from a standard multivariate normal.



Example (White noise vs VAR(1) in a 3D model (HLLb))

Rejection frequencies (out of $N = 1000$ replications), under values $\ell \mathbf{B}$, $\ell = 0, 1, 1.5$ of the VAR(1) autoregression matrix, for a trivariate model ($d = 3$), and various innovation densities of the Gaussian and rank- and sign-tests of white noise against VAR(1); the sample size is $n = 1400$; the nominal level is $\alpha = 5\%$.

f Test	0	B	$1.5B$	f Test	0	B	$1.5B$
<u>Normal</u>				<u>Mixture</u>			
Gaussian	0.044	0.354	0.751	Gaussian	0.053	0.490	0.885
BC vdW	0.047	0.325	0.729	BCvdW	0.056	0.736	0.987
BC Spearman	0.048	0.285	0.696	BC Spearman	0.061	0.735	0.987
Sign	0.043	0.238	0.568	Sign	0.043	0.492	0.898
<u>t_3</u>				<u>Skew-t_3</u>			
Gaussian	0.035	0.376	0.763	Gaussian	0.038	0.412	0.819
BC vdW	0.060	0.464	0.888	BC vdW	0.056	0.759	0.992
BC Spearman	0.046	0.411	0.834	BC Spearman	0.058	0.728	0.990
Sign	0.044	0.422	0.832	Sign	0.052	0.501	0.924
<u>AOs ($\mathbf{s} = (6, 6, 6)'$)</u>				<u>AOs ($\mathbf{s} = (9, 9, 9)'$)</u>			
Gaussian	0.337	0.378	0.525	Gaussian	0.745	0.768	0.854
BC vdW	0.059	0.215	0.497	BC vdW	0.059	0.223	0.499
BC Spearman	0.053	0.186	0.448	BC Spearman	0.061	0.190	0.432
Sign	0.052	0.162	0.389	Sign	0.063	0.165	0.397

Q: How to proceed?

Q: How to proceed?

Denote by

$$\mathcal{E}^{(n)} = \left\{ P_{f, \theta}^{(n)} \mid f \in \mathcal{F}, \theta \in \Theta \right\}$$

a semiparametric model with Euclidean parameter of interest θ and nuisance f (the density of the iid noise driving the data-generating process, whose CDF is F).

Q: How to proceed?

Denote by

$$\mathcal{E}^{(n)} = \left\{ P_{f, \boldsymbol{\theta}}^{(n)} \mid f \in \mathcal{F}, \boldsymbol{\theta} \in \boldsymbol{\Theta} \right\}$$

a semiparametric model with Euclidean parameter of interest $\boldsymbol{\theta}$ and nuisance f (the density of the iid noise driving the data-generating process, whose CDF is F). For $\mathbf{X}_t^{(n)} \in \mathbb{R}^d$, $d \geq 1$, let $\mathbf{X}^{(n)} := \{\mathbf{X}_1^{(n)}, \dots, \mathbf{X}_n^{(n)}\}$ be a finite realization of a *VARMA process* with distribution $P_{f, \boldsymbol{\theta}}^{(n)}$, so the residual function

$$(\mathbf{X}^{(n)}, \boldsymbol{\theta}) \mapsto \mathbf{Z}^{(n)}(\boldsymbol{\theta})$$

is defined, where $\mathbf{Z}^{(n)}(\boldsymbol{\theta}) := \{\mathbf{Z}_1^{(n)}(\boldsymbol{\theta}), \dots, \mathbf{Z}_n^{(n)}(\boldsymbol{\theta})\}$.

Univariate case, $d = 1$. For a fix θ , the residuals are $Z_1^{(n)}(\theta), \dots, Z_n^{(n)}(\theta)$ and the univariate ranks are a mapping of $Z_1^{(n)}, \dots, Z_n^{(n)}$ to a regular grid on $(0, 1)$

$$\frac{1}{n+1}, \frac{2}{n+1}, \dots, \frac{n}{n+1}.$$

This mapping is related to the definition of the empirical version $F^{(n)}$ of the residuals distribution function F , for $t = 1, \dots, n$,

$$R_t^{(n)} := (n+1)F^{(n)}(Z_t^{(n)}).$$

The corresponding (in population) distribution function is

$$F : z \mapsto F(z) := \Pr[Z \leq z]$$

an increasing mapping, with values in the unit interval $(0, 1)$.

Univariate case, $d = 1$. For a fix θ , the residuals are $Z_1^{(n)}(\theta), \dots, Z_n^{(n)}(\theta)$ and the univariate ranks are a mapping of $Z_1^{(n)}, \dots, Z_n^{(n)}$ to a regular grid on $(0, 1)$

$$\frac{1}{n+1}, \frac{2}{n+1}, \dots, \frac{n}{n+1}.$$

This mapping is related to the definition of the empirical version $F^{(n)}$ of the residuals distribution function F , for $t = 1, \dots, n$,

$$R_t^{(n)} := (n+1)F^{(n)}(Z_t^{(n)}).$$

The corresponding (in population) distribution function is

$$F : z \mapsto F(z) := \Pr[Z \leq z]$$

an increasing mapping, with values in the unit interval $(0, 1)$.

Remark

F strongly exploits the left-to-right nature of the order relation " \leq ", a feature that does not make any sense in higher dimension (no left, no right, no up, no down!).

Let's consider the *center-outward* distribution function

$$\mathbf{F}_{\pm} := 2F - 1,$$

which is strictly increasing, with values in $(-1, 1)$, the unit ball \mathbb{S}_1 in \mathbb{R} .

We consider an empirical version $\mathbf{F}_{\pm}^{(n)}$ of \mathbf{F}_{\pm} defined as the monotone increasing mapping of $Z_1^{(n)}, \dots, Z_n^{(n)}$ to the regular grid (let $\lfloor n/2 \rfloor =: n_R$)

$$\left\{ 0, \pm \frac{1}{n_R + 1}, \dots, \pm \frac{n_R}{n_R + 1} \right\} \quad \text{when } n \text{ is odd } (n = 2n_R + 1),$$


$$\left\{ \pm \frac{1}{n_R + 1}, \dots, \pm \frac{n_R}{n_R + 1} \right\} \quad \text{when } n \text{ is even } (n = 2n_R)$$

of the interval $(-1, 1)$ (the unit ball in \mathbb{R}). So, for $i = 1, \dots, n$

$$\mathbf{S}_i^{(n)} R_{\pm; i}^{(n)} := (n_R + 1) \mathbf{F}_{\pm}^{(n)}(Z_i^{(n)}).$$

Clearly, F and \mathbf{F}_{\pm} convey the same information and the “center” is

$$\text{Med}(P_{f, \theta}) = F^{-1}(1/2) = \mathbf{F}_{\pm}^{-1}(0).$$

Mapping induced by $F_{\pm}^{(n)}$:

Remark

*One can prove that $\mathbf{F}_{\pm}^{(n)}$ is the solution to an optimal transportation problem. In measure transportation words, $\mathbf{F}_{\pm}^{(n)}$ achieves the **optimal assignment with quadratic cost function** from the $Z_t^{(n)}$'s to the n points of the grid over $(-1, 1)$.*

Hallin (2019): \mathbf{F}_{\pm} is the unique gradient of a convex function (Kantorovich potential) pushing F forward to $\mathcal{U}_{(-1,1)}$.


Remark

*One can prove that $\mathbf{F}_{\pm}^{(n)}$ is the solution to an optimal transportation problem. In measure transportation words, $\mathbf{F}_{\pm}^{(n)}$ achieves the **optimal assignment with quadratic cost function** from the $Z_t^{(n)}$'s to the n points of the grid over $(-1, 1)$. **Hallin (2019)**: \mathbf{F}_{\pm} is the unique gradient of a convex function (Kantorovich potential) pushing F forward to $\mathcal{U}_{(-1,1)}$.*

\Rightarrow This characterization can be adopted as the definition of \mathbf{F}_{\pm} . The huge advantage of this measure transportation-based definition is that **it does not involve the canonical ordering of \mathbb{R} , and therefore readily extends to \mathbb{R}^d , $d \geq 2$.**

Multivariate case, $d \geq 2$. Let \mathbb{S}_d denote the open unit ball in \mathbb{R}^d . The *center-outward distribution function* \mathbf{F}_\pm is the unique gradient of a convex function mapping \mathbb{R}^d to \mathbb{S}_d and **pushing F forward to the spherical uniform distribution U_d over \mathbb{S}_d** . This is an optimal mapping in the sense of **quadratic cost function**.

Multivariate case, $d \geq 2$. Let \mathbb{S}_d denote the open unit ball in \mathbb{R}^d . The *center-outward distribution function* \mathbf{F}_\pm is the unique gradient of a convex function mapping \mathbb{R}^d to \mathbb{S}_d and **pushing F forward to the spherical uniform distribution \mathbf{U}_d over \mathbb{S}_d** . This is an optimal mapping in the sense of **quadratic cost function**.



Based on this empirical center-outward distribution function, the *center-outward ranks* are

$$R_{\pm,t}^{(n)} := (n_R + 1) \|F_{\pm}^{(n)}(Z_t^{(n)})\|, \quad (3)$$

and the *center-outward signs* are

$$S_{\pm,t}^{(n)} := S_{\pm,t}^{(n)}(\theta) := F_{\pm}^{(n)}(Z_t^{(n)}) / [F_{\pm}^{(n)}(Z_t^{(n)}) \neq \mathbf{0}] / \|F_{\pm}^{(n)}(Z_t^{(n)})\|. \quad (4)$$

In the above definitions, we let n factorize into $n = n_R n_S + n_0$, for $n_R, n_S, n_0 \in \mathbb{N}$ and $0 \leq n_0 < \min\{n_R, n_S\}$, where $n_R \rightarrow \infty$ and $n_S \rightarrow \infty$ as $n \rightarrow \infty$.

Based on this empirical center-outward distribution function, the *center-outward ranks* are

$$R_{\pm,t}^{(n)} := (n_R + 1) \| \mathbf{F}_{\pm}^{(n)}(\mathbf{Z}_t^{(n)}) \|, \quad (3)$$

and the *center-outward signs* are

$$\mathbf{S}_{\pm,t}^{(n)} := \mathbf{S}_{\pm,t}^{(n)}(\theta) := \mathbf{F}_{\pm}^{(n)}(\mathbf{Z}_t^{(n)}) / [\mathbf{F}_{\pm}^{(n)}(\mathbf{Z}_t^{(n)}) \neq \mathbf{0}] / \| \mathbf{F}_{\pm}^{(n)}(\mathbf{Z}_t^{(n)}) \|. \quad (4)$$

In the above definitions, we let n factorize into $n = n_R n_S + n_0$, for $n_R, n_S, n_0 \in \mathbb{N}$ and $0 \leq n_0 < \min\{n_R, n_S\}$, where $n_R \rightarrow \infty$ and $n_S \rightarrow \infty$ as $n \rightarrow \infty$.

Remark

The center-outward distribution function \mathbf{F}_{\pm} is a homeomorphism between $\mathbb{R}^d \setminus \mathbf{F}_{\pm}^{-1}(\{\mathbf{0}\})$ and $\mathbb{S}_d \setminus \{\mathbf{0}\}$; see [Figalli 2018](#). This ensures the existence of closed and nested quantile contours, obtained as the images under \mathbf{F}_{\pm}^{-1} of nested hyperspheres.

The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

Sketch of the idea

- *Given some observations from a multivariate time series, we set up a VARMA model (or VAR for testing)*

The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

Sketch of the idea

- Given some observations from a multivariate time series, we set up a VARMA model (or VAR for testing)
- We prove *the LAN and derive the central sequence* (Prop 2.1 in HLLa)

The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

Sketch of the idea

- Given some observations from a multivariate time series, we set up a VARMA model (or VAR for testing)
- We prove *the LAN and derive the central sequence* (Prop 2.1 in HLLa)
- By means of $\mathbf{F}_{\pm}^{(n)}$, we obtain $R_{\pm,t}^{(n)}$ and $\mathbf{S}_{\pm,t}^{(n)}$ of the innovations

The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

Sketch of the idea

- Given some observations from a multivariate time series, we set up a VARMA model (or VAR for testing)
- We prove *the LAN and derive the central sequence* (Prop 2.1 in HLLa)
- By means of $\mathbf{F}_{\pm}^{(n)}$, we obtain $R_{\pm,t}^{(n)}$ and $\mathbf{S}_{\pm,t}^{(n)}$ of the innovations
- We project the *LAN central sequence in the space of the center-outward ranks and signs* (see Corollary 4.1 in HLLb and section 3.4 in HLLa) \Rightarrow obtain a new central sequence, depending on the center-outward quantities and on a reference density. Then,

The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

Sketch of the idea

- Given some observations from a multivariate time series, we set up a VARMA model (or VAR for testing)
- We prove *the LAN and derive the central sequence* (Prop 2.1 in HLLa)
- By means of $\mathbf{F}_{\pm}^{(n)}$, we obtain $R_{\pm,t}^{(n)}$ and $\mathbf{S}_{\pm,t}^{(n)}$ of the innovations
- We project the *LAN central sequence in the space of the center-outward ranks and signs* (see Corollary 4.1 in HLLb and section 3.4 in HLLa) \Rightarrow obtain a new central sequence, depending on the center-outward quantities and on a reference density. Then,
 - ▶ R-estimation (HLLa): we use the new central sequence to define an R-estimator (one-step estimation) and derive its **asymptotics** (root-n consistency and asymptotic normality). For the sake of simplicity we use an elliptic reference density


The combination of Le Cam's theory with these measure transportation results yields our novel center-outward ranks- and signs-based procedures.

Sketch of the idea

- Given some observations from a multivariate time series, we set up a VARMA model (or VAR for testing)
- We prove *the LAN and derive the central sequence* (Prop 2.1 in HLLa)
- By means of $\mathbf{F}_{\pm}^{(n)}$, we obtain $R_{\pm,t}^{(n)}$ and $\mathbf{S}_{\pm,t}^{(n)}$ of the innovations
- We project the *LAN central sequence in the space of the center-outward ranks and signs* (see Corollary 4.1 in HLLb and section 3.4 in HLLa) \Rightarrow obtain a new central sequence, depending on the center-outward quantities and on a reference density. Then,
 - ▶ *R-estimation (HLLa):* we use the new central sequence to define an R-estimator (one-step estimation) and derive its **asymptotics** (root-n consistency and asymptotic normality). For the sake of simplicity we use an elliptic reference density
 - ▶ *Testing (HLLb):* we use the new central sequence to define a class of tests and derive their **asymptotics** under the null (level) and under a sequence of local alternatives (power).


Example (Ex 2: US M1SL & CPIAUCSL)

M1SL series in levels (left panel, in Billions of Dollars) and CPIAUCSL series in levels (right panel). **Monthly** records.



Example (Ex 2: US M1SL & CPIAUCSL)

M1SL series in levels (left panel, in Billions of Dollars) and CPIAUCSL series in levels (right panel). **Monthly** records.



Comment

- The differentiated series seem to be correlated and display common movements (mainly in opposite directions)
- Starting from 2005, the trajectories of the differentiated series look increasingly asymmetric and spiky: March-September-2020 reveals a cluster of outlying values due to COVID-19-related policy decisions. \Rightarrow No Gaussian or elliptical innovation density!!

Example (Ex 2 (cont'd): VAR order selection for M1SL and CPIAUCSL)

Values of test statistics and p -values (in parenthesis) of the Gaussian, vdW, Spearman, sign tests, along with their p -values (in brackets) under the null hypotheses of a VAR(p_0) model ($p_0 = 0, \dots, 7$).

p_0	Gaussian	vdW	Spearman	Sign
0	25.05 (4.91×10^{-5})	224.67 (0)	238.71 (0)	356.90 (0)
1	25.83 (3.42×10^{-5})	16.77 (0.002)	15.42 (0.004)	26.16 (2.94×10^{-5})
2	2.30 (0.681)	62.35 (9.30×10^{-13})	71.57 (1.07×10^{-14})	16.56 (0.002)
3		48.59 (7.18×10^{-10})	56.90 (1.30×10^{-11})	47.99 (9.49×10^{-10})
4		10.74 (0.030)	13.14 (0.011)	26.58 (2.41×10^{-5})
5		19.02 (7.81×10^{-5})	21.37 (2.67×10^{-5})	16.57 (0.002)
6		7.58 (0.108)	24.14 (7.49×10^{-5})	32.58 (1.45×10^{-6})
7			4.27 (0.371)	8.60 (0.072)

Example (Ex 2 (cont'd): VAR order selection for M1SL and CPIAUCSL)

Values of test statistics and p -values (in parenthesis) of the Gaussian, vdW, Spearman, sign tests, along with their p -values (in brackets) under the null hypotheses of a VAR(p_0) model ($p_0 = 0, \dots, 7$).


p_0	Gaussian	vdW	Spearman	Sign
0	25.05 (4.91×10^{-5})	224.67 (0)	238.71 (0)	356.90 (0)
1	25.83 (3.42×10^{-5})	16.77 (0.002)	15.42 (0.004)	26.16 (2.94×10^{-5})
2	2.30 (0.681)	62.35 (9.30×10^{-13})	71.57 (1.07×10^{-14})	16.56 (0.002)
3		48.59 (7.18×10^{-10})	56.90 (1.30×10^{-11})	47.99 (9.49×10^{-10})
4		10.74 (0.030)	13.14 (0.011)	26.58 (2.41×10^{-5})
5		19.02 (7.81×10^{-5})	21.37 (2.67×10^{-5})	16.57 (0.002)
6		7.58 (0.108)	24.14 (7.49×10^{-5})	32.58 (1.45×10^{-6})

Comment

- sharp contrast between the conclusions of the Gaussian and rank-based methods; 2 lags vs 6-7 lags.
- this is in line with the simulation exercises: our educated guess is that the combination of skewness, kurtosis, and outliers (all due to the COVID-19 periods) are blurring the conclusions of Gaussian tests.


Example (Ex 2 (cont'd): VAR order selection for M1SL and CPIAUCSL)

To see this, we consider the scatter plot of the fitted residuals (VAR(7) model)



Example (Ex 2 (cont'd): VAR order selection for M1SL and CPIAUCSL)

To see this, we consider the scatter plot of the fitted residuals (VAR(7) model)



Take home message

Exploiting some results from measure transportation theory we defined:

Take home message

Exploiting some results from measure transportation theory we defined:

- *R-estimation on VARMA models (linear time series models)*
- *Rank and sign semiparametric tests for $\text{VAR}(p)$ vs $\text{VAR}(p+1)$.*

Those procedures are based on measure transportation results: optimal transport of the innovations to the unit ball, center-outward ranks and signs.

Take home message

Exploiting some results from measure transportation theory we defined:

- *R-estimation on VARMA models (linear time series models)*
- *Rank and sign semiparametric tests for $\text{VAR}(p)$ vs $\text{VAR}(p+1)$.*

Those procedures are based on measure transportation results: optimal transport of the innovations to the unit ball, center-outward ranks and signs.

OT results (Kantorovich dual problem) are also related to saddlepoint approximations; see the working paper:

On some connections between Esscher's tilting, saddlepoint approximations, and optimal transportation: a statistical perspective, by La Vecchia & Ronchetti;

Take home message

Exploiting some results from measure transportation theory we defined:

- *R-estimation on VARMA models (linear time series models)*
- *Rank and sign semiparametric tests for $\text{VAR}(p)$ vs $\text{VAR}(p+1)$.*

Those procedures are based on measure transportation results: optimal transport of the innovations to the unit ball, center-outward ranks and signs.

OT results (Kantorovich dual problem) are also related to saddlepoint approximations; see the working paper:

On some connections between Esscher's tilting, saddlepoint approximations, and optimal transportation: a statistical perspective, by La Vecchia & Ronchetti;
code available https://github.com/dvdlvc/MyGitHub/tree/Saddlepoint_MeasureTransportation.

Thank you

For questions: davide.lavecchia@unige.ch

Write for the log-likelihood ratio math detail

$$L_{\theta_0 + n^{-1/2}\tau^{(n)}/\theta_0; f}^{(n)} := \log dP_{\theta_0 + n^{-1/2}\tau^{(n)}/\theta_0; f}^{(n)} / dP_{\theta_0; f}^{(n)},$$

where $\tau^{(n)}$ is a bounded sequence of $\mathbb{R}^{p_1 d^2}$. Then,

Proposition

Under some regularity assumptions on the VAR process and on f , for any bounded sequence $\tau^{(n)}$ in $\mathbb{R}^{p_1 d^2}$, under $P_{\theta_0; f}^{(n)}$, as $n \rightarrow \infty$,

$$L_{\theta_0 + n^{-1/2}\tau^{(n)}/\theta_0; f}^{(n)} = \tau^{(n)'} \Delta_f^{(n)}(\theta_0) - \frac{1}{2} \tau^{(n)'} \Lambda_f(\theta_0) \tau^{(n)} + o_P(1), \quad (5)$$

and $\Delta_f^{(n)}(\theta_0) \rightarrow \mathcal{N}(\mathbf{0}, \Lambda_f(\theta_0))$.

Parametrically efficient (in the Hájek-Le Cam asymptotic sense) and **distribution-free** inference procedures in LAN families (with given f) are possible when the LAN central sequence $\Delta_f^{(n)}(\theta_0)$ can be expressed in terms of signs and ranks.

More precisely when there exists some $\underline{\Delta}_f^{(n)}(\theta_0)$ measurable with respects to the ranks and signs of the residuals $\mathbf{Z}_t^{(n)}(\theta_0)$ such that

$$\underline{\Delta}_f^{(n)}(\theta_0) - \Delta_f^{(n)}(\theta_0) = o_P(1)$$

under $P_{\theta_0;f}^{(n)}$ as $n \rightarrow \infty$.

In mathematical detail:

$$\Delta_f^{(n)}(\theta) := \mathbf{M}'_{\theta} \mathbf{P}'_{\theta} \mathbf{Q}_{\theta}^{(n)'} \Gamma_f^{(n)}(\theta), \quad (6)$$

where \mathbf{M}_{θ} , \mathbf{P}_{θ} , and $\mathbf{Q}_{\theta}^{(n)}$ (see the JASA paper for an explicit form) do not depend on f and

$$\Gamma_f^{(n)}(\theta) := ((n-1)^{1/2}(\text{vec}\Gamma_{1,f}^{(n)}(\theta))', \dots, (n-i)^{1/2}(\text{vec}\Gamma_{i,f}^{(n)}(\theta))', \dots, (\text{vec}\Gamma_{n-1,f}^{(n)}(\theta))')'$$

with the so-called *f-cross-covariance matrices*

$$\Gamma_{i,f}^{(n)}(\theta) := (n-i)^{-1} \sum_{t=i+1}^n \varphi_f(\mathbf{Z}_t^{(n)}(\theta)) \mathbf{Z}_{t-i}^{(n)'}(\theta).$$

In mathematical detail:

$$\Delta_f^{(n)}(\theta) := \mathbf{M}'_{\theta} \mathbf{P}'_{\theta} \mathbf{Q}_{\theta}^{(n)'} \Gamma_f^{(n)}(\theta), \quad (6)$$

where \mathbf{M}_{θ} , \mathbf{P}_{θ} , and $\mathbf{Q}_{\theta}^{(n)}$ (see the JASA paper for an explicit form) do not depend on f and

$$\Gamma_f^{(n)}(\theta) := ((n-1)^{1/2}(\text{vec} \Gamma_{1,f}^{(n)}(\theta))', \dots, (n-i)^{1/2}(\text{vec} \Gamma_{i,f}^{(n)}(\theta))', \dots, (\text{vec} \Gamma_{n-1,f}^{(n)}(\theta))')'$$

with the so-called *f-cross-covariance matrices*

$$\Gamma_{i,f}^{(n)}(\theta) := (n-i)^{-1} \sum_{t=i+1}^n \varphi_f(\mathbf{Z}_t^{(n)}(\theta)) \mathbf{Z}_{t-i}^{(n)'}(\theta).$$

A rank-based, hence distribution-free, *f*-cross-covariance matrices of the form ($i = 1, \dots, n-1$) is (for an adequate multivariate reference density f)

$$\tilde{\Gamma}_{i,f}^{(n)}(\theta) := (n-i)^{-1} \sum_{t=i+1}^n \varphi_f \left(\mathbf{F}_{\pm}^{-1} \left(\frac{R_{\pm,t}^{(n)}}{n_R + 1} \mathbf{S}_{\pm,t}^{(n)} \right) \right) \mathbf{F}_{\pm}^{-1'} \left(\frac{R_{\pm,t-i}^{(n)}}{n_R + 1} \mathbf{S}_{\pm,t-i}^{(n)} \right).$$

In the JASA paper, we concentrate on rank-based cross-covariance matrices of the form $(i = 1, \dots, n - 1)$ of type:

$$\tilde{\Gamma}_{i,J_1,J_2}^{(n)}(\theta) := (n - i)^{-1} \sum_{t=i+1}^n J_1 \left(\frac{R_{\pm,t}^{(n)}}{n_R + 1} \right) J_2 \left(\frac{R_{\pm,t-i}^{(n)}}{n_R + 1} \right) \mathbf{s}_{\pm,t}^{(n)} \mathbf{s}_{\pm,t-i}^{(n)'}.$$

to which $\tilde{\Gamma}_{i,f}^{(n)}(\theta)$ reduces, with

$$J_1(u) = \varphi_{\mathfrak{f}}(F_{d;\mathfrak{f}}^{\star-1}(u))$$

and

$$J_2(u) = F_{d;\mathfrak{f}}^{\star-1}(u),$$

in the case of a **spherical reference f** with radial density \mathfrak{f} , yielding a rank-based version $\tilde{\Delta}_f^{(n)}$ of $\Delta_f^{(n)}$.

Jump Back