

# Discussion

## ■ *iNtr*

- \* used vigorously for 2015 NRI;
- \* will ultimately replace the current in-house comparison program.

## ■ *viscover*

- \* can be generalized to visualize and integrate different data sources in R.
  - Web Services seen in other official statistics: Normalized Difference Vegetation Index (NDVI), Vegetation Condition Index (VCI), National Land Cover Data (NLCD), etc.

# References

1. Groves, Robert M., and Lars Lyberg. "Total survey error: Past, present, and future." *Public opinion quarterly* 74.5 (2010): 849-879.
2. Brackstone, Gordon. "Managing data quality in a statistical agency." (2003).
3. Karr, Alan F., Ashish P. Sanil, and David L. Banks. "Data quality: A statistical perspective." *Statistical Methodology* 3.2 (2006): 137-173.
4. Williams, Matthew, and Emily Berg. "Incorporating User Input Into Optimal Constraining Procedures for Survey Estimates". *Journal of Official Statistics* 29.3 (2013): 375-396.
5. Joblin, Mitchell, and Wolfgang Mauerer. "An Interactive Survey Application for Validating Social Network Analysis Techniques." *R Journal* 8.1 (2016).
6. U.S. Department of Agriculture. 2018. Summary Report: 2015 National Resources Inventory, Natural Resources Conservation Service, Washington, DC, and Center for Survey Statistics and Methodology, Iowa State University, Ames, Iowa.
7. X. Lyu, E. J. Berg, and H. Hofmann. Empirical bayes small area prediction of sheet and rill erosion under a zero-inflated lognormal model. 2018. Manuscript submitted for publication.