Math and Theory: collection overview

Dev D. Goyal

26 October 2026

Abstract

The catalogue contains seven result families and seven preprints in mathematics and theoretical computer science. Six preprints have complete Lean proofs of their principal claims. The Gyires preprint proves inequalities for structured matrices and develops related polynomial bounds. Its principal order-five claims, the all-order four-block theorem, and the exceptional-vector theorem have complete Lean proofs. The results comprise counterexamples, exact special cases, and sharp sequence bounds. The collection makes no claim of first priority. The papers have not been peer reviewed.

Contents

1 How to use the collection **1**

2 Mathematics **2**
2.1 Real positive-definite counterexamples to extended q-permanent monotonicity . 2
2.2 Gyires's permanent inequality for structured matrices 2
2.3 An entropy increase for a structurally disassortative DomiRank graph 2
2.4 Diagonal concavity does not imply concavity for sums of elementary symmetric polynomials 3
2.5 Sharp root log-concavity of alternating Motzkin sums 3
2.6 Counterexamples to a late-iterate ordering conjecture for generalized Hofstadter functions 3

3 Theoretical computer science **3**
3.1 Counterexamples to a best-of-n win-rate bound at every budget 3

4 Verification and authorship **3**

1 How to use the collection

The manuscript catalogue links each paper to its formal statements and independent computations. The verification guide records the commands and their results. The formal library uses one Lake project, pinned to Lean 4.35.0 and an exact Mathlib revision. The 29 selected formal statements include equality, sharpness, and auxiliary positivity claims within these seven papers.

2 Mathematics

2.1 Real positive-definite counterexamples to extended q-permanent monotonicity

For every real parameter $q > 1$, some real symmetric positive-definite matrix has a negative q-permanent derivative at q. An explicit matrix of order five decreases from $q = 14$ to $q = 15$. The construction also gives correlation matrices with no zero entries and integer-entry examples. The dimension can depend on q. The real-symmetric question on $[-1, 1]$ remains outside the result.

Scope: Counterexample and all-parameter extension.

Manuscript Formal statement and prior work.

2.2 Gyires's permanent inequality for structured matrices

Every nonnegative real circulant matrix A of order five satisfies $24\,\mathrm{per}(AA^T) \le 625\,\mathrm{per}(A)^2$. Equality holds exactly for constant matrices. This proves the order-five circulant case of Gyires's ratio inequality. A bound in sorted-gap coordinates has sharp coefficient 484. The conjecture for arbitrary doubly stochastic matrices remains open.

Scope: Structured matrix inequalities and related polynomial bounds. The principal order-five claims, the all-order four-block theorem, and the exceptional-vector theorem have Lean proofs.

Manuscript Formal statement and prior work.

Broader arguments in the same preprint

The manuscript also proves results for circulants of orders six and seven, with sharp sorted-gap margins. It treats $aI + bQ + cQ^2$ in every dimension, where Q is any permutation matrix and the coefficients are nonnegative. Further results cover four-block doubly stochastic matrices with unequal block sizes and boundary entries, matrices with one exceptional column or row, and every rank-at-most-two doubly stochastic matrix through order fifteen.

The supporting sections prove structural, scalar, deletion, balancing, local polynomial, and configuration-model results. Exact counterexamples identify the limits of auxiliary assertions. The complete support record gives the source map and exact reproduction commands.

The all-order four-block and exceptional-vector theorems have complete Lean proofs. Their inequalities and equality cases are among the 29 selected formal statements. The principal higher-order circulant, permutation-polynomial, and finite rank-two theorems remain incomplete in Lean. The all-order rank-two problem and the unrestricted Gyires conjecture remain unproved here.

2.3 An entropy increase for a structurally disassortative DomiRank graph

A connected graph with 14 vertices and 32 edges has negative degree–log-degree edge covariance. All resolvent scores are positive on $[251/1000, 63/250]$ inside the stable domain. Its entropy nevertheless increases by more than $1/10000$ in natural-logarithm units. This refutes the source's assertion for every positive-score interval.

Scope: Counterexample to an interval monotonicity conjecture.

Manuscript Formal statement and prior work.

2.4 Diagonal concavity does not imply concavity for sums of elementary symmetric polynomials

For $M > 0$, let $f_M = 1 + M\sigma_1 + \sigma_2 + M\sigma_3 + \sigma_4$ in four variables, where σ_j is the elementary symmetric polynomial of degree j. The fourth root $f_M^{1/4}$ is concave on the positive diagonal exactly when $M \geq 1$. It is concave on the full positive orthant exactly when $M = 1$. Thus every $M > 1$ disproves Lachaume's criterion with strictly positive coefficients.

Scope: Counterexample and complete parameter classification.

Manuscript Formal statement and prior work.

2.5 Sharp root log-concavity of alternating Motzkin sums

Let $T_n = \sum_{j=0}^{n}(-1)^{n-j}M_j$, where M_j is the Motzkin number. The sequence $T_n^{1/n}$ is strictly log-concave on its tail starting at five. The inequality reverses at center five, so no earlier tail is log-concave. This strengthens Zhao's proposed tail starting at nine. The source recurrence and finite sum are linked in Lean.

Scope: Conjecture resolution with an optimal threshold.

Manuscript Formal statement and prior work.

2.6 Counterexamples to a late-iterate ordering conjecture for generalized Hofstadter functions

For every $k \geq 4$, some input n satisfies $F_k^{\circ(2k+1)}(n) > F_{k+1}^{\circ(2k+2)}(n)$. Here the superscript denotes repeated composition. The proof constructs inverse iterates with exact gap $k-3$. The witness $F_4^{\circ 9}(132) = 8 > 7 = F_5^{\circ 10}(132)$ gives the first instance. The result concerns infinitely many parameters. It does not assert infinitely many violating inputs for each fixed parameter.

Scope: Counterexamples for every parameter k≥4.

Manuscript Formal statement and prior work.

3 Theoretical computer science

3.1 Counterexamples to a best-of-n win-rate bound at every budget

For every $n \geq 2$ and $\delta > 0$, a finite positive reference distribution with largest atom at most δ violates the proposed best-of-n divergence–win-rate curve. A four-response best-of-two example gives exact probabilities and a strict logarithmic certificate. The all-budget proof uses finite interval bounds and gives a quantitative positive gap. It does not determine the sharp replacement curve.

Scope: Disproof for every budget n≥2 and every positive atom cap.

Manuscript Formal statement and prior work.

4 Verification and authorship

Lean checks the proofs against their formal statements. Separate Challenge modules state the intended claims. Comparator checks these against their proved Solutions. The verification record identifies the runs of Lean, NanoDa, and con-ron. Independent exact and numerical programs test finite certificates and examples. These finite computations do not replace the general proofs.

Dev D. Goyal directed the research and is responsible for publication. AI tools contributed mathematical arguments, Lean formalizations, computations, and editing. The authorship disclosure identifies these tools and their roles. These preprints have not been peer reviewed.