Math and Theory: Collection Overview

Dev D. Goyal

26 October 2026

Abstract

The catalogue contains seven result families and seven preprints in mathematics and theoretical computer science. Six preprints have complete Lean proofs of their principal claims. The Gyires preprint combines verified order-five results with completed broader ordinary-proof arguments. Lean verification, novelty assessment, and independent human expert review of those additions remain incomplete. The results comprise counterexamples, exact special cases, and sharp sequence bounds. The collection makes no claim of first priority. Machine checking and AI review do not constitute independent human expert peer review.

Contents

1 How to use the collection **1**

2 Mathematics **2**
- 2.1 Real positive-definite counterexamples to extended q-permanent monotonicity . 2
- 2.2 Gyires's permanent inequality for structured matrices 2
- 2.3 An entropy increase for a structurally disassortative DomiRank graph 2
- 2.4 Diagonal concavity does not imply concavity for sums of elementary symmetric polynomials 2
- 2.5 Sharp root log-concavity of alternating Motzkin sums 3
- 2.6 Counterexamples to a late-iterate ordering conjecture for generalized Hofstadter functions 3

3 Theoretical computer science **3**
- 3.1 Counterexamples to a best-of-n win-rate bound at every budget 3

4 Verification and authorship **3**

1 How to use the collection

The manuscript catalogue links each paper to its formal statements and independent computations. The verification guide records the commands and their results. The project uses one Lake installation, pinned to Lean 4.35.0 and an exact Mathlib revision. The 21 selected submission statements include equality, sharpness, and auxiliary positivity claims. They are grouped into seven papers, not presented as 21 independent discoveries.

2 Mathematics

2.1 Real positive-definite counterexamples to extended q-permanent monotonicity

For every real parameter $q > 1$, some real symmetric positive-definite matrix has a negative q-permanent derivative at q. An explicit matrix of order five decreases from $q = 14$ to $q = 15$. The construction also gives correlation matrices with no zero entries and integer-entry examples. The dimension can depend on q. The real-symmetric question on $[-1, 1]$ remains outside the result.

Scope: Counterexample and all-parameter extension.

Manuscript Formal statement and prior work.

2.2 Gyires's permanent inequality for structured matrices

Every nonnegative real circulant matrix A of order five satisfies $24\,\mathrm{per}(AA^T) \le 625\,\mathrm{per}(A)^2$. Equality holds exactly for constant matrices. This proves the order-five circulant case of Gyires's ratio inequality. A bound in sorted-gap coordinates has sharp coefficient 484. The conjecture for arbitrary doubly stochastic matrices remains open.

Scope: Verified order-five claims and the completed broader matrix and polynomial arguments in one continuous preprint.

Manuscript Formal statement and prior work.

Broader arguments in the same preprint

The continuous manuscript also includes circulants of orders six and seven, with sharp sorted-gap margins; the all-order family $aI + bQ + cQ^2$ for any permutation matrix Q and nonnegative coefficients; four-block doubly stochastic matrices with unequal block sizes and boundary entries; matrices with one exceptional column or row; and every rank-at-most-two doubly stochastic matrix through order fifteen.

The supporting sections give completed structural, scalar, deletion, balancing, local polynomial, and configuration-model arguments. Exact counterexamples state the limits of failed auxiliary claims. The complete support record gives the source map and exact reproduction commands.

The broader claims have incomplete Lean verification. They are outside the 21 selected formal statements. The earlier order-five verification does not certify these additions. The all-order rank-two problem and the unrestricted Gyires conjecture remain unproved here.

2.3 An entropy increase for a structurally disassortative DomiRank graph

A connected graph with 14 vertices and 32 edges has negative degree–log-degree edge covariance. All resolvent scores are positive on $[251/1000, 63/250]$ inside the stable domain. Its entropy nevertheless increases by more than $1/10000$ in natural-logarithm units. This refutes the source's assertion for every positive-score interval.

Scope: Counterexample to an interval monotonicity conjecture.

Manuscript Formal statement and prior work.

2.4 Diagonal concavity does not imply concavity for sums of elementary symmetric polynomials

For $M > 0$, let $f_M = 1 + M\sigma_1 + \sigma_2 + M\sigma_3 + \sigma_4$ in four variables, where σ_j is the elementary symmetric polynomial of degree j. The fourth root $f_M^{1/4}$ is concave on the positive diagonal

exactly when $M \geq 1$. It is concave on the full positive orthant exactly when $M = 1$. Thus every $M > 1$ disproves Lachaume's criterion with strictly positive coefficients.

Scope: Counterexample and complete parameter classification.

Manuscript Formal statement and prior work.

2.5 Sharp root log-concavity of alternating Motzkin sums

Let $T_n = \sum_{j=0}^{n} (-1)^{n-j} M_j$, where M_j is the Motzkin number. The sequence $T_n^{1/n}$ is strictly log-concave on its tail starting at five. The inequality reverses at center five, so no earlier tail is log-concave. This strengthens Zhao's proposed tail starting at nine. The source recurrence and finite sum are linked in Lean.

Scope: Conjecture resolution with an optimal threshold.

Manuscript Formal statement and prior work.

2.6 Counterexamples to a late-iterate ordering conjecture for generalized Hofstadter functions

For every $k \geq 4$, some input n satisfies $F_k^{\circ(2k+1)}(n) > F_{k+1}^{\circ(2k+2)}(n)$. Here the superscript denotes repeated composition. The proof constructs inverse iterates with exact gap $k-3$. The witness $F_4^{\circ 9}(132) = 8 > 7 = F_5^{\circ 10}(132)$ gives the first instance. The result concerns infinitely many parameters. It does not assert infinitely many violating inputs for each fixed parameter.

Scope: Counterexamples for every parameter k≥4.

Manuscript Formal statement and prior work.

3 Theoretical computer science

3.1 Counterexamples to a best-of-n win-rate bound at every budget

For every $n \geq 2$ and $\delta > 0$, a finite positive reference distribution with largest atom at most δ violates the proposed best-of-n divergence–win-rate curve. A four-response best-of-two example gives exact probabilities and a strict logarithmic certificate. The all-budget proof uses finite interval bounds and gives a quantitative positive gap. It does not determine the sharp replacement curve.

Scope: Disproof for every budget n≥2 and every positive atom cap.

Manuscript Formal statement and prior work.

4 Verification and authorship

Lean checks the proofs against their formal statements. Separate Challenge modules state the intended claims. Comparator checks these against their proved Solutions. The verification record identifies the runs of Lean, NanoDa, and con-ron. Independent exact and numerical programs test finite certificates and examples. These finite computations do not replace the general proofs.

Dev D. Goyal directed the research and is responsible for publication. The original research agent was ChatGPT (OpenAI), using the GPT-6 model with Pro reasoning. The earlier Gyires development also records work by Claude Opus 5.5, used with Claude Code. Codex, using GPT-6, audited and revised the collection. The authorship record states the precise review scope and identifies the final independent AI reviewer.

These papers have not been peer reviewed. The recorded personal review of the earlier Gyires proofs does not extend to this collection's new material. No mathematical-physics

result is included in this edition. New families can use the same structure after they pass the publication checks.