Depth-one Artin–Schreier defect towers are composita

Dev D. Goyal

30 October 2026

Abstract

Let k be an algebraically closed field of characteristic $p > 0$ and let K be a perfect henselian subfield of the Hahn field $k((y^{\mathbb{Q}}))$ that contains $k(y)$ and whose elements have no finite accumulation points in their supports. Nart and Novacoski proved that every finite compositum of Artin–Schreier extensions of K has depth one, and conjectured the converse. We prove the converse: a finite Artin–Schreier tower M/K of depth one is a compositum of Artin–Schreier extensions of K. More precisely, for every $\theta \in M \setminus K$ the truncation of θ at the first finite accumulation point of its support generates a nontrivial elementary abelian subextension of $K(\theta)/K$, and depth one forces this subextension to be all of M. The extraction step adapts an Ore-relation argument of Kedlaya. Both implications are formalized in Lean 4 in the setting of Nart and Novacoski, the algebraic closure of K inside the Hahn field with the Hahn valuation, with depth one expressed as minimum Okutsu length one; the identification of that minimum with depth is their Theorem 1.3.

Verification status. A fresh replay on 30 October 2026 compiled all 20 modules and a regenerated audit one at a time into a new directory, against a Mathlib checkout built from source at the pinned revision. All 435 declarations defined in these modules depend only on `propext`, `Classical.choice` and `Quot.sound`. Both implications are formalized; the identification of minimum Okutsu length with depth is cited, and Section 6 states the formal scope.

1 The theorem

Let k be an algebraically closed field of characteristic $p > 0$ and let $H = k((y^{\mathbb{Q}}))$ be the field of Hahn series with well-ordered rational support. Write v for the valuation by least exponent. Suppose

$$k(y) \subseteq K \subseteq H,$$

that K is perfect and henselian for v, and that every $a \in K$ has only finitely many support exponents below every real bound; for Hahn series this says that the support has no finite accumulation point. These are the standing assumptions of Nart and Novacoski [3, §3].

An *Artin–Schreier tower* is a chain $K = K_0 \subset K_1 \subset \cdots \subset K_r = M$ in which each step is generated by a root of $X^p - X - a_i$ with $a_i \in K_{i-1}$. Redundant steps can be removed, so we assume every step has degree p and $r \geq 1$. Such towers can be realized in H by the explicit root formula of Lemma 3.2, and by henselianity the valuation does not depend on the realization.

For an algebraic element θ write $\deg_K \theta = [K(\theta) : K]$. Nart and Novacoski define the depth of a primitive generator through Mac Lane–Vaquié chains and characterize it through Okutsu

sequences [3, Definition 1.2, Theorem 1.3]. For a nontrivial finite immediate extension, a primitive generator θ has depth one exactly when

$$\text{for every algebraic } b \text{ with } \deg_K b < \deg_K \theta \text{ there is } a \in K \text{ with } v(\theta - b) \le v(\theta - a). \tag{1}$$

The depth of M/K is the minimum depth over all primitive generators. As in [3, §3], distances are measured in the algebraic closure $\overline{K}$ of K inside H; the argument below only uses approximations lying in M itself.

Theorem 1. *Under these assumptions, let M/K be a nontrivial finite Artin–Schreier tower. If $\operatorname{depth}(M/K) = 1$, then M is a compositum of Artin–Schreier extensions of K.*

Nart and Novacoski proved the reverse implication [3, Theorem 3.15] and conjectured Theorem 1 [3, Conjecture 4.1]. Together they give:

Corollary 1. *Under the standing assumptions, a nontrivial finite Artin–Schreier tower M/K has depth one if and only if M is a compositum of Artin–Schreier extensions of K.*

The proof gives a stronger statement, Proposition 4.4: for every $\theta \in M \setminus K$, the first accumulation block of θ generates a nontrivial elementary abelian Galois subextension of $K(\theta)/K$.

2 Monomials and supports of the base

Put $\Gamma = vK \subseteq \mathbb{Q}$. Since K is perfect, Γ is p-divisible.

Lemma 2.1. *For every $q \in \Gamma$, the monomial y^q lies in K.*

Proof. Choose $a \in K$ with $v(a) = q$ and leading coefficient $\lambda \in k^\times$, and write $q = m/(p^r n)$ with $p \nmid n$. Perfection gives $y^{m/p^r} \in K$, so $u = a^n / y^{m/p^r} \in K$ is a unit with residue λ^n. Hensel's lemma applied to $X^n - u$ at the simple root λ of its reduction gives $b \in K$ with $b^n = u$ and residue λ. In H, a/y^q is another nth root of u with residue λ; their quotient is an nth root of unity, these all lie in k, and the only one with residue 1 is 1. Hence $b = a/y^q$ and $y^q = a/b \in K$. □

Lemma 2.2. *Every exponent in the support of an element of K lies in Γ, and every finite truncation of an element of K lies in K.*

Proof. If an exponent outside Γ occurs, let q be the least one. The earlier terms form a finite sum, since K has no finite accumulation points, and this sum lies in K by Lemma 2.1. Subtracting it leaves an element of K of valuation $q \notin \Gamma$, which is impossible. Finite truncations are then finite k-linear combinations of monomials in K. □

3 Accumulation points in an Artin–Schreier tower

Say that a Hahn series s has property C_Γ if the closure in $\mathbb{R}$ of its support is contained in Γ. The closure must be taken in $\mathbb{R}$: closure in $\mathbb{Q}$ would miss irrational accumulation points.

Lemma 3.1. *Property C_Γ is preserved by sums and products.*

Proof. The support of a sum lies in the union of the supports, and the closure of a finite union is the union of the closures. The support of a product st lies in $S+T$ with $S = \operatorname{supp} s$ and $T = \operatorname{supp} t$, both bounded below. If $s_n + t_n$ converges with $s_n \in S$, $t_n \in T$, both coordinate sequences are bounded, and a subsequence gives limits in $\operatorname{cl} S$ and $\operatorname{cl} T$. Hence $\operatorname{cl}(S+T) \subseteq \operatorname{cl} S + \operatorname{cl} T \subseteq \Gamma$. □

Lemma 3.2. *If a has property C_Γ, then every root in H of $X^p - X - a$ has property C_Γ.*

Proof. Write $a = a^- + a_0 + a^+$ by negative, zero and positive exponents, and choose $c \in k$ with $c^p - c = a_0$. One root is

$$\beta = \sum_{n\geq 1} (a^-)^{1/p^n} + c - \sum_{n\geq 0} (a^+)^{p^n}.$$

Both sums are Hahn-summable. The supports of the negative terms lie in $[v(a^-)/p^n, 0)$, so away from 0 only finitely many n contribute; if $a^+ \neq 0$ has least exponent $\varepsilon > 0$, only finitely many n with $p^n\varepsilon$ below a given bound contribute to the positive sum. Hence the union of the supports is well ordered and each exponent receives finitely many contributions, and applying Frobenius termwise gives $\beta^p - \beta = a$. The closure of the negative part lies in $\{0\} \cup \bigcup_{n\geq 1} p^{-n} \operatorname{cl}(\operatorname{supp} a^-)$, and the positive part is locally finite on bounded intervals with closure in $\bigcup_n p^n \operatorname{cl}(\operatorname{supp} a^+)$. Since Γ is p-divisible and closed under multiplication by p, all these points lie in Γ. The other roots differ from β by elements of $\mathbb{F}_p$. □

Proposition 3.3. *Every element of M has property C_Γ. In particular its support and all its finite accumulation points lie in Γ, and M/K is immediate.*

Proof. Elements of K have property C_Γ by Lemma 2.2 and the absence of finite accumulation points. If $K_i = K_{i-1}(\beta)$, Lemma 3.2 gives C_Γ for β, every element of K_i is a polynomial in β of degree less than p over K_{i-1}, and Lemma 3.1 completes the induction. Consequently $vM \subseteq \Gamma = vK$, and the residue field of M is k because $k \subseteq K \subseteq M \subseteq H$ and H has residue field k. □

The tower hypothesis is used here. In characteristic 2 the series $s = \sum_{n\geq 1} y^{(1-4^{-n})/3}$ satisfies $s^4 + ys + y = 0$ and has support in $\mathbb{Z}[1/2]$ but the accumulation point $1/3$, so support in Γ alone does not control accumulation points.

4 The first accumulation block

Fix $\theta \in M \setminus K$ and put $n = \deg_K \theta > 1$.

Lemma 4.1. *The support of θ has a finite accumulation point. If δ is the least one, then $\delta \in \Gamma$, the truncation*

$$\alpha = \operatorname{trn}_\delta(\theta) = \sum_{q<\delta} \theta_q\, y^q$$

has infinite support, and every proper lower truncation of α is finite.

Proof. If the support had no finite accumulation point, the finite truncations of θ would lie in K by Lemma 2.1 and Proposition 3.3 and would approximate θ arbitrarily well. As K is perfect, θ is separable; let κ be the largest value of $v(\theta - \theta')$ over the other conjugates θ'. A truncation $a \in K$ with $v(\theta - a) > \kappa$ would give $K(\theta) \subseteq K(a) = K$ by Krasner's lemma, a contradiction. A well-ordered subset of $\mathbb{R}$ cannot accumulate from the right, and its set of finite accumulation points

is closed, so a least one δ exists; below every $r < \delta$ the support is finite and below δ it is infinite. Proposition 3.3 gives $\delta \in \Gamma$. □

Lemma 4.2. *For every $a \in K$ we have $v(\theta - a) < \delta$, and these values are cofinal below δ.*

Proof. If $v(\theta - a) \geq \delta$, then a and θ agree below δ and $\operatorname{supp} a$ contains the infinite bounded set $\operatorname{supp} \alpha$, which is impossible for elements of K. For $r < \delta$, a finite truncation of θ at a bound between r and δ lies in K and approximates θ beyond r. □

Proposition 4.3. *The element $z = y^{-\delta}\alpha$ satisfies $P(z) = d$ for some $d \in K$ and some additive polynomial $P(X) = \sum_j \mu_j X^{p^j} \in k[X]$ with $\mu_0 \neq 0$.*

Proof. The powers $\theta, \theta^p, \theta^{p^2}, \ldots$ lie in the n-dimensional K-space $K(\theta)$, so there is a nontrivial relation $\sum_i a_i \theta^{p^i} = 0$ with $a_i \in K$; discard the zero coefficients. Let $q_i = v(a_i)$, let $\lambda_i \in k^\times$ be the leading coefficient of a_i, and put

$$c = \min_i (q_i + p^i \delta), \qquad I = \{i : q_i + p^i \delta = c\}.$$

Then $c \in \Gamma$. Write $\theta = \alpha + \eta$ with $v(\eta) \geq \delta$. Since Frobenius is additive, $a_i \theta^{p^i} = a_i \alpha^{p^i} + a_i \eta^{p^i}$ with $v(a_i \eta^{p^i}) \geq c$, so the tails do not contribute below c.

Expand each a_i into monomials $a_{i,q} y^q$. The term $a_{i,q} y^q \alpha^{p^i}$ has support in $(q + p^i v(\alpha),\ q + p^i \delta)$, so only finitely many q contribute below c, because a_i has finitely many support exponents below any bound. Since $q \geq q_i$ and $q_i + p^i \delta \geq c$, a contributing monomial has $q + p^i \delta \geq c$, with equality exactly when $i \in I$ and $q = q_i$. In that case the whole term $\lambda_i y^{q_i} \alpha^{p^i}$ lies below c. In every other case $q + p^i \delta > c$, and the part of $y^q \alpha^{p^i}$ below c comes from a proper lower truncation of α, which is finite. All these finite terms have exponents in Γ and so lie in K by Lemma 2.1. Truncating the relation below c therefore gives

$$\sum_{i \in I} \lambda_i y^{q_i} \alpha^{p^i} = R \in K.$$

As $q_i = c - p^i \delta$, dividing by $y^c \in K$ gives $Q(z) = y^{-c} R \in K$ with $Q(X) = \sum_{i \in I} \lambda_i X^{p^i} \neq 0$. If $m = \min I$, perfection of k gives P with nonzero linear coefficient and $Q = P^{p^m}$, perfection of K gives $d \in K$ with $d^{p^m} = y^{-c} R$, and injectivity of Frobenius on H gives $P(z) = d$. □

No algebraicity of α is assumed; the equation proves it. This extraction from an Ore relation follows the minimal-face argument of Kedlaya [2, Lemma 9.3]; the specialization used here is that all truncated terms other than the leading ones are finite.

Proposition 4.4. *The field $E = K(\alpha) = K(z)$ is a nontrivial elementary abelian Galois extension of K contained in $K(\theta)$, and distinct K-conjugates of α differ by elements of valuation δ.*

Proof. The roots of P form a finite $\mathbb{F}_p$-subspace C of k, because k is algebraically closed and P is separable and additive, and the roots of $P(X) - d$ are $z + C \subseteq K(z)$. So the minimal polynomial of z splits into distinct linear factors over $K(z)$ and E/K is Galois. The map $\sigma \mapsto \sigma(z) - z$ from $\operatorname{Gal}(E/K)$ to C is an injective homomorphism, since every σ fixes k, so the group is elementary abelian. The extension is nontrivial because α has infinite bounded support, which no element of K has. Two distinct conjugates of α differ by y^δ times a nonzero element of k.

For containment, let $b = \theta_\delta y^\delta \in K$, with $\theta_\delta = 0$ if $\delta \notin \operatorname{supp} \theta$. Then $v(\theta - b - \alpha) > \delta$, while all conjugates of α lie at distance exactly δ from α. Krasner's lemma gives $K(\alpha) \subseteq K(\theta - b) = K(\theta)$. □

5 Proof of Theorem 1

Let θ be a primitive generator of M/K of depth one, and let α, δ, E be as above. If $[E : K] < [M : K]$, then α is algebraic of degree less than $\deg_K \theta$ and

$$v(\theta - \alpha) \geq \delta > v(\theta - a) \qquad \text{for every } a \in K$$

by Lemma 4.2, contradicting (1). Hence $[E : K] = [M : K]$, and since $E \subseteq K(\theta) = M$ we get $E = M$. By Proposition 4.4, M/K is elementary abelian of exponent p, so it is the compositum of its cyclic subextensions of degree p. In characteristic p each of these is an Artin–Schreier extension: if σ generates its group, the trace of 1 is $p = 0$, additive Hilbert 90 gives γ with $\sigma(\gamma) - \gamma = 1$, and $\gamma^p - \gamma$ is fixed by σ. □

The argument uses only one primitive generator realizing depth one, not a particular generator of the tower. A second route, which only uses $\operatorname{supp} \theta \subseteq \Gamma$, obtains $\delta \in \mathbb{Q}$ from the extraction itself (equality of two of the affine functions $q_i + p^i\delta$), works over the tame extension $T = K(y^\delta)$, and descends to K through the decomposition $\mathrm{Gal}(NT/K) \cong \mathrm{Gal}(N/K) \times \mathrm{Gal}(T/K)$ for a Galois p-extension N containing M, using a normalized trace over $\mathrm{Gal}(T/K)$; we record it only as a check.

6 Formal verification

The Lean project pins Lean 4.19.0 and the Mathlib commit

`c44e0c8ee63ca166450922a373c7409c5d26b00b`.

Its 20 modules work with the actual Hahn field `HahnSeries` with rational exponents and coefficients in k. They construct Artin–Schreier roots by summable Hahn families, prove the support closure invariant of Proposition 3.3, the first accumulation cut, the Ore extraction of Proposition 4.3, the Galois and containment statements of Proposition 4.4 with Krasner's lemma in an arbitrary ambient field, and the lower-degree obstruction of Section 5.

The main endpoint is `okutsu_minimum_one_iff_asCompositum` in `HahnClosureForward.lean`. Its hypotheses are that k is algebraically closed of characteristic p; that K is an intermediate field of $k \subseteq H$ which is perfect, contains y and consists of left-finite series; that the restricted valuation satisfies `RationalValuedHenselian`, which says that it has a unique extension with values in $\mathbb{Q} \cup \{\infty\}$ to every algebraic extension; and that $M \neq K$ is the top of a literal Artin–Schreier tower inside H. It works in the algebraic closure `hahnClosureK` of K inside H, with the Hahn valuation, exactly as in [3, §3], and proves that 1 is the least length of an Okutsu sequence over all primitive generators of M if and only if the Artin–Schreier compositum of M is all of M. The two implications are `asCompositum_of_hahnClosure_okutsu_minimum` (the converse, Theorem 1) and `hahnClosure_okutsu_minimum_of_asCompositum` (the forward implication, [3, Theorem 3.15], for any nontrivial finite-dimensional M inside H generated by Artin–Schreier roots).

The forward proof follows [3]. Each Artin–Schreier generator is moved by an element of K to the explicit negative root of a finite negative element of K; this uses henselian splitting of elements of nonnegative valuation, proved with Krasner's lemma. A combination of these roots with coefficients in k that avoid finitely many linear relations is a primitive element whose nontrivial conjugates differ from it by nonzero constants. Krasner's lemma, after a constant shift in the boundary case, shows that no element of smaller degree comes within valuation zero of it, while partial inverse-Frobenius sums in K approximate it at every negative level. An earlier endpoint, `asCompositum_`

`of_algebraicClosure_okutsu_minimum`, proves the converse with Okutsu sequences in an abstract algebraic closure of K.

The following are not formalized. The identification of depth defined by Mac Lane–Vaquié chains with the minimum length of Okutsu sequences is [3, Theorem 1.3]. Henselianity enters only as the unique extension of the valuation to algebraic extensions, the standard definition of a henselian valued field [1, §4.1]; its equivalence with other forms of henselianity is not proved in Lean. The verifier `scripts/verify_cached.py` compiles every module into a fresh directory and checks that every audited declaration depends only on `propext`, `Classical.choice` and `Quot.sound`. Lean verification establishes only the formal statements described here.

7 Prior work and limits

Nart and Novacoski introduced depth for defect extensions, proved that composita of Artin–Schreier extensions have depth one and stated Theorem 1 as Conjecture 4.1 of [3], supporting it with a cyclic tower of depth two. The Artin–Schreier root formula, the support estimates, Krasner's lemma and the Ore relation are standard, and the minimal-face extraction is Kedlaya's [2, Lemma 9.3]. What is offered is the first-block statement, Proposition 4.4, over these bases and its application to all towers and all primitive generators. Searches in October 2026 found no other proof of the conjecture, but a search cannot establish that none exists, and the published journal version of [3] could not be opened. No priority is claimed, and no expert has reviewed the result.

Research process and AI disclosure

This is fully autonomous agentic research. Dev D. Goyal is responsible for publication. No expert validation or external peer review is claimed, and the novelty assessment is provisional.

The theorem, its proof, the Lean formalization, the literature searches and an earlier version of this note were produced by autonomous AI research agents in an earlier session. The records of that session do not name the AI system used.

Opus 5.5 by Anthropic used with Claude Code assessed the package for this release. It checked the argument step by step, read the arXiv source of [3] (version 2) and checked the depth criterion (1), Theorem 1.3, Theorem 3.15 and Conjecture 4.1 against the statements used here, and compared the Lean endpoints and their hypotheses with Theorem 1. It formalized the converse in the source's ambient field, the forward implication and their equivalence (the modules `HahnClosureDepth` and `HahnClosureForward`), ran the fresh Lean replay described in Section 6, and rewrote the note in this form. AI checks are not human peer review. Lean verification establishes only the formal statements described in Section 6.

References

[1] A. J. Engler and A. Prestel, *Valued Fields*, Springer, 2005.

[2] K. S. Kedlaya, On the algebraicity of generalized power series, *Beitr. Algebra Geom.* **58** (2017), 499–527. arXiv:1508.01836.

[3] E. Nart and J. Novacoski, Depth of Artin–Schreier defect towers, *J. Pure Appl. Algebra* **230** (2026), 108184. doi:10.1016/j.jpaa.2026.108184; arXiv:2503.18827v2 (16 October 2026).