# Proofs and corrections of OEIS conjectures

26 papers in 7 families

Dev D. Goyal · October 2026

This collection settles conjectures recorded in 26 entries of the On-Line Encyclopedia of Integer Sequences (OEIS). Most of the sequences are coefficients of formal power series defined by functional equations, and most of the conjectures concern their residues modulo small integers. In most entries the conjectured statements are proved. In four entries (A380680, A396798, A396843 and A396844) a conjectured statement is false; the papers give counterexamples and prove the correct statements. Each paper is self-contained, and every theorem in it is formalized in Lean 4 with Mathlib; the formal proofs use only Lean's standard axioms. All statements were also checked numerically against the published OEIS data.

The results were obtained in an autonomous AI research effort directed by Dev D. Goyal: ChatGPT (OpenAI, GPT-6 with Pro reasoning) found the proofs and wrote the Lean formalization from Dev D. Goyal's prompts, and Claude Opus 5.5, used with Claude Code, checked the work independently and wrote the papers in their present form. Dev D. Goyal has verified the results. The papers have not been peer reviewed.

## 1. The family $A_m(xA_m - mxA_m^2) = x^2$

Parity, residues modulo $m^2$ and binary-digit divisibility for the unique normalized solution of $A_m(xA_m - mxA_m^2) = x^2$, with corrected residue classifications for the two entries.

**A396843.** Odd coefficients, remainders modulo 9, and binary digits for the series $A(xA - 3xA^2) = x^2$

Every coefficient is odd; the exact residues modulo 9 (correcting the entry's second conjecture); $3^{s_2(n-1)}$ divides $a(n)$, where $s_2$ is the binary digit sum. *(proved; one statement corrected.)*

Paper · OEIS A396843 · Lean

**A396844.** Remainders modulo 16 for the series $A(xA - 4xA^2) = x^2$

The exact residues modulo 16; the conjecture fails first at $n = 17$. *(corrected.)*

Paper · OEIS A396844 · Lean

## 2. Consecutive iterates $F = x + F^{\circ r}\, F^{\circ(r+1)}$

For every integer $k$, the $k$-th iterate of $F_r$ modulo $2r$ is $x/(1 - kx)$ when $r$ is odd, with an explicit correction term when $r$ is even.

**A396099.** The iterates of a self-referential power series modulo 4

All four conjectured congruences modulo 4 for the series and its iterates. *(proved.)*

Paper · OEIS A396099 · Lean

**A396798.** The iterates of a self-referential power series modulo 8

Seven of the eight conjectured congruences modulo 8; the remaining one holds from $n = 2$, not from $n = 1$ as stated. *(proved; one statement corrected.)*

Paper · OEIS A396798 · Lean

**A396807.** The iterates of a self-referential power series modulo 10

For every integer $k$, the coefficients of the $k$-th iterate satisfy $[x^n]\, A^{\circ k} \equiv k^{n-1} \pmod{10}$. *(proved.)*

Paper · OEIS A396807 · Lean

## 3. Scaled inverses $A(x - x\,A(qx)/q) = x$

Every coefficient is 1 modulo $q+1$; for even $q$, the odd coefficients sit exactly at the powers of 2.

**A393856.** Congruences for the series $A$ with $A(x - x\,A(4x)/4) = x$
$a(n) \equiv 1 \pmod 5$, and $a(n)$ is odd exactly when $n$ is a power of 2. *(proved.)*
Paper · OEIS A393856 · Lean

**A393857.** Congruences for the series $A$ with $A(x - x\,A(5x)/5) = x$
$a(n) \equiv 1 \pmod 6$. *(proved.)*
Paper · OEIS A393857 · Lean

**A393858.** Congruences for the series $A$ with $A(x - x\,A(6x)/6) = x$
$a(n) \equiv 1 \pmod 7$, and $a(n)$ is odd exactly when $n$ is a power of 2. *(proved.)*
Paper · OEIS A393858 · Lean

**A393859.** Congruences for the series $A$ with $A(x - x\,A(7x)/7) = x$
$a(n) \equiv 1 \pmod 8$. *(proved.)*
Paper · OEIS A393859 · Lean

## 4. Theta series modulo 4

Series built from the Jacobi theta series $\theta_3$ whose coefficients are 2 modulo 4 exactly at squares (or twice squares) and divisible by 4 elsewhere.

**A378580.** Remainders modulo 4 for the series with $A(x/A(x)) = \theta_3(x)$
$A(x/A(x)) = \theta_3$: for $n > 0$, $a(n) \equiv 2 \pmod 4$ if $n$ is a square, and $4 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A378580 · Lean

**A378581.** Remainders modulo 4 for the series with $A(x\,A(x)) = \theta_3(x)$
$A(x\,A(x)) = \theta_3$: the same square pattern. *(proved.)*
Paper · OEIS A378581 · Lean

**A380060.** Square indices and remainders modulo 4 in a theta-type series
For $n > 0$, $a(n) \equiv 2 \pmod 4$ if $n$ is a square, and $4 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A380060 · Lean

**A380680.** Twice the squares, not the squares: remainders modulo 4 in a theta-type series
The square conjecture is false ($a(1) = 4$, $a(2) = 6$); in fact $a(n) \equiv 2 \pmod 4$ exactly when $n = 2k^2$. *(conjecture disproved; corrected statement proved.)*
Paper · OEIS A380680 · Lean

## 5. Catalan-type equations

Equations whose solutions reduce, modulo 4 or 8, to expressions in the Catalan series, whose coefficients are odd exactly at powers of 2.

**A393172.** The Catalan numbers inside the equation $A(x - x^2) = x + A(x)^2$
For $n > 1$, $a(n) \equiv 2 \pmod 4$ if $n = 2^k$, and $4 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A393172 · Lean

**A393171.** Remainders modulo 8 of a series built from the Catalan numbers
For $n > 2$, $a(n) \equiv 4 \pmod 8$ if $n = 2^k$ or $2^k + 1$, and $8 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A393171 · Lean

**A392210.** The odd-indexed terms of a nested square are even
$a(2n-1)$ is even for $n > 1$ (an elementary observation recorded in the entry). *(proved.)*
Paper · OEIS A392210 · Lean

**A389536.** Remainders modulo 4 for the equation $A(x^2 + 2x^3) = x\,A(x)$
For $n > 1$, $a(n) \equiv 2 \pmod 4$ if $n = 2^k + 1$, and $4 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A389536 · Lean

**A397588.** Odd coefficients at powers of two for $A = x + (xA^2)'$
$a(n)$ is odd exactly when $n$ is a power of 2, and $3 \mid a(n)$ for $n > 1$. *(proved.)*
Paper · OEIS A397588 · Lean

**A195194.** Parity of an infinitely nested series reversion
The infinitely nested reversion is well defined, and $a(n)$ is odd exactly when $n$ is a power of 2. *(proved.)*
Paper · OEIS A195194 · Lean

## 6. Reversions with an exact halving step

Equations whose coefficient recursion appears to divide by 2; the division is shown to be exact before reducing.

**A389537.** Congruences modulo 4 for the series with $A\big(A(x) - x\big) = A(x)^2$
For $n > 2$, $a(n) \equiv 2 \pmod 4$ if $n = 3 \cdot 2^k$, and $4 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A389537 · Lean

**A389542.** Congruences modulo 4 for the series with $A(A(x)^2 - x^2) = 4A(x)^3$
For $n > 1$, $a(n) \equiv 2 \pmod 4$ if $n = 2^k + 1$, and $4 \mid a(n)$ otherwise. *(proved.)*
Paper · OEIS A389542 · Lean

**A389474.** A power series whose square is $x^2 + 2x^3$ modulo 9
$A^2 \equiv x^2 + 2x^3 \pmod 9$, and $A^{3k} \equiv A(x^3)^k \pmod 3$. *(proved.)*
Paper · OEIS A389474 · Lean

**A389475.** Two congruences modulo 3 for the series with $A(3x^2) = 2x^2 + A^2 - 2A^3$
$A^3 \equiv A(x^3)$ and $A^2 \equiv x^2 + 2A(x^3) \pmod 3$. *(proved.)*
Paper · OEIS A389475 · Lean

## 7. Other results

Three further entries, each with its own method.

**A396093.** When are the coefficients of $B(B(B(x)))$ odd?
$a(n)$ is odd exactly when $n \equiv 1, 3, 7, 9 \pmod{10}$. *(proved.)*
Paper · OEIS A396093 · Lean

**A396102.** Every coefficient is 1 modulo 3: the series with $A(A(A(x))) = (1 + x)\,A(A(x))$
$a(n) \equiv 1 \pmod 3$, together with a theorem for the whole family $F^{\circ r} = (1 + x)F^{\circ(r-1)}$. *(proved.)*
Paper · OEIS A396102 · Lean

**A380056.** The coefficients of $(e^x - 1)/\cos 2x$ modulo 4 and 5
$a(n) \equiv 1 \pmod 4$, and $a(n) \equiv 2 \cdot 2^n + 3 \cdot 4^n \pmod 5$, so $5 \mid a(n)$ exactly when $4 \mid n$. *(proved.)*
Paper · OEIS A380056 · Lean