The Catalan numbers inside the equation $A(x - x^2) = x + A(x)^2$

A proof of a conjecture in OEIS A393172

Dev D. Goyal

October 2026

Abstract

There is exactly one power series $A(x) = x + 2x^2 + 8x^3 + 42x^4 + \cdots$ that satisfies the equation $A(x - x^2) = x + A(x)^2$. Its coefficients $a(n)$ form the sequence A393172 of the On-Line Encyclopedia of Integer Sequences (OEIS). Paul D. Hanna conjectured that, for $n > 1$, the coefficient $a(n)$ leaves remainder 2 on division by 4 when n is a power of 2, and is divisible by 4 otherwise. We prove this conjecture. The key observation is that, once every coefficient is replaced by its remainder modulo 4, the series $A(x)$ becomes $2C(x) - x$, where $C(x) = x + x^2 + 2x^3 + 5x^4 + \cdots$ is the generating function of the Catalan numbers. The conjecture then follows from the classical fact that a Catalan number is odd only at positions given by powers of 2. The result has also been checked by computer and proved formally in the Lean proof assistant.

How this paper was produced. This paper reports the result of an autonomous AI research effort. ChatGPT (OpenAI), using the GPT-6 model with Pro reasoning and working from prompts by Dev D. Goyal, found the proof, wrote its Lean 4 formalization, and wrote a first draft of the paper. Dev D. Goyal chose the problems and asked for checks and revisions, but did not supply proof ideas. Claude Opus 5.5, used with Claude Code, then checked the work independently: it rebuilt the formal proofs from source and confirmed that they rely on no unproved assumptions, recomputed the sequence from its definition and tested every claim against the published data, and read every step of the proofs. It also rewrote this paper for clarity. Dev D. Goyal has verified the results and is responsible for them. The paper has not been peer reviewed. The section "About this paper" at the end gives the details.

1 Introduction

1.1 The sequence

The *On-Line Encyclopedia of Integer Sequences* (OEIS) [4] is a large public database of integer sequences, used by mathematicians and scientists to identify sequences and to record what is known about them. Many entries also record *conjectures*: patterns that the contributor observed in the data but could not prove.

This paper is about entry A393172 [1], contributed by Paul D. Hanna in February 2026. The entry reads, in part:

```
G.f.  A(x) satisfies A(x-x^2) = x + A(x)^2.

Conjecture:  for n > 1, a(n) == 2 (mod 4) iff n is a power of 2, otherwise
a(n) is divisible by 4.
```

```
(3) A(x) = C(x) + A(C(x))^2, where C(x) = x + C(x)^2 is a g.f.  of the
Catalan numbers (A000108).
```

— OEIS A393172, Paul D. Hanna, February 2026

Here "G.f." stands for *generating function*: the sequence $a(1), a(2), a(3), \ldots$ is stored as the coefficients of the power series $A(x) = a(1)\,x + a(2)\,x^2 + a(3)\,x^3 + \cdots$. In the equation, $A(x - x^2)$ means that the polynomial $x - x^2$ is substituted for x in A, while $A(x)^2$ is the ordinary square $A(x) \cdot A(x)$. Section 2 explains these operations. The equation determines all the coefficients, one at a time (Theorem 3.1). The first terms, starting at $n = 1$, are 1, 2, 8, 42, 260, 1808, 13744, 112290, 974880, 8922276, ...

Formula (3) of the entry brings in the *Catalan numbers* $1, 1, 2, 5, 14, 42, 132, \ldots$ (OEIS A000108), one of the most common sequences in combinatorics. For example, they count binary trees: there are $1, 1, 2, 5, 14, \ldots$ trees with $1, 2, 3, 4, 5, \ldots$ leaves in which every branching point splits into a left and a right branch. We store them in the series

$$C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + 42x^6 + \cdots ,$$

and write c_n for its coefficient of x^n. This C is the only series without constant term that satisfies $C = x + C^2$. (In the language of trees: a tree is either a single leaf, counted by x, or a branching point with two smaller trees attached, counted by C^2.)

1.2 The question

Hanna's conjecture says that, for $n > 1$, the remainder of $a(n)$ on division by 4 is 2 when n is a power of 2, and 0 otherwise. The first row of the table shows these remainders. The indices printed in bold are the powers of 2. The second row anticipates the main result: it shows the remainders modulo 4 of the coefficients of $2C(x) - x$, which are 1 for $n = 1$ and $2c_n$ for $n > 1$.

n	**1**	**2**	3	**4**	5	6	7	**8**	9	10	11	12	13	14	15	**16**
$a(n)$ mod 4	**1**	**2**	0	**2**	0	0	0	**2**	0	0	0	0	0	0	0	**2**
$[x^n]\,(2C{-}x)$ mod 4	**1**	**2**	0	**2**	0	0	0	**2**	0	0	0	0	0	0	0	**2**

n	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**32**
$a(n)$ mod 4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**2**
$[x^n]\,(2C{-}x)$ mod 4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**2**

n	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
$a(n)$ mod 4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$[x^n]\,(2C{-}x)$ mod 4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

1.3 What this paper proves

Main result (Theorem 3.1). *Modulo* 4, *the series* $A(x)$ *equals* $2C(x) - x$. *In other words,* $a(n) - 2c_n$ *is divisible by* 4 *for every* $n > 1$.

Since $2c_n$ leaves remainder 2 on division by 4 when c_n is odd, and remainder 0 when c_n is even, the remainder of $a(n)$ depends only on whether c_n is odd. A classical fact (Lemma 2.3) says that c_n is odd exactly when n is a power of 2. Together these prove Hanna's conjecture (Corollary 3.2).

1.4 What was known

The OEIS entry records several equivalent forms of the defining equation, among them formula (3) quoted above, and a recurrence for $a(n)$ due to Seiichi Manyama. The parity of the Catalan numbers is classical; see for example [3]. We are not aware of an earlier proof of the conjecture. The companion entry A393171 [2], also by Hanna, contains a conjecture modulo 8 about the series $x + A(x)^2$; it is proved in a separate paper of this collection, using the result proved here.

1.5 Why it is interesting

Equations in which an unknown power series is substituted into something, or something is substituted into it, are called *functional equations*. They arise naturally when counting objects that are built recursively, such as trees, and the OEIS contains many sequences defined in this way. Their coefficients usually grow quickly and look irregular. Their remainders modulo small powers of 2, however, often show striking patterns, and explaining such a pattern is a way of understanding the sequence. In this case the explanation is that the Catalan numbers are hidden inside A: the substitution $x \mapsto x - x^2$ is exactly the one that the Catalan series undoes. The method of proof (guess a simple series that solves the equation modulo 4, and use the fact that the solution is unique) is a general one, and it is used throughout this collection of papers.

1.6 How to read this paper

Section 2 explains the few notions we need: power series, substitution, remainders, and the Catalan numbers modulo 2. Section 3 states the results precisely, and Section 4 proves them. Section 5 describes how the results were checked, by computer and by a formal proof in Lean. Section 6 discusses the scope of the result and some open questions.

2 Background

This section collects the definitions and facts used later. Readers who know formal power series and congruences can skip it and return when needed.

2.1 Formal power series

A *formal power series* is an expression

$$U(x) = u_0 + u_1 x + u_2 x^2 + u_3 x^3 + \cdots ,$$

determined by its sequence of coefficients $u_0, u_1, u_2, \ldots$. No number is ever substituted for x, so questions of convergence do not arise. We write $[x^n]\, U = u_n$ for the coefficient of x^n in U; two series are equal when all their coefficients are equal.

Series are added coefficient by coefficient and multiplied like polynomials:

$$[x^n]\,(UV) = u_0 v_n + u_1 v_{n-1} + \cdots + u_n v_0.$$

Each coefficient of a product is a finite sum. For example, $(1 - x)(1 + x + x^2 + x^3 + \cdots) = 1$, so $1 + x + x^2 + \cdots$ is the *multiplicative inverse* of $1 - x$, written $\frac{1}{1-x}$. More generally, every series D with constant term 1 has a unique multiplicative inverse $\frac{1}{D}$, whose coefficients are

determined one at a time by $D \cdot \frac{1}{D} = 1$ using only addition, subtraction and multiplication. Every fraction in this paper has a denominator of this kind, so it denotes a power series.

We write $U = O(x^d)$ when all coefficients of U below degree d are zero. So $U - V = O(x^d)$ means that U and V agree up to degree $d-1$. A series of the form $x + u_2x^2 + u_3x^3 + \cdots$, with constant term 0 and coefficient of x equal to 1, is called *normalized*.

2.2 Substituting one series into another

If $V(x)$ has constant term 0, we can substitute V for x in U:

$$U(V(x)) = u_0 + u_1V(x) + u_2V(x)^2 + u_3V(x)^3 + \cdots .$$

This is called *composition*, also written $U \circ V$. It makes sense because $V(x)^k$ has no terms below x^k, so only the finitely many terms with $k \le n$ affect the coefficient of x^n. For example, if $U = x + x^2$ and $V = x^2$, then $U(V(x)) = x^2 + x^4$. Composition is associative: $U \circ (V \circ W) = (U \circ V) \circ W$.

Substituting a series into itself repeatedly gives its *iterates*: $U^{\circ 2}(x) = U(U(x))$, $U^{\circ 3}(x) = U(U(U(x)))$, and so on, with $U^{\circ 0}(x) = x$ and $U^{\circ 1} = U$. They must not be confused with *powers* $U^2 = U \cdot U$, $U^3 = U \cdot U \cdot U$. (The OEIS sometimes writes $A^k(x)$ for the k-th iterate. In this paper the small circle always means an iterate, and a plain exponent always means a power.)

2.3 Undoing a substitution

A normalized series can always be "undone" by another normalized series.

Lemma 2.1 (Compositional inverse)**.** *For every normalized series U there is exactly one normalized series V with $U(V(x)) = x$, and this V also satisfies $V(U(x)) = x$. If U has integer coefficients, so does V.*

We call V the *compositional inverse* of U. The OEIS calls it the *series reversion* of U. For example, the compositional inverse of $x - x^2$ is $C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + \cdots$, whose coefficients are the Catalan numbers. Indeed $C - C^2 = x$, which says exactly that substituting C into $x - x^2$ gives x.

Proof. Write $V = x + v_2x^2 + v_3x^3 + \cdots$ with unknown coefficients. In $U(V) = V + u_2V^2 + u_3V^3 + \cdots$, the coefficient of x^n (for $n \ge 2$) equals v_n plus terms that involve only $v_2, \ldots, v_{n-1}$. Setting it equal to 0 therefore determines v_n from the earlier coefficients, using only additions, subtractions and multiplications. So there is exactly one V with $U \circ V = x$, and it has integer coefficients if U does. Applying the same construction to V gives a normalized W with $V \circ W = x$. Then $U = U \circ (V \circ W) = (U \circ V) \circ W = W$, so $V \circ U = V \circ W = x$ as well. □

2.4 Remainders and congruences

For an integer $m \ge 2$ we write $a \equiv b \pmod m$ ("a is congruent to b modulo m") when $a - b$ is divisible by m, that is, when a and b leave the same remainder after division by m. For example, $17 \equiv 5 \pmod 4$, since both leave remainder 1, and $-3 \equiv 1 \pmod 4$. The remainder of a on division by m is called its *residue* modulo m; it is one of $0, 1, \ldots, m-1$. Saying that a is odd is the same as saying $a \equiv 1 \pmod 2$.

Addition and multiplication are compatible with congruences, so the residues $0, 1, \ldots, m-1$, added and multiplied modulo m, form a ring, written $\mathbb{Z}/m\mathbb{Z}$. When $m = p$ is a prime we also write $\mathbb{F}_p$. In $\mathbb{F}_2 = \{0, 1\}$ we have $1 + 1 = 0$, so addition and subtraction coincide.

For power series with integer coefficients, $U \equiv V \pmod{m}$ means that *every* coefficient of $U - V$ is divisible by m. Equivalently, the two series become equal when every coefficient is replaced by its residue; we call this *reducing* the series modulo m. The key fact is the following.

Reduction modulo m is compatible with all the operations above. To reduce a sum, a product, a substitution, or an inverse, we may reduce the ingredients first and then compute in $\mathbb{Z}/m\mathbb{Z}$. This holds because every coefficient of the result is obtained from finitely many coefficients of the ingredients by adding, subtracting and multiplying, and these operations are compatible with remainders.

Rings. Several lemmas below hold over every *commutative ring*, that is, every set with an addition, subtraction and multiplication satisfying the usual rules of arithmetic (for example $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{Z}/m\mathbb{Z}$). No division is used; this matters because in $\mathbb{Z}/m\mathbb{Z}$ division is not always possible (for instance, 2 has no inverse in $\mathbb{Z}/4\mathbb{Z}$).

2.5 The method

Almost every proof in this paper follows the same three steps.

1. *Uniqueness.* Show that the defining equation determines the coefficients one at a time, using only addition, subtraction and multiplication. Then the equation has exactly one solution of the required shape, in every commutative ring, including $\mathbb{Z}/4\mathbb{Z}$.
2. *Reduction.* Reducing the integer solution modulo 4 gives a solution of the reduced equation, because reduction is compatible with all operations.
3. *A good guess.* Find a simple series that satisfies the reduced equation and check that it does. By uniqueness, this simple series *is* the reduction of the sequence, and its coefficients can simply be read off.

The creative part is step 3: finding a simple series that works. Step 1 sometimes needs care, when the equation seems to require a division. For a first example, take the Catalan series $C = x + C^2$ from above. Its coefficients $1, 1, 2, 5, 14, 42, \ldots$ are determined one at a time by this equation. Modulo 2, the series $S = x + x^2 + x^4 + x^8 + \cdots$ satisfies the same equation: in $\mathbb{F}_2$, $S^2 = x^2 + x^4 + x^8 + \cdots$ (see Lemma 2.2), so $x + S^2 = S$. Hence $C \equiv S \pmod{2}$, and the Catalan coefficient $[x^n]\, C$ is odd exactly when n is a power of 2.

2.6 Squaring modulo 2

Lemma 2.2 (Frobenius identity)**.** *Let p be a prime. For every power series U with coefficients in $\mathbb{F}_p$, we have $U(x)^p = U(x^p)$.*

Proof. In $\mathbb{F}_p$, the binomial theorem simplifies to $(a + b)^p = a^p + b^p$, because every binomial coefficient $\binom{p}{i}$ with $0 < i < p$ is divisible by p. Also $c^p = c$ for every $c \in \mathbb{F}_p$ (Fermat's little theorem). So $\left(\sum_n u_n x^n\right)^p = \sum_n u_n^p x^{pn} = \sum_n u_n x^{pn} = U(x^p)$. Only finitely many terms matter for any given coefficient, so this is a polynomial identity. □

For $p = 2$ this says that squaring a series modulo 2 just doubles every exponent: for instance, $(x + x^3)^2 \equiv x^2 + x^6 \pmod{2}$. In particular, the square of any series has, modulo 2, no terms of odd degree.

2.7 The Catalan numbers modulo 2

Lemma 2.3. *In every commutative ring the equation $U = x+U^2$ has exactly one solution with $U(0) = 0$. Over the integers it is the Catalan series $C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + \cdots$, and*

$$C(x) \equiv x + x^2 + x^4 + x^8 + \cdots = \sum_{k\geq 0} x^{2^k} \pmod 2.$$

So $[x^n]\,C$ is odd exactly when n is a power of 2.

Proof. The coefficient of x^n in U^2 involves only $u_1, \ldots, u_{n-1}$, so the equation determines u_n one at a time: $u_1 = 1$ and $u_n = \sum_{i=1}^{n-1} u_i u_{n-i}$. Over $\mathbb{F}_2$, the series $S = \sum_k x^{2^k}$ satisfies $x + S^2 = x + \sum_k x^{2^{k+1}} = S$ by Lemma 2.2, so by uniqueness $C \equiv S \pmod 2$. □

The parity of the Catalan numbers is classical; see for example [3]. (Here the Catalan number $\frac{1}{n}\binom{2n-2}{n-1}$ appears as the coefficient of x^n, so "n a power of 2" corresponds to the familiar statement that the k-th Catalan number is odd exactly when $k+1$ is a power of 2.)

3 Results

Throughout, C is the Catalan series, the unique normalized series with $C = x+C^2$ (Lemma 2.3), and $c_n = [x^n]\,C$. The defining equation of the sequence is

$$A(x - x^2) = x + A(x)^2. \tag{1}$$

Theorem 3.1. *(a) Equation* (1) *has exactly one normalized solution A with integer coefficients. (More generally, it has exactly one normalized solution with coefficients in any commutative ring.) The sequence A393172 is $a(n) = [x^n]\,A$.*

(b) Modulo 4*,*

$$A(x) \equiv 2C(x) - x \pmod 4.$$

Corollary 3.2 (Hanna's conjecture)**.** *For every $n > 1$: if n is a power of* 2 *then $a(n) \equiv 2 \pmod 4$, and otherwise $a(n)$ is divisible by* 4*.*

Proof. For $n > 1$ the coefficient of x^n in $2C(x) - x$ is $2c_n$, so Theorem 3.1(b) gives $a(n) \equiv 2c_n \pmod 4$. If c_n is odd, then $2c_n \equiv 2 \pmod 4$; if c_n is even, then $2c_n \equiv 0 \pmod 4$. By Lemma 2.3, c_n is odd exactly when n is a power of 2. □

For example, $c_8 = 429$ is odd, and indeed $a(8) = 112290 = 4 \cdot 28072 + 2$; while $c_5 = 14$ is even, and $a(5) = 260 = 4 \cdot 65$.

4 Proofs

The idea in brief. The substitution $x \mapsto x - x^2$ is undone by the Catalan series, because $C(x) - C(x)^2 = x$. Substituting C into (1) therefore turns it into the equation $A = C + A(C)^2$, which fixes the coefficients of A one at a time; so there is exactly one solution, in every commutative ring. For the congruence we guess the answer $D = 2C - x$ and check that it satisfies (1) modulo 4. Two short computations do this: substituting $x - x^2$ into D gives exactly $x + x^2$, and squaring D gives x^2 plus multiples of 4. By uniqueness, A and D agree modulo 4.

Lemma 4.1 (An equation with a square)**.** *Let T and P be normalized series with coefficients in a commutative ring R. There is exactly one normalized series G with coefficients in R such that*

$$G(x) = T(x) + \big(G(P(x))\big)^2.$$

If T and P have integer coefficients, then so does G, and reducing G modulo m gives the solution of the same equation over $\mathbb{Z}/m\mathbb{Z}$ (with T and P reduced modulo m).

Proof. The point is that the coefficient of x^n on the right-hand side depends only on the coefficients of G below degree n. To see this, let G and H be two series with constant term 0 that agree below degree r, so $G - H = O(x^r)$. Since P starts with x, the power P^k has no terms below x^k, and so $G(P) - H(P) = O(x^r)$ as well. Moreover $G(P) + H(P)$ has constant term 0. Hence

$$G(P)^2 - H(P)^2 = \big(G(P) - H(P)\big)\big(G(P) + H(P)\big) = O(x^{r+1}).$$

In other words, changing G in degree r and above changes the right-hand side only in degree $r + 1$ and above.

So the equation fixes the coefficients of G one at a time. The constant term is 0, the coefficient of x is the coefficient of x in T, which is 1, and for $n \geq 2$ the coefficient of x^n in G must equal the coefficient of x^n in $T + G(P)^2$, which has already been computed from the earlier coefficients. Only additions and multiplications are used, so this works in every commutative ring, and gives integer coefficients when T and P have them. Finally, reducing the integer solution modulo m gives a normalized series that satisfies the reduced equation (reduction is compatible with all the operations involved), and by uniqueness it is *the* solution over $\mathbb{Z}/m\mathbb{Z}$. □

Lemma 4.2 (Two forms of the equation)**.** *Let R be a commutative ring and A a normalized series over R. Then A satisfies* (1) *if and only if*

$$A(x) = C(x) + \big(A(C(x))\big)^2. \tag{2}$$

Proof. Write $N(x) = x - x^2$. Then $N(C(x)) = C - C^2 = x$. By Lemma 2.1, C is the compositional inverse of N, so also $C(N(x)) = x$. We use two simple facts about substitution: it can be done in steps, $(U \circ V) \circ W = U \circ (V \circ W)$; and substituting into a sum or a product is the same as adding or multiplying the substituted terms.

Suppose $A(N) = x + A^2$. Substitute C for x on both sides. The left side becomes $A(N(C)) = A(x)$, and the right side becomes $C + A(C)^2$. This is (2). Conversely, if (2) holds, substitute N for x: the left side becomes $A(N)$, and the right side becomes $C(N) + A(C(N))^2 = x + A^2$. This is (1). □

Proof of Theorem 3.1. (a) By Lemma 4.2, the normalized solutions of (1) are exactly the normalized solutions of (2). By Lemma 4.1 with $T = P = C$, there is exactly one, in every commutative ring. Over the integers it has integer coefficients, because C does.

(b) By part (a) applied to the ring $\mathbb{Z}/4\mathbb{Z}$, and the last sentence of Lemma 4.1, the reduction of A modulo 4 is the only normalized series over $\mathbb{Z}/4\mathbb{Z}$ that satisfies (1). So it is enough to check that the normalized series $D = 2C - x$ satisfies (1) in $\mathbb{Z}/4\mathbb{Z}$. On the left side, using $C(x - x^2) = x$,

$$D(x - x^2) = 2\,C(x - x^2) - (x - x^2) = 2x - x + x^2 = x + x^2.$$

On the right side,

$$x + D^2 = x + (2C - x)^2 = x + 4C^2 - 4xC + x^2 = x + x^2 \quad \text{in } \mathbb{Z}/4\mathbb{Z},$$

because the terms $4C^2$ and $4xC$ vanish modulo 4. The two sides agree, so $A \equiv D = 2C - x \pmod 4$. □

5 Checking the results

A mathematical proof can contain mistakes, so the results were checked in two independent ways.

Numerical check. We solved (1) coefficient by coefficient, with exact integer arithmetic. This reproduces all 24 terms printed in the OEIS entry. We then checked that every coefficient of $A(x) - (2C(x) - x)$ below degree 200 is divisible by 4, and that the rule of Corollary 3.2 holds for every n with $1 < n < 200$. The computation was done with the PARI/GP computer algebra system, using code that was written separately from the formal proofs (file `verification/independent_checks.gp` in the accompanying repository [9]). A computation can only test finitely many cases, so it is a check on the definitions and on the indexing, not a proof.

Formal proof in Lean. Lean [7] is a *proof assistant*: a program in which mathematical statements and their proofs are written in a precise formal language, and every step is checked mechanically down to the basic rules of logic. All results of this paper are proved in Lean 4 (version 4.19.0) on top of Mathlib [8], Lean's library of formalized mathematics. The statements are collected in the file `lean/Entries/A393172.lean` of the repository, in the namespace `OEIS.A393172`. There the Catalan series is called `catalan`, and the main proofs are in the module `lean/Families/CatalanComposition.lean`. The table lists where each result of this paper is proved.

Statement in this paper	Lean theorem(s)
Theorem 3.1(a) for integer coefficients; A is normalized and satisfies (1)	`first_existence`, `first_definition`
Theorem 3.1(b): $A \equiv 2C - x \pmod 4$	`complete_series_mod_four`
Corollary 3.2, also for any normalized integer solution of (1)	`first_residue`, `divisibility_complement`, `first_arbitrary_source`

The Lean project was rebuilt from source for this paper, and Lean was asked to list the basic assumptions (axioms) that each of these theorems ultimately depends on. Every one depends only on the three standard axioms of Lean's logic (`propext`, `Classical.choice`, `Quot.sound`); no step is left unproved. A formal proof certifies exactly the statement that Lean checks. Whether that statement says what the OEIS entry means is a separate question, so the Lean definitions were also compared, side by side, with the definition in the OEIS entry (Section 1.1).

6 Discussion

Scope. Theorem 3.1 determines the remainder of every coefficient $a(n)$ modulo 4, so it settles the conjecture of the entry completely. It says nothing about the size of $a(n)$, and nothing about remainders modulo higher powers of 2.

Modulo 8. A computer calculation suggests a similar, but more intricate, rule modulo 8. For every n with $1 < n < 200$ we found: $a(n) \equiv 2 \pmod 8$ when n is a power of 2; $a(n) \equiv 4 \pmod 8$ when $n = 2^k\,(4^j + 1)$ for some $j \geq 1$ and $k \geq 0$ (that is, $n = 5, 10, 17, 20, 34, 40, 65, 68, 80, \ldots$); and $a(n) \equiv 0 \pmod 8$ otherwise. We have not tried to prove this. A proof would probably need finer information about the Catalan numbers, namely their remainders modulo 4.

The companion sequence. The entry A393171 [2] is about the series $B = x + A^2$. Squaring is useful here: if $A = D + 2^k K$ for some series K with integer coefficients and some $k \geq 1$, then $A^2 = D^2 + 2^{k+1}DK + 2^{2k}K^2$, so A^2 is known modulo 2^{k+1} once A is known modulo 2^k. In this way Theorem 3.1 determines B modulo 8, which proves Hanna's conjecture for A393171 (see the companion paper). The rule modulo 8 suggested above would, in the same way, determine B modulo 16.

About this paper

How the work was done. This paper is one of 26 papers on conjectures from the OEIS that came out of an autonomous AI research effort directed by Dev D. Goyal [9]. The roles were as follows.

- *Dev D. Goyal* started the project, chose which problems to pursue, asked for proofs, formal verification and revisions, and decided what to publish. Dev D. Goyal did not supply proof strategies or intermediate arguments.
- *ChatGPT* (OpenAI), using the GPT-6 model with Pro reasoning, worked autonomously from those prompts. It searched the OEIS for suitable conjectures, found the proofs, wrote the Lean 4 formalization, and wrote a first draft of this paper.
- *Claude Opus 5.5*, used with Claude Code, checked the work independently and then rewrote this paper. It rebuilt the Lean project from source and confirmed that the theorems listed in Section 5 rely only on Lean's standard axioms; compared the Lean definitions with the OEIS definition; recomputed the sequence with separately written PARI/GP code, compared it with the terms published in the OEIS, and tested every claim of the paper on many terms; and read every step of the proofs. It then rewrote the text to make it self-contained and readable. The theorems and the proof strategies are ChatGPT's. In rewriting, Claude Opus 5.5 expanded the explanations and, in places, wrote out an argument in more detail or by a more elementary route. The Discussion section was added by Claude Opus 5.5; the further observations in it were checked by computer and are not part of the Lean formalization.

Dev D. Goyal has verified the results and is responsible for this paper.

Review status. This paper has not been peer reviewed, and it has not been submitted to a journal or to the OEIS.

Priority. An OEIS entry that labels a statement as a conjecture shows only that its contributor did not know a proof; it does not show that no proof existed. During the research only limited searches for earlier work were made. No claim of priority is made.

Corrections. Please report any error in the issue tracker of the repository [9], where the Lean sources and the numerical checks are also available.

Acknowledgements

The sequence, its formulas and the conjecture are due to Paul D. Hanna [1]. The recurrence in the entry was contributed by Seiichi Manyama.

References

[1] P. D. Hanna, entry A393172 (G.f. $A(x)$ satisfies $A(x - x^2) = x + A(x)^2$), *The On-Line Encyclopedia of Integer Sequences*, February 2026. `https://oeis.org/A393172`. Text as of October 2026.

[2] P. D. Hanna, entry A393171 (G.f. satisfies $A(x) = x + A(C(x))^2$, where $C(x) = x + C(x)^2$), *The On-Line Encyclopedia of Integer Sequences*, February 2026. `https://oeis.org/A393171`. Text as of October 2026.

[3] E. Deutsch and B. E. Sagan, Congruences for Catalan and Motzkin numbers and related sequences, *Journal of Number Theory* 117 (2006), 191–215.

[4] N. J. A. Sloane and The OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*. `https://oeis.org`.

[5] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999. (Chapter 5 treats formal power series, composition and compositional inverses.)

[6] I. Niven, Formal power series, *American Mathematical Monthly* 76 (1969), 871–889.

[7] L. de Moura and S. Ullrich, The Lean 4 theorem prover and programming language, in *Automated Deduction – CADE 28*, Lecture Notes in Computer Science 12699, Springer, 2021, 625–635.

[8] The mathlib Community, The Lean mathematical library, in *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, 367–381. Version used: commit `c44e0c8ee63c`.

[9] D. D. Goyal, *Proofs and corrections of OEIS conjectures, with Lean 4 formalizations*, repository, 2026. `https://github.com/dvgyl/oeis-a396843-proof`.