Congruences for the series A with $A\big(x - x\,A(5x)/5\big) = x$

A proof of a conjecture in OEIS A393857

Dev D. Goyal

October 2026

Abstract

The sequence A393857 of the On-Line Encyclopedia of Integer Sequences (OEIS) lists the coefficients of the power series $A(x) = x + x^2 + 7x^3 + 205x^4 + \cdots$ defined by the equation $A\big(x - x\,A(5x)/5\big) = x$. The unknown series A occurs both as the outer function and inside its own argument, and its coefficients grow rapidly. Paul D. Hanna conjectured that every coefficient leaves remainder 1 on division by 6. We prove this. In fact we prove that, with 5 replaced by any positive integer q, the series becomes $x + x^2 + x^3 + \cdots$ once its coefficients are reduced modulo $q + 1$. The proof rests on the congruence $q \equiv -1 \pmod{q+1}$: for $q = -1$ the equation has an explicit solution. The results have also been checked by computer and proved formally in the Lean proof assistant.

How this paper was produced. This paper reports the result of an AI research effort directed by Dev D. Goyal. ChatGPT (OpenAI), using the GPT-6 model with Pro reasoning and working autonomously from prompts by Dev D. Goyal, found the proof, wrote its Lean 4 formalization, and wrote a first draft of the paper. Dev D. Goyal chose the problems and asked for proofs, formal verification and revisions, but did not supply proof ideas. Claude Opus 5.5, used with Claude Code, then checked the work independently: it rebuilt the formal proofs from source and confirmed that they rely only on Lean's three standard axioms, recomputed the sequence from its definition compared it with the published terms and tested every claim numerically, and read every step of the proofs. It also wrote the present version of this paper. Dev D. Goyal has reviewed the proofs and is responsible for them; this review is not an independent expert validation. The paper has not been peer reviewed. The section "About this paper" at the end gives the details.

1 Introduction

1.1 The sequence

The *On-Line Encyclopedia of Integer Sequences* (OEIS) [2] is a large public database of integer sequences. Many entries also record *conjectures*: patterns that the contributor observed in the data but did not prove.

This paper is about entry A393857 [1], contributed by Paul D. Hanna in March 2026. It defines a power series

$$A(x) = a(1)\,x + a(2)\,x^2 + a(3)\,x^3 + \cdots$$

by the equation

$$A\big(x - x\,A(5x)/5\big) = x. \tag{1}$$

Here $A(5x)$ is the series A with x replaced by $5x$, so its coefficient of x^n is $5^n\,a(n)$. Dividing by 5 gives the series whose coefficients are $5^{\,n-1}a(n)$, which are still integers. The left side of

(1) means: substitute the series $x - x\,A(5x)/5$ into A. (Section 2 explains what substituting one series into another means.) So the equation says that A undoes a substitution that is itself built from A. Because A occurs inside its own argument, it is not obvious that the equation has a solution, or only one; Theorem 3.1 shows that it has exactly one solution of the form $x +$ (higher powers of x).

The first terms, starting at $n = 1$, are 1, 1, 7, 205, 26869, 16984687, 53213653657, ... They grow rapidly: $a(15)$ already has 64 digits. The entry reads:

```
G.f.  A(x) satisfies A( x - x*A(5*x)/5 ) = x.

Conjecture:  a(n) == 1 (mod 6) for n >= 1.
```

— OEIS A393857, Paul D. Hanna, March 2026

Here "G.f." stands for *generating function*, the series whose coefficients are the terms of the sequence. The entry also records other formulas for $A(x)$, for example as an infinitely nested sequence of substitutions; we do not need them.

1.2 The question

Hanna conjectured, from the computed terms, that every $a(n)$ leaves remainder 1 on division by 6. The table shows the remainders of the first 32 terms modulo 6.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$a(n)$ mod 6	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
n	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
$a(n)$ mod 6	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

1.3 What this paper proves

Main result (Corollary 3.4). *The conjecture is true:* $a(n) \equiv 1 \pmod 6$ *for every* $n \geq 1$.

The number 5 plays no special role. Hanna contributed a family of such sequences, in which 5 is replaced by another positive integer q, and we treat all of them at once. Theorem 3.2 shows that, for every $q \geq 1$, all coefficients leave remainder 1 on division by $q + 1$. Theorem 3.3 shows that, for every even q, a coefficient is odd exactly when its index is a power of 2.

The proof rests on two observations. Modulo $q + 1$ we have $q \equiv -1$, and for $q = -1$ the equation has the explicit solution $x + x^2 + x^3 + \cdots$. Similarly, modulo 2 an even q is congruent to 0, and for $q = 0$ the equation becomes the classical equation of the Catalan numbers, whose parity is well known.

1.4 What was known

Hanna contributed the four entries A393856–A393859, for $q = 4, 5, 6, 7$, in March 2026, each with the conjectures for its own value of q. The OEIS also lists the members for $q = 0, 1, 2, 3$, which were contributed earlier. For $q = 0$, where $A(qx)/q$ is replaced by x (its value at $q = 0$ according to formula (2) below), the equation becomes $A(x - x^2) = x$; its solution is the generating series of the Catalan numbers (A000108). The cases $q = 1, 2, 3$ are A088714, A195195 and A195196. We are not aware of an earlier proof of the conjectures. The tools we use, formal power series and their reduction modulo an integer, are standard; see [4, 3].

1.5 Motivation

Equations in which an unknown series is substituted into an expression built from itself are common in combinatorics, where they describe structures defined recursively, such as trees. The Catalan numbers, the case $q = 0$ of this family, are the classic example. Hanna has contributed many such equations to the OEIS. For $q \geq 1$ the coefficients grow rapidly and no closed formula for them is known to us, yet their remainders follow simple rules. The proof explains these rules: working modulo $q + 1$ replaces the parameter q by -1, and, for even q, working modulo 2 replaces it by 0; at these two values the equation can be solved explicitly. The same approach, reduction modulo a number for which the equation simplifies, is used in other papers of this collection.

1.6 How to read this paper

Section 2 explains the few notions we need: power series, substitution, and remainders. Section 3 makes the equation precise for every q and states the results, and Section 4 proves them. Section 5 describes how the results were checked, by computer and by a formal proof in Lean. Section 6 discusses the scope of the results, a further congruence, and open questions.

2 Background

This section collects the definitions and facts used later. Readers who know formal power series and congruences can skip it and return when needed.

2.1 Formal power series

A *formal power series* is an expression

$$U(x) = u_0 + u_1 x + u_2 x^2 + u_3 x^3 + \cdots ,$$

determined by its sequence of coefficients $u_0, u_1, u_2, \ldots$; see [4, 3]. No number is ever substituted for x, so questions of convergence do not arise. We write $[x^n]\, U = u_n$ for the coefficient of x^n in U; two series are equal when all their coefficients are equal.

Series are added coefficient by coefficient and multiplied like polynomials:

$$[x^n]\,(UV) = u_0 v_n + u_1 v_{n-1} + \cdots + u_n v_0.$$

Each coefficient of a product is a finite sum. For example, $(1 - x)(1 + x + x^2 + x^3 + \cdots) = 1$, so $1 + x + x^2 + \cdots$ is the *multiplicative inverse* of $1 - x$, written $\frac{1}{1-x}$. More generally, every series D with constant term 1 has a unique multiplicative inverse $\frac{1}{D}$, whose coefficients are determined one at a time by $D \cdot \frac{1}{D} = 1$ using only addition, subtraction and multiplication. Whenever a quotient of power series appears in this paper, its denominator has constant term 1, so the quotient is again a power series.

A *commutative ring* is a set with an addition, a subtraction and a multiplication that satisfy the usual rules of arithmetic; examples are $\mathbb{Z}$, $\mathbb{Q}$ and the integers modulo m. For a commutative ring R we write $R[[x]]$ for the set of power series with coefficients in R; for instance, writing $\mathbb{F}_2$ for the integers modulo 2, $\mathbb{F}_2[[x]]$ consists of the series with coefficients modulo 2. We write $U(0)$ for the constant term u_0 of U, and $U = O(x^d)$ when the coefficients of $1, x, \ldots, x^{d-1}$ in U are all zero. Thus $U - V = O(x^d)$ means that U and V have the same coefficient of x^n for every $n < d$. A series with constant term 0 and coefficient of x equal to 1, that is, a series of the form $x + u_2 x^2 + u_3 x^3 + \cdots$, is called *normalized*.

2.2 Substituting one series into another

If $V(x)$ has constant term 0, we can substitute V for x in U:

$$U(V(x)) = u_0 + u_1 V(x) + u_2 V(x)^2 + u_3 V(x)^3 + \cdots .$$

This is called *composition*, also written $U \circ V$. It makes sense because $V(x)^k$ has no terms below x^k, so only the finitely many terms with $k \le n$ affect the coefficient of x^n. For example, if $U = x + x^2$ and $V = x^2$, then $U(V(x)) = x^2 + x^4$. Composition is associative: $U \circ (V \circ W) = (U \circ V) \circ W$.

Substituting a series into itself repeatedly gives its *iterates*: $U^{\circ 2}(x) = U(U(x))$, $U^{\circ 3}(x) = U(U(U(x)))$, and so on, with $U^{\circ 0}(x) = x$ and $U^{\circ 1} = U$. They must not be confused with *powers* $U^2 = U \cdot U$, $U^3 = U \cdot U \cdot U$. (The OEIS sometimes writes $A^k(x)$ for the k-th iterate. In this paper the small circle always means an iterate, and a plain exponent always means a power.)

2.3 Undoing a substitution

A normalized series can always be "undone" by another normalized series.

Lemma 2.1 (Compositional inverse)**.** *For every normalized series U there is exactly one normalized series V with $U(V(x)) = x$, and this V also satisfies $V(U(x)) = x$. If U has integer coefficients, so does V.*

We call V the *compositional inverse* of U. The OEIS calls it the *series reversion* of U. For example, the compositional inverse of $x - x^2$ is $C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + \cdots$, whose coefficients are the Catalan numbers. Indeed $C - C^2 = x$, which says exactly that substituting C into $x - x^2$ gives x.

Proof. Write $V = x + v_2 x^2 + v_3 x^3 + \cdots$ with unknown coefficients. In $U(V) = V + u_2 V^2 + u_3 V^3 + \cdots$, the coefficient of x^n (for $n \ge 2$) equals v_n plus terms that involve only $v_2, \ldots, v_{n-1}$. Setting it equal to 0 therefore determines v_n from the earlier coefficients, using only additions, subtractions and multiplications. So there is exactly one V with $U \circ V = x$, and it has integer coefficients if U does. Applying the same construction to V gives a normalized W with $V \circ W = x$. Then $U = U \circ (V \circ W) = (U \circ V) \circ W = W$, so $V \circ U = V \circ W = x$ as well. □

2.4 Remainders and congruences

For an integer $m \ge 2$ we write $a \equiv b \pmod m$ ("a is congruent to b modulo m") when $a - b$ is divisible by m, that is, when a and b leave the same remainder after division by m. The remainder of a on division by m is called its *residue* modulo m; it is one of $0, 1, \ldots, m-1$.

Addition and multiplication are compatible with congruences, so the residues $0, 1, \ldots, m-1$, added and multiplied modulo m, form a ring, written $\mathbb{Z}/m\mathbb{Z}$. When $m = p$ is a prime we also write $\mathbb{F}_p$. In $\mathbb{F}_2 = \{0, 1\}$ we have $1 + 1 = 0$, so addition and subtraction coincide.

For power series with integer coefficients, $U \equiv V \pmod m$ means that *every* coefficient of $U - V$ is divisible by m. Equivalently, the two series become equal when every coefficient is replaced by its residue; we call this *reducing* the series modulo m. The key fact is the following.

Reduction modulo m is compatible with all the operations above. To reduce a sum, a product, a substitution, or an inverse, we may reduce the ingredients first and then compute in $\mathbb{Z}/m\mathbb{Z}$.

This holds because every coefficient of the result is obtained from finitely many coefficients of the ingredients by adding, subtracting and multiplying, and these operations are compatible with remainders.

Rings. Some statements below hold over every commutative ring, for example $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{Z}/m\mathbb{Z}$. No division is used in them; this matters because in $\mathbb{Z}/m\mathbb{Z}$ division is not always possible (for instance, 2 has no inverse in $\mathbb{Z}/4\mathbb{Z}$).

2.5 The method

The main proofs follow three steps.

1. *Uniqueness.* Show that the defining equation determines the coefficients one at a time, using only addition, subtraction and multiplication. Then the equation has exactly one solution of the required shape, in every commutative ring, including $\mathbb{Z}/m\mathbb{Z}$.
2. *Reduction.* Reducing the integer solution modulo m gives a solution of the reduced equation, because reduction is compatible with all operations.
3. *A candidate solution.* Find an explicit series that satisfies the reduced equation. By uniqueness, it is the reduction of the integer solution, and its coefficients can be read off.

Finding the candidate in step 3 is usually the main difficulty.

3 Results

To treat the whole family at once, we replace 5 by a parameter q. First we make the equation precise. For a series $U = u_1x + u_2x^2 + u_3x^3 + \cdots$ without constant term and a number $q \neq 0$, the series $U(qx)/q$ has the coefficients $q^{n-1}u_n$. We take this formula as the definition, for every q:

$$L_q(U) = \sum_{n\geq 1} q^{n-1}u_n\, x^n = u_1x + q\,u_2x^2 + q^2u_3x^3 + \cdots . \tag{2}$$

No division by q occurs, so $L_q(U)$ makes sense for every q, including $q = 0$ (where it equals u_1x), and for coefficients in every commutative ring, such as $\mathbb{Z}/m\mathbb{Z}$. The equation of the family is

$$A(x - x\,L_q(A)) = x, \qquad A \text{ normalized.} \tag{3}$$

For $q = 5$ this is equation (1) of the entry.

Theorem 3.1 (Exactly one solution)**.** *Let q be an element of a commutative ring R. There is exactly one normalized series G with coefficients in R that satisfies* (3)*. When $R = \mathbb{Z}$ and q is an integer, we call this solution G_q. For every $m \geq 2$, the reduction of G_q modulo m is the solution of* (3) *over $\mathbb{Z}/m\mathbb{Z}$, with q replaced by its residue.*

So the sequence of OEIS A393857 is $a(n) = [x^n]\,G_5$.

Theorem 3.2 (All coefficients are 1 modulo $q+1$)**.** *For every integer $q \geq 1$,*

$$G_q(x) \equiv \frac{x}{1-x} = x + x^2 + x^3 + \cdots \pmod{q+1}.$$

That is, every coefficient $[x^n]\,G_q$ with $n \geq 1$ leaves remainder 1 on division by $q+1$.

Theorem 3.3 (Parity)**.** *If q is even, then*

$$G_q(x) \equiv x + x^2 + x^4 + x^8 + \cdots = \sum_{k\geq 0} x^{2^k} \pmod 2,$$

so $[x^n]\,G_q$ is odd if and only if n is a power of 2. If q is odd, every $[x^n]\,G_q$ with $n \geq 1$ is odd.

Corollary 3.4 (Hanna's conjecture)**.** *For the sequence of OEIS A393857: $a(n) \equiv 1 \pmod 6$ for every $n \geq 1$.*

4 Proofs

The idea in brief. The equation determines the coefficients of A one at a time, without any division, so it has exactly one solution over every commutative ring, including $\mathbb{Z}/(q+1)\mathbb{Z}$ and $\mathbb{Z}/2\mathbb{Z}$ (Theorem 3.1). Modulo $q+1$ we may replace q by -1, and then a direct calculation shows that $x + x^2 + x^3 + \cdots$ is a solution. Modulo 2, for even q, we may replace q by 0, and then the equation says that A undoes the substitution $x \mapsto x - x^2$, which is exactly what the Catalan series does. By uniqueness, these explicit series are the reductions of G_q.

Proof of Theorem 3.1. Write $G = x + g_2x^2 + g_3x^3 + \cdots$ with unknown coefficients, and put $V = x - x\,L_q(G)$. By (2), the coefficient g_k enters $L_q(G)$ only in degree k, and therefore enters V only in degree $k+1$, because of the factor x. Also, $V = x + O(x^2)$ is normalized. Now look at the coefficient of x^n, for some $n \geq 2$, in

$$G(V) = V + g_2V^2 + g_3V^3 + \cdots + g_nV^n + \cdots .$$

- The term g_nV^n contributes exactly g_n, because $V^n = x^n + O(x^{n+1})$.
- The terms g_kV^k with $k > n$ contribute nothing, because V^k has no terms below x^k.
- The remaining terms, V and g_kV^k with $2 \leq k < n$, involve g_k with $k < n$, and the coefficients of V up to degree n, which involve only $g_2, \ldots, g_{n-1}$.

So the coefficient of x^n in $G(V)$ is g_n plus an expression in $q, g_2, \ldots, g_{n-1}$ built with additions, subtractions and multiplications. Setting it equal to the coefficient of x^n in x, which is 0, determines g_n. Doing this for $n = 2, 3, 4, \ldots$ in turn gives exactly one solution, in every commutative ring.

Reduction modulo m is compatible with all the operations in (3): with L_q, which multiplies the coefficients by powers of q, with products, and with substitution. So the reduction of G_q satisfies the reduced equation, and by the uniqueness just proved (over $\mathbb{Z}/m\mathbb{Z}$) it is its solution. □

Proof of Theorem 3.2. We work over $\mathbb{Z}/(q+1)\mathbb{Z}$, where $q = -1$. Let $H = x/(1-x) = x + x^2 + x^3 + \cdots$. All coefficients of H (except the constant term) are 1, so by (2)

$$L_{-1}(H) = x - x^2 + x^3 - x^4 + \cdots = \frac{x}{1+x}, \qquad \text{hence} \qquad x - x\,L_{-1}(H) = x - \frac{x^2}{1+x} = \frac{x}{1+x}.$$

Substituting this into H gives

$$H\Big(\frac{x}{1+x}\Big) = \frac{x/(1+x)}{1 - x/(1+x)} = \frac{x/(1+x)}{1/(1+x)} = x.$$

So H satisfies the reduced equation. By Theorem 3.1, H is the reduction of G_q modulo $q+1$. Every coefficient of H of positive degree is 1. □

Proof of Theorem 3.3. If q is odd, then $q+1$ is even. By Theorem 3.2, every coefficient of positive degree is 1 plus a multiple of $q+1$, hence odd.

Now let q be even, and work over $\mathbb{F}_2$, where $q = 0$. By (2), $L_0(U) = u_1 x$, which is x for a normalized U, so over $\mathbb{F}_2$ the equation becomes $G(x - x^2) = x$. Thus G is the compositional inverse of $x - x^2$ (Lemma 2.1), which is the reduction of the Catalan series C (Section 2). It is a classical fact that $C \equiv x + x^2 + x^4 + x^8 + \cdots \pmod 2$: indeed, $S = \sum_k x^{2^k}$ satisfies $S - S^2 = x$ over $\mathbb{F}_2$, because squaring a series over $\mathbb{F}_2$ doubles all its exponents (there $(a+b)^2 = a^2 + 2ab + b^2 = a^2 + b^2$ and $c^2 = c$), so $S^2 = \sum_{k\geq 1} x^{2^k}$; and the equation $U - U^2 = x$ determines U uniquely, one coefficient at a time. Hence $G_q \equiv S \pmod 2$. □

Corollary 3.4 is the case $q = 5$ of Theorems 3.2 and 3.3. (For odd q the parity statement adds nothing new: it already follows from the congruence modulo $q+1$. Since $6 = 2 \cdot 3$, the corollary also says that every $a(n)$ is odd and leaves remainder 1 on division by 3.)

5 Checking the results

A mathematical proof can contain mistakes, so the results were checked in two independent ways.

Numerical check. We solved (1) coefficient by coefficient with exact integer arithmetic, up to $n = 25$; this reproduces all 14 terms printed in the OEIS entry. We then ran the same computation modulo $12 = 2(q+1)$, which is legitimate because the construction uses only additions, subtractions and multiplications (Theorem 3.1). It agrees with the exact values for $n \leq 25$, and it confirms the congruence of Corollary 3.4 for every $n < 300$. The computation was done with the PARI/GP computer algebra system, using code that was written separately from the formal proofs (file `verification/independent_checks.gp` in the accompanying repository [7]). A computation can only test finitely many cases, so it is a check on the definitions and on the indexing, not a proof.

Formal proof in Lean. Lean [5] is a *proof assistant*: a program in which mathematical statements and their proofs are written in a precise formal language, and every step is checked mechanically down to the basic rules of logic. The theorems of Section 3 are proved in Lean 4 (version 4.35.0) on top of Mathlib [6], Lean's library of formalized mathematics. The statements are collected in the file `lean/Entries/A393857.lean` of the repository, in the namespace `OEIS.A393857`. In Lean, the operation L_q of (2) is called `scaleTail q`. The general theorems for every q are `integer_mod_parameter_add_one` (Theorem 3.2) and `integer_coeff_odd_iff` (Theorem 3.3 for even q), in the namespace `Hanna.Related.ScaledInverse`, in the files `lean/Hanna/Related/ScaledInverse.lean` and `ScaledParity.lean` of the same folder. The table lists where each result of this paper is proved.

Statement in this paper	Lean theorem(s)
Theorem 3.1, over every commutative ring	`source_existence_uniqueness`
G_5 is normalized and satisfies (1)	`source_normalized`, `source_equation`
Corollary 3.4: $a(n) \equiv 1 \pmod 6$ for $n \geq 1$	`a393857_mod_six`

The Lean project was rebuilt from source, and the axioms on which each of these theorems depends were listed with Lean's `#print axioms` command. Every one depends only on

the three standard axioms of Lean's logic (`propext`, `Classical.choice`, `Quot.sound`); no step is left unproved. A formal proof certifies exactly the statement that Lean checks. Whether that statement says what the OEIS entry means is a separate question, so the Lean definitions were also compared with the definition in the OEIS entry (Section 1.1). The principal theorems are stated once more, using only Mathlib's definitions, in the file `lean/Challenges/ScaledInverses.lean`, and Lean's Comparator tool confirms that the proved theorems have exactly these statements.

6 Discussion

Scope. Corollary 3.4 settles the conjecture of the entry completely, for every index n. The theorems describe the remainders modulo $q+1$ and modulo 2; they say nothing about the size of the coefficients.

A further congruence. The argument used for Theorem 3.3 works modulo q itself. In arithmetic modulo q the parameter q is 0, so the equation becomes $G(x-x^2)=x$ again, and G_q agrees modulo q with the Catalan series:

$$[x^n]\,G_q \equiv \frac{1}{n}\binom{2n-2}{n-1} \pmod{q} \qquad (n \geq 1).$$

For example, for $q=5$ and $n=5$, the term $a(5)=26869$ and the Catalan number $\frac{1}{5}\binom{8}{4}=14$ both leave remainder 4 on division by 5. Since q and $q+1$ have no common factor, a number is determined modulo $q(q+1)$ by its remainders modulo q and modulo $q+1$ (the Chinese remainder theorem). So, together with Theorem 3.2, this determines every coefficient of G_q modulo $q(q+1)=30$ when $q=5$. We checked this congruence by computer for $q=2,3,\ldots,7$ and every $n<300$. It is not part of the Lean formalization.

The rest of the family. The theorems hold for every $q \geq 1$. The OEIS lists the members $q=1,2,3$ as A088714, A195195 and A195196; we checked that their published terms agree with the coefficients of G_1, G_2 and G_3 (for A088714 after shifting the index by one). For instance, A195195 is defined by the case $q=2$ of the family equation (3), so our results show that all its terms leave remainder 1 on division by 3, and that a term is odd exactly when its index is a power of 2.

Open questions. We have not studied the remainders modulo other numbers, such as $36=(q+1)^2$ for $q=5$. It would be interesting to know whether they are also given by an explicit series.

About this paper

How the work was done. This paper is one of 28 papers on conjectures from the OEIS that came out of an AI research effort directed by Dev D. Goyal [7]. The roles were as follows.

- *Dev D. Goyal* started the project, chose which problems to pursue, asked for proofs, formal verification and revisions, and decided what to publish. Dev D. Goyal did not supply proof strategies or intermediate arguments.
- *ChatGPT* (OpenAI), using the GPT-6 model with Pro reasoning, worked autonomously from those prompts. It searched the OEIS for suitable conjectures, found the proofs, wrote the Lean 4 formalization, and wrote a first draft of this paper.

- *Claude Opus 5.5*, used with Claude Code, checked the work independently. It rebuilt the Lean project from source and confirmed that the theorems listed in Section 5 rely only on Lean's standard axioms; compared the Lean definitions with the OEIS definition; recomputed the sequence with separately written PARI/GP code, compared it with the terms published in the OEIS, and tested every claim of the paper on many terms; and read every step of the proofs. It then wrote the present version of the paper, expanding the explanations and, in places, writing out an argument in more detail or by a more elementary route. The theorems of Section 3 and the proof strategies are ChatGPT's. The Discussion section, including the short arguments in it, was written by Claude Opus 5.5; its observations were checked by computer and are not part of the Lean formalization.

Dev D. Goyal has reviewed the proofs and is responsible for this paper; this review is not an independent expert validation.

Review status. This paper has not been peer reviewed, and it has not been submitted to a journal or to the OEIS.

Priority. An OEIS entry that labels a statement as a conjecture shows only that its contributor did not know a proof; it does not show that no proof existed. During the research only limited searches for earlier work were made. No claim of priority is made.

Corrections. Please report any error in the issue tracker of the repository [7], where the Lean sources and the numerical checks are also available.

Acknowledgements

The sequence and the conjecture are due to Paul D. Hanna [1], who contributed the entries A393856–A393859 ($q = 4, 5, 6, 7$), as well as the earlier members A088714, A195195 and A195196.

References

[1] P. D. Hanna, entry A393857 (G.f. $A(x)$ satisfies $A\big(x - x\,A(5x)/5\big) = x$), *The On-Line Encyclopedia of Integer Sequences*, March 2026. `https://oeis.org/A393857`. Text as of October 2026.

[2] N. J. A. Sloane and The OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*. `https://oeis.org`.

[3] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999. (Chapter 5 treats formal power series, composition and compositional inverses.)

[4] I. Niven, Formal power series, *American Mathematical Monthly* 76 (1969), 871–889.

[5] L. de Moura and S. Ullrich, The Lean 4 theorem prover and programming language, in *Automated Deduction – CADE 28*, Lecture Notes in Computer Science 12699, Springer, 2021, 625–635.

[6] The mathlib Community, The Lean mathematical library, in *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, 367–381. Version used: commit `56a005855ea1`.

[7] D. D. Goyal, *Proofs and corrections of OEIS conjectures*, repository, 2026. `https://github.com/dvgyl/oeis-proofs`.