Remainders modulo 8 of a series built from the Catalan numbers

A proof of a conjecture in OEIS A393171

Dev D. Goyal

October 2026

Abstract

The sequence A393171 of the On-Line Encyclopedia of Integer Sequences (OEIS), contributed by Paul D. Hanna, consists of the coefficients $a(n)$ of the power series $B(x) = x + x^2 + 4x^3 + 20x^4 + 116x^5 + \cdots$ defined by the equation $B(x) = x + B(C(x))^2$, where $C(x)$ is the generating function of the Catalan numbers. Hanna conjectured that, for $n > 2$, the coefficient $a(n)$ leaves remainder 4 on division by 8 when n or $n - 1$ is a power of 2, and is divisible by 8 otherwise. We prove this conjecture. The proof goes through the companion sequence A393172, whose series $A(x)$ satisfies $A(x - x^2) = x + A(x)^2$: we show that $B = x + A^2$, and that $A = 2C - x + 4K$ for a series K with integer coefficients. Squaring then gives B modulo 8 in closed form, $B \equiv x + x^2 + 4(C - x - xC)$, from which the conjecture can be read off. The result has also been checked by computer and proved formally in the Lean proof assistant.

How this paper was produced. This paper reports the result of an AI research effort directed by Dev D. Goyal. ChatGPT (OpenAI), using the GPT-6 model with Pro reasoning and working autonomously from prompts by Dev D. Goyal, found the proof, wrote its Lean 4 formalization, and wrote a first draft of the paper. Dev D. Goyal chose the problems and asked for proofs, formal verification and revisions, but did not supply proof ideas. Claude Opus 5.5, used with Claude Code, then checked the work independently: it rebuilt the formal proofs from source and confirmed that they rely only on Lean's three standard axioms, recomputed the sequence from its definition compared it with the published terms and tested every claim numerically, and read every step of the proofs. It also wrote the present version of this paper. Dev D. Goyal has verified the results and is responsible for them. The paper has not been peer reviewed. The section "About this paper" at the end gives the details.

1 Introduction

1.1 The sequence

The *On-Line Encyclopedia of Integer Sequences* (OEIS) [4] is a large public database of integer sequences. Many entries also record *conjectures*: patterns that the contributor observed in the data but did not prove.

This paper is about entry A393171 [1], contributed by Paul D. Hanna in February 2026. The entry reads, in part:

```
G.f.  satisfies A(x) = x + A(C(x))^2, where C(x) = x + C(x)^2 is a g.f.
of the Catalan numbers (A000108).

Conjecture:  for n > 2, a(n) == 4 (mod 8) iff n = 2^k or n = 2^k + 1 for k
>= 1, otherwise a(n) is divisible by 8.
```

```
(1) A(x-x^2) = x-x^2 + A(x)^2.
```

— OEIS A393171, Paul D. Hanna, February 2026

Here "G.f." stands for *generating function*: the terms $a(1), a(2), a(3), \ldots$ of the sequence are the coefficients of a power series. To avoid confusion with the equation numbers of this paper, we refer to the formula numbered (1) in the entry as (E1). The companion entry A393172 [2] is about a closely related series, which it also calls $A(x)$. To avoid a clash of names, in this paper we write

$$B(x) = a(1)\,x + a(2)\,x^2 + a(3)\,x^3 + \cdots$$

for the series of A393171, and we reserve the letter A for the series of A393172 (defined in Section 3). So the defining equation reads

$$B(x) = x + \big(B(C(x))\big)^2.$$

Here $B(C(x))$ is the series obtained by substituting $C(x)$ for x in B, and $B(C(x))^2$ is its square. Section 2 explains these operations. The first terms, starting at $n = 1$, are 1, 1, 4, 20, 116, 752, 5328, 40644, 330324, 2838296, ...

The equation involves the *Catalan numbers* $1, 1, 2, 5, 14, 42, 132, \ldots$ (OEIS A000108), one of the most common sequences in combinatorics. For example, they count the full binary trees with $1, 2, 3, 4, 5, \ldots$ leaves, that is, the rooted trees in which every internal vertex has exactly two children, a left one and a right one. Their generating function is

$$C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + 42x^6 + \cdots,$$

which is the only series without constant term that satisfies $C = x + C^2$. We write c_n for its coefficient of x^n.

1.2 The question

Hanna's conjecture says that for $n > 2$ the remainder of $a(n)$ on division by 8 is 4 when n is a power of 2 or one more than a power of 2, and 0 otherwise. The first row of the table shows the computed remainders of $a(n)$; the indices $n > 2$ of the form 2^k or $2^k + 1$ are printed in bold. The second row anticipates the main result: it shows the remainder of $4(c_n - c_{n-1})$ modulo 8.

n	1	2	**3**	**4**	**5**	6	7	**8**	**9**	10	11	12	13	14	15	**16**
$a(n)$ mod 8	1	1	**4**	**4**	**4**	0	0	**4**	**4**	0	0	0	0	0	0	**4**
$4(c_n - c_{n-1})$ mod 8	–	–	**4**	**4**	**4**	0	0	**4**	**4**	0	0	0	0	0	0	**4**

n	**17**	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**32**
$a(n)$ mod 8	**4**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**4**
$4(c_n - c_{n-1})$ mod 8	**4**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**4**

n	**33**	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
$a(n)$ mod 8	**4**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$4(c_n - c_{n-1})$ mod 8	**4**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

1.3 What this paper proves

Main result (Theorem 3.1). *Modulo* 8,

$$B(x) \equiv x + x^2 + 4\big(C(x) - x - x\,C(x)\big) \pmod 8.$$

In particular, $a(n) \equiv 4\,(c_n - c_{n-1}) \pmod 8$ *for every* $n > 2$.

The number $4(c_n - c_{n-1})$ leaves remainder 4 on division by 8 when $c_n - c_{n-1}$ is odd, and remainder 0 when it is even. A classical fact (Lemma 2.3) says that c_n is odd exactly when n is a power of 2. Together these prove Hanna's conjecture (Corollary 3.2). Along the way we prove the relations between the two entries: $B = x + A^2$, $B(C(x)) = A$, and formula (E1) of the entry, which in our notation reads $B(x - x^2) = x - x^2 + B(x)^2$.

1.4 What was known

The two OEIS entries record the relations between A and B as formulas, without proofs; they follow quickly from the definitions, and we prove them here (Theorem 3.1(b)). The parity of the Catalan numbers is classical; see for example [3]. We are not aware of an earlier proof of the conjecture. The conjecture of the companion entry A393172, about A modulo 4, is proved in a separate paper of this collection; to keep the present paper self-contained, we repeat that proof here (Proposition 4.2).

1.5 Motivation

Equations in which an unknown power series appears in a substitution, either as the inner or as the outer series, are called *functional equations*. They arise naturally when counting objects that are built recursively, such as trees, and the OEIS contains many sequences defined in this way. Their coefficients usually grow quickly and look irregular, but their remainders modulo small powers of 2 often show patterns. This example illustrates a technique for passing from one power of 2 to the next: a congruence modulo 4 for one series, written as an exact equation with integer coefficients, becomes a congruence modulo 8 for its square. The technique avoids division by 2, which is not possible in arithmetic modulo 8.

1.6 How to read this paper

Section 2 explains the notions we need: power series, substitution, remainders, and the Catalan numbers modulo 2. Section 3 states the results precisely, and Section 4 proves them. Section 5 describes how the results were checked, by computer and by a formal proof in Lean. Section 6 discusses the scope of the result and some open questions.

2 Background

This section collects the definitions and facts used later. Readers who know formal power series and congruences can skip it and return when needed.

2.1 Formal power series

A *formal power series* is an expression

$$U(x) = u_0 + u_1 x + u_2 x^2 + u_3 x^3 + \cdots ,$$

determined by its sequence of coefficients $u_0, u_1, u_2, \ldots$; see [6, 5]. No number is ever substituted for x, so questions of convergence do not arise. We write $[x^n]\,U = u_n$ for the coefficient of x^n in U; two series are equal when all their coefficients are equal.

Series are added coefficient by coefficient and multiplied like polynomials:

$$[x^n]\,(UV) = u_0v_n + u_1v_{n-1} + \cdots + u_nv_0.$$

Each coefficient of a product is a finite sum. For example, $(1-x)(1+x+x^2+x^3+\cdots) = 1$, so $1+x+x^2+\cdots$ is the *multiplicative inverse* of $1-x$, written $\frac{1}{1-x}$. More generally, every series D with constant term 1 has a unique multiplicative inverse $\frac{1}{D}$, whose coefficients are determined one at a time by $D \cdot \frac{1}{D} = 1$ using only addition, subtraction and multiplication. Whenever a quotient of power series appears in this paper, its denominator has constant term 1, so the quotient is again a power series.

A *commutative ring* is a set with an addition, a subtraction and a multiplication that satisfy the usual rules of arithmetic; examples are $\mathbb{Z}$, $\mathbb{Q}$ and the integers modulo m. For a commutative ring R we write $R[[x]]$ for the set of power series with coefficients in R; for instance, writing $\mathbb{F}_2$ for the integers modulo 2, $\mathbb{F}_2[[x]]$ consists of the series with coefficients modulo 2. We write $U(0)$ for the constant term u_0 of U, and $U = O(x^d)$ when the coefficients of $1, x, \ldots, x^{d-1}$ in U are all zero. Thus $U - V = O(x^d)$ means that U and V have the same coefficient of x^n for every $n < d$. A series with constant term 0 and coefficient of x equal to 1, that is, a series of the form $x + u_2x^2 + u_3x^3 + \cdots$, is called *normalized*.

2.2 Substituting one series into another

If $V(x)$ has constant term 0, we can substitute V for x in U:

$$U(V(x)) = u_0 + u_1V(x) + u_2V(x)^2 + u_3V(x)^3 + \cdots .$$

This is called *composition*, also written $U \circ V$. It makes sense because $V(x)^k$ has no terms below x^k, so only the finitely many terms with $k \le n$ affect the coefficient of x^n. For example, if $U = x + x^2$ and $V = x^2$, then $U(V(x)) = x^2 + x^4$. Composition is associative: $U \circ (V \circ W) = (U \circ V) \circ W$.

Substituting a series into itself repeatedly gives its *iterates*: $U^{\circ 2}(x) = U(U(x))$, $U^{\circ 3}(x) = U(U(U(x)))$, and so on, with $U^{\circ 0}(x) = x$ and $U^{\circ 1} = U$. They must not be confused with *powers* $U^2 = U \cdot U$, $U^3 = U \cdot U \cdot U$. (The OEIS sometimes writes $A^k(x)$ for the k-th iterate. In this paper the small circle always means an iterate, and a plain exponent always means a power.)

2.3 Undoing a substitution

A normalized series can always be "undone" by another normalized series.

Lemma 2.1 (Compositional inverse)**.** *For every normalized series U there is exactly one normalized series V with $U(V(x)) = x$, and this V also satisfies $V(U(x)) = x$. If U has integer coefficients, so does V.*

We call V the *compositional inverse* of U. The OEIS calls it the *series reversion* of U. For example, the compositional inverse of $x - x^2$ is $C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + \cdots$, whose coefficients are the Catalan numbers. Indeed $C - C^2 = x$, which says exactly that substituting C into $x - x^2$ gives x.

Proof. Write $V = x + v_2x^2 + v_3x^3 + \cdots$ with unknown coefficients. In $U(V) = V + u_2V^2 + u_3V^3 + \cdots$, the coefficient of x^n (for $n \ge 2$) equals v_n plus terms that involve only $v_2, \ldots, v_{n-1}$.

Setting it equal to 0 therefore determines v_n from the earlier coefficients, using only additions, subtractions and multiplications. So there is exactly one V with $U \circ V = x$, and it has integer coefficients if U does. Applying the same construction to V gives a normalized W with $V \circ W = x$. Then $U = U \circ (V \circ W) = (U \circ V) \circ W = W$, so $V \circ U = V \circ W = x$ as well. $\square$

2.4 Remainders and congruences

For an integer $m \geq 2$ we write $a \equiv b \pmod{m}$ ("a is congruent to b modulo m") when $a - b$ is divisible by m, that is, when a and b leave the same remainder after division by m. The remainder of a on division by m is called its *residue* modulo m; it is one of $0, 1, \ldots, m-1$.

Addition and multiplication are compatible with congruences, so the residues $0, 1, \ldots, m-1$, added and multiplied modulo m, form a ring, written $\mathbb{Z}/m\mathbb{Z}$. When $m = p$ is a prime we also write $\mathbb{F}_p$. In $\mathbb{F}_2 = \{0, 1\}$ we have $1 + 1 = 0$, so addition and subtraction coincide.

For power series with integer coefficients, $U \equiv V \pmod{m}$ means that *every* coefficient of $U - V$ is divisible by m. Equivalently, the two series become equal when every coefficient is replaced by its residue; we call this *reducing* the series modulo m. The key fact is the following.

Reduction modulo m is compatible with all the operations above. To reduce a sum, a product, a substitution, or an inverse, we may reduce the ingredients first and then compute in $\mathbb{Z}/m\mathbb{Z}$. This holds because every coefficient of the result is obtained from finitely many coefficients of the ingredients by adding, subtracting and multiplying, and these operations are compatible with remainders.

Rings. Some statements below hold over every commutative ring, for example $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{Z}/m\mathbb{Z}$. No division is used in them; this matters because in $\mathbb{Z}/m\mathbb{Z}$ division is not always possible (for instance, 2 has no inverse in $\mathbb{Z}/4\mathbb{Z}$).

2.5 The method

The main proofs follow three steps.

1. *Uniqueness.* Show that the defining equation determines the coefficients one at a time, using only addition, subtraction and multiplication. Then the equation has exactly one solution of the required shape, in every commutative ring, including $\mathbb{Z}/8\mathbb{Z}$.
2. *Reduction.* Reducing the integer solution modulo 8 gives a solution of the reduced equation, because reduction is compatible with all operations.
3. *A candidate solution.* Find an explicit series that satisfies the reduced equation. By uniqueness, it is the reduction of the integer solution, and its coefficients can be read off.

Finding the candidate in step 3 is usually the main difficulty.

2.6 The Frobenius identity

Lemma 2.2 (Frobenius identity)**.** *Let p be a prime. For every power series U with coefficients in $\mathbb{F}_p$, we have $U(x)^p = U(x^p)$.*

Proof. In $\mathbb{F}_p$, the binomial theorem simplifies to $(a+b)^p = a^p + b^p$, because every binomial coefficient $\binom{p}{i}$ with $0 < i < p$ is divisible by p. Also $c^p = c$ for every $c \in \mathbb{F}_p$ (Fermat's little theorem). So $\left(\sum_n u_n x^n\right)^p = \sum_n u_n^p x^{pn} = \sum_n u_n x^{pn} = U(x^p)$. The first equality holds for

polynomials by repeated use of $(a+b)^p = a^p + b^p$; it holds for power series because each coefficient of $U(x)^p$ involves only finitely many coefficients of U. □

For $p = 2$ this says that squaring a series modulo 2 doubles every exponent: for instance, $(x + x^3)^2 \equiv x^2 + x^6 \pmod 2$. In particular, the square of any series has, modulo 2, no terms of odd degree.

2.7 The Catalan numbers modulo 2

Lemma 2.3. *In every commutative ring the equation $U = x + U^2$ has exactly one solution with $U(0) = 0$. Over the integers it is the Catalan series $C(x) = x + x^2 + 2x^3 + 5x^4 + 14x^5 + \cdots$, and*

$$C(x) \equiv x + x^2 + x^4 + x^8 + \cdots = \sum_{k\ge 0} x^{2^k} \pmod 2.$$

So $[x^n]\,C$ is odd exactly when n is a power of 2.

Proof. The coefficient of x^n in U^2 involves only $u_1, \ldots, u_{n-1}$, so the equation determines u_n one at a time: $u_1 = 1$ and $u_n = \sum_{i=1}^{n-1} u_i u_{n-i}$. Over $\mathbb{F}_2$, the series $S = \sum_k x^{2^k}$ satisfies $x + S^2 = x + \sum_k x^{2^{k+1}} = S$ by Lemma 2.2, so by uniqueness $C \equiv S \pmod 2$. □

The parity of the Catalan numbers is classical; see for example [3]. (Here the Catalan number $\frac{1}{n}\binom{2n-2}{n-1}$ appears as the coefficient of x^n, so "n a power of 2" corresponds to the familiar statement that the k-th Catalan number is odd exactly when $k+1$ is a power of 2.)

3 Results

Throughout, C is the Catalan series, the unique normalized series with $C = x + C^2$ (Lemma 2.3), and $c_n = [x^n]\,C$. We also need the series of the companion entry A393172: A denotes the unique normalized series with integer coefficients that satisfies

$$A(x - x^2) = x + A(x)^2 \tag{1}$$

(Proposition 4.2 below shows that there is exactly one). The series of A393171 is defined by

$$B(x) = x + \big(B(C(x))\big)^2. \tag{2}$$

Theorem 3.1. *(a) Equation (2) has exactly one normalized solution B with integer coefficients (and, more generally, exactly one with coefficients in any commutative ring). The sequence A393171 is $a(n) = [x^n]\,B$.*

(b) The series A and B are related by

$$B = x + A^2, \qquad B(C(x)) = A(x), \qquad B(x - x^2) = x - x^2 + B(x)^2.$$

(c) Modulo 8,

$$B(x) \equiv x + x^2 + 4\big(C(x) - x - x\,C(x)\big) \pmod 8.$$

Corollary 3.2 (Hanna's conjecture)**.** *For every $n > 2$: if $n = 2^k$ or $n = 2^k + 1$ for some $k \ge 1$, then $a(n) \equiv 4 \pmod 8$; otherwise $a(n)$ is divisible by 8.*

Proof. Let $n > 2$. The coefficient of x^n in $x + x^2$ is 0, and in $4(C - x - xC)$ it is $4(c_n - c_{n-1})$, because the coefficient of x^n in $x\,C(x)$ is c_{n-1}. So Theorem 3.1(c) gives $a(n) \equiv 4(c_n - c_{n-1}) \pmod 8$. This is 4 modulo 8 if $c_n - c_{n-1}$ is odd, and 0 if it is even. Now $c_n - c_{n-1}$ is odd exactly when one of c_n, c_{n-1} is odd and the other is even. By Lemma 2.3, c_j is odd exactly when j is a power of 2. Since $n > 2$, the numbers n and $n - 1$ are not both powers of 2 (the only consecutive powers of 2 are 1 and 2). So $a(n) \equiv 4 \pmod 8$ exactly when n or $n - 1$ is a power of 2, that is, when $n = 2^k$ or $n = 2^k + 1$; and since $n > 2$, in both cases $k \geq 1$. □

For example, $a(5) = 116 = 8 \cdot 14 + 4$, and indeed $c_5 - c_4 = 14 - 5 = 9$ is odd; while $a(6) = 752 = 8 \cdot 94$, and $c_6 - c_5 = 42 - 14 = 28$ is even.

4 Proofs

The idea in brief. First we prove the result for the companion series A: modulo 4 it equals $2C - x$. Written over the integers, this says $A = 2C - x + 4K$ for some series K with integer coefficients. Next we show that $B = x + A^2$. Squaring $2C - x + 4K$ gives $(2C - x)^2$ plus multiples of 8, so modulo 8 the unknown series K disappears, and B is determined. The identity $C^2 = C - x$ simplifies the result.

Lemma 4.1 (An equation with a square)**.** *Let T and P be normalized series with coefficients in a commutative ring R. There is exactly one normalized series G with coefficients in R such that*

$$G(x) = T(x) + \big(G(P(x))\big)^2.$$

If T and P have integer coefficients, then so does G, and reducing G modulo m gives the solution of the same equation over $\mathbb{Z}/m\mathbb{Z}$ (with T and P reduced modulo m).

Proof. The point is that the coefficient of x^n on the right-hand side depends only on the coefficients of G below degree n. To see this, let G and H be two series with constant term 0 that agree below degree r, so $G - H = O(x^r)$. Since P starts with x, the power P^k has no terms below x^k, and so $G(P) - H(P) = O(x^r)$ as well. Moreover $G(P) + H(P)$ has constant term 0. Hence

$$G(P)^2 - H(P)^2 = \big(G(P) - H(P)\big)\big(G(P) + H(P)\big) = O(x^{r+1}).$$

In other words, changing G in degree r and above changes the right-hand side only in degree $r + 1$ and above.

So the equation fixes the coefficients of G one at a time. The constant term is 0, the coefficient of x is the coefficient of x in T, which is 1, and for $n \geq 2$ the coefficient of x^n in G must equal the coefficient of x^n in $T + G(P)^2$, which has already been computed from the earlier coefficients. Only additions and multiplications are used, so this works in every commutative ring, and gives integer coefficients when T and P have them. Finally, reducing the integer solution modulo m gives a normalized series that satisfies the reduced equation (reduction is compatible with all the operations involved), and by uniqueness it is *the* solution over $\mathbb{Z}/m\mathbb{Z}$. □

Proposition 4.2 (The companion sequence A393172)**.** *Write $N(x) = x - x^2$. Equation* (1) *has exactly one normalized solution A with integer coefficients. It also satisfies $A = C + A(C)^2$, and $A \equiv 2C - x \pmod 4$.*

Proof. Since $N(C) = C - C^2 = x$, the series C is the compositional inverse of N, and also $C(N(x)) = x$ (Lemma 2.1). Substitution can be done in steps, and substituting into a sum or product is the same as adding or multiplying the substituted terms. So, substituting C for x in $A(N) = x + A^2$ gives $A = C + A(C)^2$, and substituting N for x in $A = C + A(C)^2$ gives back $A(N) = x + A^2$. The two equations therefore have the same normalized solutions, and by Lemma 4.1 (with $T = P = C$) the second has exactly one, in every commutative ring.

Over $\mathbb{Z}/4\mathbb{Z}$ the solution is the reduction of the integer A (Lemma 4.1), so it suffices to check that $D = 2C - x$ satisfies (1) in $\mathbb{Z}/4\mathbb{Z}$. Indeed $D(N) = 2C(N) - N = 2x - (x - x^2) = x + x^2$, and $x + D^2 = x + 4C^2 - 4xC + x^2 = x + x^2$ in $\mathbb{Z}/4\mathbb{Z}$. Hence $A \equiv 2C - x \pmod 4$. □

Proof of Theorem 3.1. (a) This is Lemma 4.1 with $T = x$ and $P = C$.

(b) Put $B' = x + A^2$. Substituting C for x and using $A = C + A(C)^2$ (Proposition 4.2),

$$B'(C(x)) = C + A(C)^2 = A.$$

Hence $B' = x + A^2 = x + B'(C)^2$, so B' is a normalized solution of (2). By the uniqueness in part (a), $B = B'$. This proves $B = x + A^2$ and $B(C) = A$. Finally, (1) says that $A(N) = x + A^2 = B$; substituting N for x in $B = x + A^2$ therefore gives $B(N) = N + A(N)^2 = N + B^2$, which is the third relation.

(c) By Proposition 4.2, every coefficient of $A - (2C - x)$ is divisible by 4. So $A = 2C - x + 4K$, where K is a series with *integer* coefficients. Squaring, over the integers,

$$A^2 = (2C - x)^2 + 8K\,(2C - x) + 16K^2.$$

The last two terms are divisible by 8. Using $C^2 = C - x$,

$$(2C - x)^2 = 4C^2 - 4xC + x^2 = x^2 + 4\big(C - x - xC\big).$$

So $A^2 \equiv x^2 + 4(C - x - xC) \pmod 8$, and by part (b), $B = x + A^2 \equiv x + x^2 + 4(C - x - xC) \pmod 8$. □

The proof does not divide by 2 or 4 in arithmetic modulo 8, where such divisions are not defined. The information modulo 4 was first turned into an exact statement over the integers (the series K), and only then reduced modulo 8.

5 Checking the results

A mathematical proof can contain mistakes, so the results were checked in two independent ways.

Numerical check. We computed B by repeating the substitution $B \mapsto x + B(C)^2$ with exact integer arithmetic, starting from $B = x$; each round fixes at least one more coefficient. This reproduces all 24 terms printed in the OEIS entry. We also computed the series A of A393172 separately, from its own equation (1), and checked that $B = x + A^2$ for all coefficients below degree 200, and that the rule of Corollary 3.2 holds for every n with $2 < n < 200$. The computation was done with the PARI/GP computer algebra system, using code that was written separately from the formal proofs (file `verification/independent_checks.gp` in the accompanying repository [9]). A computation can only test finitely many cases, so it is a check on the definitions and on the indexing, not a proof.

Formal proof in Lean. Lean [7] is a *proof assistant*: a program in which mathematical statements and their proofs are written in a precise formal language, and every step is checked mechanically down to the basic rules of logic. The theorems of Section 3 are proved in Lean 4 (version 4.19.0) on top of Mathlib [8], Lean's library of formalized mathematics. The statements are collected in the file `lean/Entries/A393171.lean` of the repository, in the namespace `OEIS.A393171`. There the Catalan series is called `catalan`, and B is defined as $x + A^2$; the theorems show that this series satisfies the equation of the entry and is its only normalized solution. The main proofs are in the module `lean/Families/CatalanComposition.lean`. The table lists where each result of this paper is proved.

Statement in this paper	Lean theorem(s)
Theorem 3.1(a) for integer coefficients, and the same for the equation $B(x - x^2) = x - x^2 + B^2$; B is normalized and satisfies both	`second_named_existence`, `second_existence`, `second_definition`
Theorem 3.1(b): $B = x + A^2$ and $B(C) = A$	`link_to_first_sequence`, `link_composed_catalan`
Theorem 3.1(c): the formula modulo 8	`complete_series_mod_eight`
Corollary 3.2, also for any normalized integer solution of $B(x - x^2) = x - x^2 + B^2$	`second_residue`, `divisibility_complement`, `second_arbitrary_source`

The Lean project was rebuilt from source, and the axioms on which each of these theorems depends were listed with Lean's `#print axioms` command. Every one depends only on the three standard axioms of Lean's logic (`propext`, `Classical.choice`, `Quot.sound`); no step is left unproved. A formal proof certifies exactly the statement that Lean checks. Whether that statement says what the OEIS entry means is a separate question, so the Lean definitions were also compared with the definition in the OEIS entry (Section 1.1).

6 Discussion

Scope. Theorem 3.1 determines the remainder of every coefficient $a(n)$ modulo 8, so it settles the conjecture of the entry completely. It says nothing about the size of $a(n)$, and nothing about remainders modulo 16 or higher powers of 2.

Going further. The step from A to $B = x + A^2$ always gains one power of 2: if $A = D + 2^k K$ with $k \geq 1$ and K a series with integer coefficients, then $A^2 = D^2 + 2^{k+1} DK + 2^{2k} K^2 \equiv D^2 \pmod{2^{k+1}}$. So any rule for A modulo 8 would give a rule for B modulo 16. The companion paper on A393172 records such a rule for A modulo 8, found by computer and checked for all $n < 200$, but not proved. Proving it would therefore also describe B modulo 16.

About this paper

How the work was done. This paper is one of 29 papers on conjectures from the OEIS that came out of an AI research effort directed by Dev D. Goyal [9]. The roles were as follows.

- *Dev D. Goyal* started the project, chose which problems to pursue, asked for proofs, formal verification and revisions, and decided what to publish. Dev D. Goyal did not supply proof strategies or intermediate arguments.

- *ChatGPT* (OpenAI), using the GPT-6 model with Pro reasoning, worked autonomously from those prompts. It searched the OEIS for suitable conjectures, found the proofs, wrote the Lean 4 formalization, and wrote a first draft of this paper.
- *Claude Opus 5.5*, used with Claude Code, checked the work independently. It rebuilt the Lean project from source and confirmed that the theorems listed in Section 5 rely only on Lean's standard axioms; compared the Lean definitions with the OEIS definition; recomputed the sequence with separately written PARI/GP code, compared it with the terms published in the OEIS, and tested every claim of the paper on many terms; and read every step of the proofs. It then wrote the present version of the paper, expanding the explanations and, in places, writing out an argument in more detail or by a more elementary route. The theorems of Section 3 and the proof strategies are ChatGPT's. The Discussion section, including the short arguments in it, was written by Claude Opus 5.5; its observations were checked by computer and are not part of the Lean formalization.

Dev D. Goyal has verified the results and is responsible for this paper.

Review status. This paper has not been peer reviewed, and it has not been submitted to a journal or to the OEIS.

Priority. An OEIS entry that labels a statement as a conjecture shows only that its contributor did not know a proof; it does not show that no proof existed. During the research only limited searches for earlier work were made. No claim of priority is made.

Corrections. Please report any error in the issue tracker of the repository [9], where the Lean sources and the numerical checks are also available.

Acknowledgements

The sequence, its formulas and the conjecture are due to Paul D. Hanna [1], as is the companion entry A393172 [2].

References

[1] P. D. Hanna, entry A393171 (G.f. satisfies $A(x) = x + A(C(x))^2$, where $C(x) = x + C(x)^2$), *The On-Line Encyclopedia of Integer Sequences*, February 2026. `https://oeis.org/A393171`. Text as of October 2026.

[2] P. D. Hanna, entry A393172 (G.f. $A(x)$ satisfies $A(x - x^2) = x + A(x)^2$), *The On-Line Encyclopedia of Integer Sequences*, February 2026. `https://oeis.org/A393172`. Text as of October 2026.

[3] E. Deutsch and B. E. Sagan, Congruences for Catalan and Motzkin numbers and related sequences, *Journal of Number Theory* 117 (2006), 191–215.

[4] N. J. A. Sloane and The OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*. `https://oeis.org`.

[5] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999. (Chapter 5 treats formal power series, composition and compositional inverses.)

[6] I. Niven, Formal power series, *American Mathematical Monthly* 76 (1969), 871–889.

[7] L. de Moura and S. Ullrich, The Lean 4 theorem prover and programming language, in *Automated Deduction – CADE 28*, Lecture Notes in Computer Science 12699, Springer, 2021, 625–635.

[8] The mathlib Community, The Lean mathematical library, in *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, 367–381. Version used: commit `c44e0c8ee63c`.

[9] D. D. Goyal, *Proofs and corrections of OEIS conjectures*, repository, 2026. `https://github.com/dvgyl/oeis-a396843-proof`.