

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	16385	Lag set, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided set, dyadic depth 20	0/373/27	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided intersection, dyadic depth 20	0/361/39	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lag set, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided set, dyadic depth 28	0/373/27	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	16385	Two-sided intersection, dyadic depth 28	0/361/39	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lag set, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided set, dyadic depth 36	0/373/27	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided intersection, dyadic depth 36	0/361/39	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lag set, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	16385	Two-sided set, arithmetic bits 160	0/373/27	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 160	0/361/39	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lag set, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided set, arithmetic bits 320	0/373/27	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 320	0/361/39	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lag set, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided set, arithmetic bits 80	0/373/27	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 80	0/361/39	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lag set, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided set, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided intersection, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lag set, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided set, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided intersection, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lag set, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided set, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided intersection, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lag set, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided set, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lag set, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided set, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, reversible, gaussian	65537	Lag set, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided set, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lag set, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided set, dyadic depth 20	153/29/18	[0.700, 0.822]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 20	151/23/26	[0.689, 0.813]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lag set, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided set, dyadic depth 28	153/29/18	[0.700, 0.822]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 28	152/22/26	[0.694, 0.818]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lag set, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	16385	Two-sided set, dyadic depth 36	153/29/18	[0.700, 0.822]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 36	152/22/26	[0.694, 0.818]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lag set, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided set, arithmetic bits 160	153/29/18	[0.700, 0.822]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 160	152/22/26	[0.694, 0.818]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lag set, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided set, arithmetic bits 320	153/29/18	[0.700, 0.822]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 320	152/22/26	[0.694, 0.818]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lag set, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided set, arithmetic bits 80	153/29/18	[0.700, 0.822]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 80	152/22/26	[0.694, 0.818]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lag set, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided set, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lag set, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided set, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lag set, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided set, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lag set, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided set, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	65537	Lag set, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided set, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lag set, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, strong delay, gaussian	65537	Two-sided set, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lag set, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided set, dyadic depth 20	48/132/20	[0.182, 0.306]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 20	45/130/25	[0.169, 0.290]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lag set, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, weak delay, gaussian	16385	Two-sided set, dyadic depth 28	48/132/20	[0.182, 0.306]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 28	45/130/25	[0.169, 0.290]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lag set, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided set, dyadic depth 36	48/132/20	[0.182, 0.306]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 36	45/130/25	[0.169, 0.290]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lag set, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided set, arithmetic bits 160	48/132/20	[0.182, 0.306]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 160	45/130/25	[0.169, 0.290]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lag set, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided set, arithmetic bits 320	48/132/20	[0.182, 0.306]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 320	45/130/25	[0.169, 0.290]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lag set, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided set, arithmetic bits 80	48/132/20	[0.182, 0.306]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 80	45/130/25	[0.169, 0.290]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lag set, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided set, dyadic depth 20	197/3/0	[0.956, 0.997]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 20	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lag set, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided set, dyadic depth 28	197/3/0	[0.956, 0.997]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 28	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lag set, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided set, dyadic depth 36	197/3/0	[0.956, 0.997]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 36	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, weak delay, gaussian	65537	Lag set, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided set, arithmetic bits 160	197/3/0	[0.956, 0.997]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 160	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lag set, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided set, arithmetic bits 320	197/3/0	[0.956, 0.997]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 199/200, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 320	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lag set, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided set, arithmetic bits 80	197/3/0	[0.956, 0.997]	Source reversibility under stated assumptions
phi 199/200, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 80	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lag set, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/136/264	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/19/381	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	16385	Two-sided set, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided intersection, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lag set, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/136/264	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/19/381	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided set, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided intersection, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lag set, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/136/264	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/19/381	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided set, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided intersection, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lag set, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/136/264	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/19/381	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided set, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lag set, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/136/264	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/19/381	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided set, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lag set, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/136/264	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/19/381	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided set, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lag set, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/395/5	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided set, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided intersection, dyadic depth 20	0/399/1	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lag set, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/395/5	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided set, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided intersection, dyadic depth 28	0/399/1	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	65537	Lag set, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/395/5	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided set, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided intersection, dyadic depth 36	0/399/1	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lag set, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/395/5	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided set, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 160	0/399/1	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lag set, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/395/5	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided set, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 320	0/399/1	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lag set, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/395/5	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, reversible, gaussian	65537	Two-sided set, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 80	0/399/1	[0.000, 0.010]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lag set, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/69/131	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided set, dyadic depth 20	115/85/0	[0.503, 0.645]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lag set, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/69/131	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided set, dyadic depth 28	115/85/0	[0.503, 0.645]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lag set, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/69/131	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided set, dyadic depth 36	115/85/0	[0.503, 0.645]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lag set, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/69/131	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided set, arithmetic bits 160	115/85/0	[0.503, 0.645]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lag set, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/69/131	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided set, arithmetic bits 320	115/85/0	[0.503, 0.645]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lag set, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/69/131	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided set, arithmetic bits 80	115/85/0	[0.503, 0.645]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lag set, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	63/137/0	[0.251, 0.385]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	10/187/3	[0.024, 0.091]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided set, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, strong delay, gaussian	65537	Lag set, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	63/137/0	[0.251, 0.385]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	10/187/3	[0.024, 0.091]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided set, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lag set, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	63/137/0	[0.251, 0.385]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	10/187/3	[0.024, 0.091]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided set, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lag set, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	63/137/0	[0.251, 0.385]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	10/187/3	[0.024, 0.091]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided set, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lag set, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	63/137/0	[0.251, 0.385]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	10/187/3	[0.024, 0.091]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, strong delay, gaussian	65537	Two-sided set, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lag set, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	63/137/0	[0.251, 0.385]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	10/187/3	[0.024, 0.091]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided set, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lag set, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/83/117	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided set, dyadic depth 20	38/162/0	[0.138, 0.252]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lag set, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/83/117	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided set, dyadic depth 28	38/162/0	[0.138, 0.252]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lag set, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/83/117	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided set, dyadic depth 36	38/162/0	[0.138, 0.252]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lag set, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/83/117	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided set, arithmetic bits 160	38/162/0	[0.138, 0.252]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lag set, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/83/117	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided set, arithmetic bits 320	38/162/0	[0.138, 0.252]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lag set, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/83/117	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/12/188	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided set, arithmetic bits 80	38/162/0	[0.138, 0.252]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	65537	Lag set, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/200/0	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/198/2	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided set, dyadic depth 20	123/77/0	[0.543, 0.683]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lag set, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/200/0	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/198/2	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided set, dyadic depth 28	123/77/0	[0.543, 0.683]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lag set, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/200/0	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/198/2	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided set, dyadic depth 36	123/77/0	[0.543, 0.683]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lag set, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/200/0	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/198/2	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	65537	Two-sided set, arithmetic bits 160	123/77/0	[0.543, 0.683]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lag set, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/200/0	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/198/2	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided set, arithmetic bits 320	123/77/0	[0.543, 0.683]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lag set, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/200/0	[0.000, 0.019]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/198/2	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 4/5, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided set, arithmetic bits 80	123/77/0	[0.543, 0.683]	Source reversibility under stated assumptions
phi 4/5, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lag set, dyadic depth 20	0/8/392	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 20	0/6/394	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided set, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided intersection, dyadic depth 20	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lag set, dyadic depth 28	0/8/392	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 28	0/6/394	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided set, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided intersection, dyadic depth 28	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lag set, dyadic depth 36	0/8/392	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail intersection, dyadic depth 36	0/6/394	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided set, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided intersection, dyadic depth 36	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lag set, arithmetic bits 160	0/8/392	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 160	0/6/394	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided set, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 160	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lag set, arithmetic bits 320	0/8/392	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 320	0/6/394	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided set, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 320	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, reversible, gaussian	16385	Lag set, arithmetic bits 80	0/8/392	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/0/400	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Lower-tail intersection, arithmetic bits 80	0/6/394	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided set, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	16385	Two-sided intersection, arithmetic bits 80	0/400/0	[0.000, 0.010]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lag set, dyadic depth 20	0/525/475	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/19/981	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/0/1000	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 20	0/891/109	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided set, dyadic depth 20	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, reversible, gaussian	65537	Two-sided intersection, dyadic depth 20	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lag set, dyadic depth 28	0/525/475	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/19/981	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/0/1000	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 28	0/891/109	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided set, dyadic depth 28	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided intersection, dyadic depth 28	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lag set, dyadic depth 36	0/525/475	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/19/981	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/0/1000	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail intersection, dyadic depth 36	0/891/109	[0.000, 0.004]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, reversible, gaussian	65537	Two-sided set, dyadic depth 36	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided intersection, dyadic depth 36	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lag set, arithmetic bits 160	0/525/475	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/19/981	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/0/1000	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 160	0/891/109	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided set, arithmetic bits 160	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 160	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lag set, arithmetic bits 320	0/525/475	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/19/981	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/0/1000	[0.000, 0.004]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 320	0/891/109	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided set, arithmetic bits 320	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 320	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lag set, arithmetic bits 80	0/525/475	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/19/981	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/0/1000	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Lower-tail intersection, arithmetic bits 80	0/891/109	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided set, arithmetic bits 80	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, reversible, gaussian	65537	Two-sided intersection, arithmetic bits 80	0/1000/0	[0.000, 0.004]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lag set, dyadic depth 20	9/0/191	[0.020, 0.084]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 20	5/1/194	[0.008, 0.058]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided set, dyadic depth 20	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 20	191/9/0	[0.916, 0.980]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lag set, dyadic depth 28	9/0/191	[0.020, 0.084]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 28	5/1/194	[0.008, 0.058]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided set, dyadic depth 28	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 28	191/9/0	[0.916, 0.980]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lag set, dyadic depth 36	9/0/191	[0.020, 0.084]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail intersection, dyadic depth 36	5/1/194	[0.008, 0.058]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided set, dyadic depth 36	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided intersection, dyadic depth 36	191/9/0	[0.916, 0.980]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lag set, arithmetic bits 160	9/0/191	[0.020, 0.084]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 160	5/1/194	[0.008, 0.058]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided set, arithmetic bits 160	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 160	191/9/0	[0.916, 0.980]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	16385	Lag set, arithmetic bits 320	9/0/191	[0.020, 0.084]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 320	5/1/194	[0.008, 0.058]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided set, arithmetic bits 320	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 320	191/9/0	[0.916, 0.980]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lag set, arithmetic bits 80	9/0/191	[0.020, 0.084]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Lower-tail intersection, arithmetic bits 80	5/1/194	[0.008, 0.058]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	16385	Two-sided set, arithmetic bits 80	194/6/0	[0.935, 0.989]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	16385	Two-sided intersection, arithmetic bits 80	191/9/0	[0.916, 0.980]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lag set, dyadic depth 20	103/0/97	[0.443, 0.587]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/2/198	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 20	168/5/27	[0.781, 0.888]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided set, dyadic depth 20	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lag set, dyadic depth 28	103/0/97	[0.443, 0.587]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/2/198	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 28	168/5/27	[0.781, 0.888]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	65537	Two-sided set, dyadic depth 28	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lag set, dyadic depth 36	103/0/97	[0.443, 0.587]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/2/198	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail intersection, dyadic depth 36	168/5/27	[0.781, 0.888]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided set, dyadic depth 36	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lag set, arithmetic bits 160	103/0/97	[0.443, 0.587]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/2/198	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 160	168/5/27	[0.781, 0.888]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided set, arithmetic bits 160	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lag set, arithmetic bits 320	103/0/97	[0.443, 0.587]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/2/198	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 320	168/5/27	[0.781, 0.888]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided set, arithmetic bits 320	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lag set, arithmetic bits 80	103/0/97	[0.443, 0.587]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/2/198	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, strong delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Lower-tail intersection, arithmetic bits 80	168/5/27	[0.781, 0.888]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided set, arithmetic bits 80	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, strong delay, gaussian	65537	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lag set, dyadic depth 20	4/0/196	[0.005, 0.051]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 20	1/1/198	[0.000, 0.028]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided set, dyadic depth 20	92/108/0	[0.389, 0.532]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 20	147/53/0	[0.668, 0.795]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lag set, dyadic depth 28	4/0/196	[0.005, 0.051]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 28	1/1/198	[0.000, 0.028]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided set, dyadic depth 28	92/108/0	[0.389, 0.532]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 28	147/53/0	[0.668, 0.795]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lag set, dyadic depth 36	4/0/196	[0.005, 0.051]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail finite cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail intersection, dyadic depth 36	1/1/198	[0.000, 0.028]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided set, dyadic depth 36	92/108/0	[0.389, 0.532]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided intersection, dyadic depth 36	147/53/0	[0.668, 0.795]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, weak delay, gaussian	16385	Lag set, arithmetic bits 160	4/0/196	[0.005, 0.051]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 160	1/1/198	[0.000, 0.028]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided set, arithmetic bits 160	92/108/0	[0.389, 0.532]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 160	147/53/0	[0.668, 0.795]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lag set, arithmetic bits 320	4/0/196	[0.005, 0.051]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 320	1/1/198	[0.000, 0.028]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided set, arithmetic bits 320	92/108/0	[0.389, 0.532]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 320	147/53/0	[0.668, 0.795]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lag set, arithmetic bits 80	4/0/196	[0.005, 0.051]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail finite cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Lower-tail intersection, arithmetic bits 80	1/1/198	[0.000, 0.028]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided set, arithmetic bits 80	92/108/0	[0.389, 0.532]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	16385	Two-sided intersection, arithmetic bits 80	147/53/0	[0.668, 0.795]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lag set, dyadic depth 20	102/5/93	[0.438, 0.582]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 20	0/4/196	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 20	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 20	86/95/19	[0.360, 0.502]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, weak delay, gaussian	65537	Two-sided set, dyadic depth 20	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 20	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lag set, dyadic depth 28	102/5/93	[0.438, 0.582]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 28	0/4/196	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 28	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 28	86/95/19	[0.360, 0.502]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided set, dyadic depth 28	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 28	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lag set, dyadic depth 36	102/5/93	[0.438, 0.582]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail finite cutoff, dyadic depth 36	0/4/196	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail analytic cutoff, dyadic depth 36	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, weak delay, gaussian	65537	Lower-tail intersection, dyadic depth 36	86/95/19	[0.360, 0.502]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided set, dyadic depth 36	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided intersection, dyadic depth 36	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lag set, arithmetic bits 160	102/5/93	[0.438, 0.582]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 160	0/4/196	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 160	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 160	86/95/19	[0.360, 0.502]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided set, arithmetic bits 160	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 160	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lag set, arithmetic bits 320	102/5/93	[0.438, 0.582]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 320	0/4/196	[0.000, 0.019]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.

Numerical

Condition	N	Method	R / NR / A	95% rejection interval	Nominal test target
phi 97/100, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 320	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 320	86/95/19	[0.360, 0.502]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided set, arithmetic bits 320	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 320	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lag set, arithmetic bits 80	102/5/93	[0.438, 0.582]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail finite cutoff, arithmetic bits 80	0/4/196	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail analytic cutoff, arithmetic bits 80	0/0/200	[0.000, 0.019]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Lower-tail intersection, arithmetic bits 80	86/95/19	[0.360, 0.502]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided set, arithmetic bits 80	199/1/0	[0.972, 1.000]	Source reversibility under stated assumptions
phi 97/100, weak delay, gaussian	65537	Two-sided intersection, arithmetic bits 80	200/0/0	[0.981, 1.000]	Source reversibility under stated assumptions

R, rejection; NR, nonrejection; A, abstention. Intervals use all records. Displayed endpoints round outward to 0.001. The nominal target does not give a size guarantee. Violated source, covariance, response, or error assumptions remove that guarantee. The baseline comparators have no finite-record source-null guarantee here.