

Span Categories & their Uses IV:

Universal Biadjointable Functors

David Martínez-Carpén

2nd of February of 2026

A **3-functor formalism** is a lax symmetric monoidal $(\infty, 2)$ -functor
 $D: \text{Span}(C, E) \rightarrow \text{Cat}_\infty$

For any $f: X \rightarrow Y$ in C , define $f^* := D(Y \xleftarrow{f} X = X): D(Y) \rightarrow D(X)$

For any $g: X \rightarrow Y$ in E , define $g_! := D(X = X \xrightarrow{g} Y): D(X) \rightarrow D(Y)$

A **6-functor formalism** is a 3-functor formalism for which the
functors $- \otimes A$, f^* and $g_!$ admit right adjoints

6-functor formalism = operations $(f^*, f_*, g_!, g^!, \otimes, \underline{\text{Hom}})$
+ relationships

A **3-functor formalism** is a lax symmetric monoidal $(\infty, 2)$ -functor
 $D: \text{Span}(\mathcal{C}, \mathcal{E}) \rightarrow \text{Cat}_\infty$

For any $f: X \rightarrow Y$ in $\mathcal{C}$, define $f^* := D(Y \xleftarrow{f} X = X): D(Y) \rightarrow D(X)$

For any $g: X \rightarrow Y$ in $\mathcal{E}$, define $g_! := D(X = X \xrightarrow{g} Y): D(X) \rightarrow D(Y)$

A **6-functor formalism** is a 3-functor formalism for which the
functors $- \otimes A$, f^* and $g_!$ admit right adjoints

⤴ Difficult to construct! Solution: Construct the pullback functor

$(-)^*: \mathcal{C}^{\text{op}} \rightarrow \text{Cat}_\infty$ and extend it to $\text{Span}(\mathcal{C}, \mathcal{E})$

One extension: Assume $\mathcal{E}$ decomposes in two classes $\mathcal{I}$ and $\mathcal{P}$

For $i \in \mathcal{I}$, i^* has left adjoint $i_\#$
For $p \in \mathcal{P}$, p^* has right adjoint p_* } $\Rightarrow (-)_! := p_* i_\# : \mathcal{C} \rightarrow \text{Cat}_\infty$

Generalisation: $\mathbb{D}$ $(\infty, 2)$ -category instead of $\mathbf{Cat}_\infty$

$\text{SPAN}_2(C, E)_{P, I} := (\infty, 2)$ -category of spans, with spans of spans
as 2-morphisms

Theorem A. C ∞ -cat, $I, P \subset E \subset C$ wide subcats s.t.

- (i) I and P are closed under base change and left cancellable
- (ii) Every morphism in E factors as $p \circ i$ with $i \in I$ and $p \in P$
- (iii) Every map in $I \cap P$ is truncated

Then the inclusion $h: C^{\text{op}} \hookrightarrow \text{SPAN}_2(C, E)_{P, I}$ induces an equivalence

$$h^*: \text{FUN}(\text{SPAN}_2(C, E)_{P, I}, \mathbb{D}) \xrightarrow{\sim} \text{FUN}_{(I, P)\text{-bidi}}(C^{\text{op}}, \mathbb{D})$$

$I = P = E$ and $(2, 2)$ -categories
Balmer and Dell'Ambrogio

$\uparrow$
 (I, P) -bidiointable functors and nat. trans.

Intuition. Let $m \leq n \leq \infty$, an (n, m) -category is a higher category having i -morphisms for $i \leq n$ which are invertible $i > m$.

Notation. $\text{categories} := (\infty, 1)$ -categories FUN, HOM categories in
 n -categories $:= (\infty, n)$ -categories a 2-categorical setting
 CAT 2-category of categories, CAT_2 3-category of 2-categories

Intuition. Let $m \leq n \leq \infty$, an (n, m) -category is a higher category having i -morphisms for $i \leq n$ which are invertible $i > m$.

Notation. categories $:= (\infty, 1)$ -categories FUN, HOM categories in a 2-categorical setting
 n -categories $:= (\infty, n)$ -categories
CAT 2-category of categories, CAT₂ 3-category of 2-categories

Goal: Given a category C and suitable subcategories I and P :

- Introduce (I, P) -biadjointable functor
- The universal biadjointable functor
- Recognition principle $\rightsquigarrow$ Used to prove Theorem A in future talks

$\mathbb{D}$ 2-category, a commutative square $\begin{array}{ccc} X' & \xrightarrow{g^*} & X \\ \downarrow f^* & & \downarrow k^* \\ Y' & \xrightarrow{h^*} & Y \end{array}$ in $\mathbb{D}$ is:

① **Vertically left adjointable** if f^* and k^* admit left adjoints $f_\#$ and $k_\#$ and the Beck-Chevalley map

$$BC_\# : k_\# h^* \rightarrow k_\# h^* f^* f_\# \simeq k_\# k^* g^* f_\# \rightarrow g^* f_\# \quad \text{is an equivalence}$$

$\mathbb{D}$ 2-category, a commutative square $\begin{array}{ccc} X' & \xrightarrow{g^*} & X \\ \downarrow f^* & & \downarrow k^* \\ Y' & \xrightarrow{h^*} & Y \end{array}$ in $\mathbb{D}$ is:

① **Vertically left adjointable** if f^* and k^* admit left adjoints $f_\#$ and $k_\#$ and the Beck-Chevalley map

$$BC_\# : k_\# h^* \rightarrow k_\# h^* f^* f_\# \simeq k_\# k^* g^* f_\# \rightarrow g^* f_\# \quad \text{is an equivalence}$$

② **Vertically right adjointable** " admit right adjoints f_* and h_*
 " $BC_* : g^* f_* \rightarrow k_* h^*$ "

$\mathbb{D}$ 2-category, a commutative square $\begin{array}{ccc} X' & \xrightarrow{g^*} & X \\ f^* \downarrow & & \downarrow k^* \\ Y' & \xrightarrow{h^*} & Y \end{array}$ in $\mathbb{D}$ is:

① **Vertically left adjointable** if f^* and k^* admit left adjoints $f_\#$ and $k_\#$ and the Beck-Chevalley map

$$BC_\# : k_\# h^* \rightarrow k_\# h^* f^* f_\# \simeq k_\# k^* g^* f_\# \rightarrow g^* f_\# \quad \text{is an equivalence}$$

② **Vertically right adjointable** " admit right adjoints f_* and k_*
 " $BC_* : g^* f_* \rightarrow k_* h^*$ "

③ **Horizontally left/right adjointable** same but for g^* and h^*

① 2-category, a commutative square $\begin{array}{ccc} X' & \xrightarrow{g^*} & X \\ \downarrow f^* & & \downarrow k^* \\ Y' & \xrightarrow{h^*} & Y \end{array}$ in $\mathbb{D}$ is:

① Vertically left adjointable if f^* and k^* admit left adjoints $f_\#$ and $k_\#$ and the Beck-Chevalley map

$$BC_\# : k_\# h^* \rightarrow k_\# h^* f^* f_\# \simeq k_\# k^* g^* f_\# \rightarrow g^* f_\# \text{ is an equivalence}$$

② Vertically right adjointable " admit right adjoints f_* and k_*
 " $BC_* : g^* f_* \rightarrow k_* h^*$ "

③ Horizontally left/right adjointable same but for g^* and h^*

④ Biadjointable if it is horizontal left adjointable and $\begin{array}{ccc} X' & \xleftarrow{g_\#} & X \\ \downarrow f^* & & \downarrow k^* \\ Y' & \xleftarrow{h_\#} & Y \end{array}$ is vertically right adjointable

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcategory closed under base change, $\mathbb{D}$ 2-category
 and $F: \mathcal{C}^{\text{op}} \rightarrow \mathbb{D}$ denote $f^* := F(f): F(Y) \rightarrow F(X) \quad \forall f: X \rightarrow Y$ in $\mathcal{C}$

① F is **left $\mathcal{I}$ -adjointable** if

$$\forall \text{ pullback } \begin{array}{ccc} X' & \xrightarrow{i} & X \\ g \downarrow & \lrcorner & \downarrow f \\ Y' & \xrightarrow{j} & Y \end{array}$$

$$i \in \mathcal{I} \implies \begin{array}{ccc} F(Y) & \xrightarrow{i^*} & F(Y') \\ \downarrow f^* & & \downarrow g^* \\ F(X) & \xrightarrow{j^*} & F(X') \end{array} \text{ is horizontally left adjointable}$$

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change, $\mathbb{D}$ 2-category and $F: \mathcal{C}^{\text{op}} \rightarrow \mathbb{D}$ denote $f^* := F(f): F(Y) \rightarrow F(X) \quad \forall f: X \rightarrow Y$ in $\mathcal{C}$

① F is **left $\mathcal{I}$ -adjointable** if

$$\forall \text{ pullback } \begin{array}{ccc} X' & \xrightarrow{i} & X \\ g \downarrow & \lrcorner & \downarrow f \\ Y' & \xrightarrow{j} & Y \end{array} \quad i \in \mathcal{I} \rightsquigarrow \begin{array}{ccc} F(Y) & \xrightarrow{i^*} & F(Y') \\ \downarrow f^* & & \downarrow g^* \\ F(X) & \xrightarrow{j^*} & F(X') \end{array} \quad \begin{array}{l} \text{is horizontally} \\ \text{left} \\ \text{adjointable} \end{array}$$

② Dually, F is **right $\mathcal{P}$ -adjointable** if

$$\forall \text{ pullback } \begin{array}{ccc} X' & \xrightarrow{g} & X \\ q \downarrow & \lrcorner & \downarrow p \\ Y' & \xrightarrow{f} & Y \end{array} \quad p \in \mathcal{P} \rightsquigarrow \begin{array}{ccc} F(Y) & \xrightarrow{g^*} & F(Y') \\ \downarrow p^* & & \downarrow q^* \\ F(X) & \xrightarrow{f^*} & F(X') \end{array} \quad \begin{array}{l} \text{is vertically} \\ \text{right} \\ \text{adjointable} \end{array}$$

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change, $\mathbb{D}$ 2-category and $F: \mathcal{C}^{\text{op}} \rightarrow \mathbb{D}$ denote $f^* := F(f): F(Y) \rightarrow F(X) \quad \forall f: X \rightarrow Y$ in $\mathcal{C}$

① F is **left $\mathcal{I}$ -adjointable** if

$$\forall \text{ pullback } \begin{array}{ccc} X' & \xrightarrow{i} & X \\ g \downarrow & \lrcorner & \downarrow f \\ Y' & \xrightarrow{j} & Y \end{array} \quad i \in \mathcal{I} \rightsquigarrow \begin{array}{ccc} F(Y) & \xrightarrow{i^*} & F(Y') \\ \downarrow f^* & & \downarrow g^* \\ F(X) & \xrightarrow{j^*} & F(X') \end{array} \quad \begin{array}{l} \text{is horizontally} \\ \text{left} \\ \text{adjointable} \end{array}$$

② Dually, F is **right $\mathcal{P}$ -adjointable** if

$$\forall \text{ pullback } \begin{array}{ccc} X' & \xrightarrow{g} & X \\ q \downarrow & \lrcorner & \downarrow p \\ Y' & \xrightarrow{f} & Y \end{array} \quad p \in \mathcal{P} \rightsquigarrow \begin{array}{ccc} F(Y) & \xrightarrow{g^*} & F(Y') \\ \downarrow p^* & & \downarrow q^* \\ F(X) & \xrightarrow{f^*} & F(X') \end{array} \quad \begin{array}{l} \text{is vertically} \\ \text{right} \\ \text{adjointable} \end{array}$$

③ F is **$(\mathcal{I}, \mathcal{P})$ -biadjointable** if it is left $\mathcal{I}$ -adjointable, right $\mathcal{P}$ -adjointable and

$$\forall \text{ pullback } \begin{array}{ccc} X' & \xrightarrow{i} & X \\ q \downarrow & \lrcorner & \downarrow p \\ Y' & \xrightarrow{j} & Y \end{array} \quad \begin{array}{l} p \in \mathcal{P} \\ i \in \mathcal{I} \end{array} \rightsquigarrow \begin{array}{ccc} F(Y) & \xrightarrow{i^*} & F(Y') \\ \downarrow p^* & & \downarrow q^* \\ F(X) & \xrightarrow{j^*} & F(X') \end{array} \quad \text{is biadjointable}$$

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change, $\mathcal{D}$ 2-category
 $F, G: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$ and $\alpha: F \Rightarrow G$ natural transformation

$\alpha: F \Rightarrow G$ is **left $\mathcal{I}$ -adjointable** if

$$\begin{array}{ccc} F(Y) & \xrightarrow{j_Y} & F(X) \\ \downarrow \alpha_Y & \searrow j_X & \downarrow \alpha_X \\ G(Y) & \xrightarrow{i_Y} & G(X) \end{array}$$

is horizontally
left
adjointable

Same for **right $\mathcal{P}$ -adjointable**

and **$(\mathcal{I}, \mathcal{P})$ -biadjointable** = left $\mathcal{I}$ -adjointable + right $\mathcal{P}$ -adjointable

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change, $\mathcal{D}$ 2-category
 $F, G: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$ and $\alpha: F \Rightarrow G$ natural transformation

$\alpha: F \Rightarrow G$ is **left $\mathcal{I}$ -adjointable** if

$$\begin{array}{ccc} F(Y) & \xrightarrow{j_Y} & F(X) \\ \downarrow \alpha_Y & \downarrow j_X & \downarrow \alpha_X \\ G(Y) & \xrightarrow{j_Y} & G(X) \end{array}$$

is horizontally left adjointable

Same for **right $\mathcal{P}$ -adjointable**

and **$(\mathcal{I}, \mathcal{P})$ -biadjointable** = left $\mathcal{I}$ -adjointable + right $\mathcal{P}$ -adjointable

$\text{Fun}_{(\mathcal{I}, \mathcal{P})\text{-badj}}(\mathcal{C}^{\text{op}}, \mathcal{D})$:= locally full sub-2-category of $\text{Fun}(\mathcal{C}^{\text{op}}, \mathcal{D})$
 spanned by $(\mathcal{I}, \mathcal{P})$ -biadjointable functors and nat. transformations

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change, $\mathcal{D}$ 2-category
 $F, G: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$ and $\alpha: F \Rightarrow G$ natural transformation

$\alpha: F \Rightarrow G$ is **left $\mathcal{I}$ -adjointable** if
$$\begin{array}{ccc} F(Y) & \xrightarrow{j_Y} & F(X) \\ \downarrow \alpha_Y & \downarrow j_X & \downarrow \alpha_X \\ G(Y) & \xrightarrow{j_Y} & G(X) \end{array}$$
 is horizontally left adjointable

Same for **right $\mathcal{P}$ -adjointable**

and **$(\mathcal{I}, \mathcal{P})$ -biadjointable** = left $\mathcal{I}$ -adjointable + right $\mathcal{P}$ -adjointable

$\text{Fun}_{(\mathcal{I}, \mathcal{P})\text{-biadj}}(\mathcal{C}^{\text{op}}, \mathcal{D}) :=$ locally full sub-2-category of $\text{Fun}(\mathcal{C}^{\text{op}}, \mathcal{D})$
 spanned by $(\mathcal{I}, \mathcal{P})$ -biadjointable functors and nat. transformations

Remark. • If $\mathcal{P}$ (resp. $\mathcal{I}$) consists only of the equivalences in $\mathcal{C}$
 $(\mathcal{I}, \mathcal{P})$ -biadjoint. = left $\mathcal{I}$ -adjoint. (resp. right $\mathcal{P}$ -adjoint.)

• Any 2-functor preserves (bi)adjointable squares

C category, $I, P \subseteq C$ wide subcats closed under base change

An (I, P) -bidualizable functor $F: C^{op} \rightarrow B$ is **universal** if the restriction

$$F^*: \text{Fun}(B, D) \rightarrow \text{Fun}_{(I, P)\text{-bidual}}(C^{op}, D)$$

is an equivalence of 2-cats for every 2-cat D .

Theorem (Existence). $\exists h: C^{op} \rightarrow B_{I, P}(C)$ universal

$$\begin{array}{ccc} C^{op} & \xrightarrow{G} & D \\ \downarrow F & \nearrow & \uparrow \\ B & \dashrightarrow & \exists \bar{G} \end{array}$$

$\mathcal{C}$ category, $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change

An $(\mathcal{I}, \mathcal{P})$ -bidualisable functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}$ is **universal** if the restriction

$$F^*: \text{FUN}(\mathcal{B}, \mathcal{D}) \rightarrow \text{FUN}_{(\mathcal{I}, \mathcal{P})\text{-bidual}}(\mathcal{C}^{\text{op}}, \mathcal{D})$$

is an equivalence of 2-cats for every 2-cat $\mathcal{D}$.

Theorem (Existence). $\exists h: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}_{\mathcal{I}, \mathcal{P}}(\mathcal{C})$ universal

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} & \xrightarrow{G} & \mathcal{D} \\ \downarrow F & \nearrow & \uparrow \\ \mathcal{B} & \dashrightarrow & \exists \bar{G} \end{array}$$

An $(\mathcal{I}, \mathcal{P})$ -bidualisable functor $F: \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ is **free**

on a class $1_a \in F(a)$ with $a \in \mathcal{C}$ if $\forall (\mathcal{I}, \mathcal{P})$ -bidualisable functor

$G: \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ the evaluation at 1_a $\text{HOM}(F, G) \xrightarrow{\sim} G(a)$ is an equivalence

Theorem (Recognition principle). An $(\mathcal{I}, \mathcal{P})$ -bidualisable functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}$

is universal $\Leftrightarrow$ it is essentially surjective and for all $a \in \mathcal{C}$

$\text{HOM}(F(a), F(-)): \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ is a free $(\mathcal{I}, \mathcal{P})$ -bidualisable functor on $\text{id}_{F(a)}$

PROOF OF THE EXISTENCE OF THE UNIVERSAL BADJ FUNCTOR

Define a 3-cat BADJDATA_2 with the following data:

(1) Objects $(\mathbb{C}, \mathbb{Q}_L, \mathbb{Q}_R, \mathbb{Q}_B)$ where $\mathbb{C}$ is a 2-cat, $\mathbb{Q}_\bullet \subset \text{Fun}([1] \times [1], \mathbb{C})$

full subcategories s.t. $\{\text{equivalences}\} \subset \mathbb{Q}_\bullet$ and $\mathbb{Q}_B \subset \mathbb{Q}_L \cap \mathbb{Q}_R$

Denote L the horizontal maps in squares of $\mathbb{Q}_L$

and R a vertical " " " " $\mathbb{Q}_R$

(2) Morphisms are 2-functors $F: \mathbb{C} \rightarrow \mathbb{C}'$ satisfying $F(\mathbb{Q}_\bullet) \subset \mathbb{Q}'_\bullet$

(3) 2-cells are nat. transf. $\alpha: F \Rightarrow G$ s.t. $\forall \ell: X \rightarrow Y$ in L

$$\begin{array}{ccc} F(X) & \xrightarrow{F(\ell)} & F(Y) \\ \alpha_X \downarrow & & \downarrow \alpha_Y \\ G(X) & \xrightarrow{G(\ell)} & G(Y) \end{array} \text{ belongs to } \mathbb{Q}_L, \text{ dually for } R \text{ and } \mathbb{Q}_R$$

(4) No condition on higher cells

PROOF OF THE EXISTENCE OF THE UNIVERSAL BADJ FUNCTOR

Ex. $\mathcal{C}$ category with $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change

$j(\mathcal{C}, \mathcal{I}, \mathcal{P}) := (\mathcal{C}^{\text{op}}, \mathcal{Q}_L, \mathcal{Q}_R, \mathcal{Q}_B) \in \text{BADJDATA}_2$ where

$\mathcal{Q}_L = \text{pullbacks with horizontal maps in } \mathcal{I}$

$\mathcal{Q}_R = \text{pullbacks with vertical maps in } \mathcal{P}$

$$\mathcal{Q}_B = \mathcal{Q}_L \cap \mathcal{Q}_R$$

PROOF OF THE EXISTENCE OF THE UNIVERSAL B_{ADJ} FUNCTOR

Ex. $\mathcal{C}$ category with $\mathcal{I}, \mathcal{P} \subseteq \mathcal{C}$ wide subcats closed under base change

$j(\mathcal{C}, \mathcal{I}, \mathcal{P}) := (\mathcal{C}^{\text{op}}, \mathcal{Q}_L, \mathcal{Q}_R, \mathcal{Q}_B) \in \text{B}_{\text{ADJ}}\text{DATA}_2$ where

$\mathcal{Q}_L =$ pullbacks with horizontal maps in $\mathcal{I}$

$\mathcal{Q}_R =$ pullbacks with vertical maps in $\mathcal{P}$

$$\mathcal{Q}_B = \mathcal{Q}_L \cap \mathcal{Q}_R$$

Ex. Any 2-cat $\mathbb{D}$ determines $(\mathbb{D}, \mathcal{Q}_L, \mathcal{Q}_R, \mathcal{Q}_B) \in \text{B}_{\text{ADJ}}\text{DATA}_2$ where

$\mathcal{Q}_L =$ horizontally left adjointable squares

$\mathcal{Q}_R =$ vertically right adjointable squares

$\mathcal{Q}_B =$ biadjointable squares

Because 2-functors and 2-natural transformations preserve adjointability, this assignment defines a 3-functor

$$(-)_{\text{badj}} : \text{CAT}_2 \longrightarrow \text{B}_{\text{ADJ}}\text{DATA}_2$$

PROOF OF THE EXISTENCE OF THE UNIVERSAL BAJT FUNCTOR

For every (C, I, P) as before and 2-cat D , the forgetful map induces an equivalence

$$\text{Hom}_{\text{BAJTDATA}_2}(j(C, I, P), D_{\text{baj}}) \xrightarrow{\gamma} \text{Fun}_{(I, P)\text{-baj}}(C^{\text{op}}, D) \subseteq \text{Fun}(C^{\text{op}}, D)$$

Lemma 1. $(-)\text{baj}$ admits a left adjoint 3-functor $\hat{B}$

PROOF OF THE EXISTENCE OF THE UNIVERSAL BAJT FUNCTOR

For every (C, I, P) as before and 2-cat D , the forgetful map induces an equivalence

$$\text{Hom}_{\text{BAJTDATA}_2}(j(C, I, P), D_{\text{baj}}) \xrightarrow{\gamma} \text{Fun}_{(I, P)\text{-baj}}(C^{\text{op}}, D) \subseteq \text{Fun}(C^{\text{op}}, D)$$

Lemma 1. $(-)_{\text{baj}}$ admits a left adjoint 3-functor $\hat{B}$

Assuming Lemma 1, then

$$\text{Hom}_{\text{CAT}_2}(\hat{B}j(C, I, P), D) \xrightarrow{\varphi} \text{Hom}_{\text{BAJTDATA}_2}(j(C, I, P), D_{\text{baj}})$$

Then, there exists $B_{I, P}(C) := \hat{B}j(C, I, P)$ and $h := \gamma \circ \varphi(\text{id})$ where h is universal and (I, P) -bajadjoinable by construction.

□

IDEA OF THE PROOF OF THE LEMMA 1

Define the *walking adjunction* $\mathbf{Adj}$ in the $\mathcal{Z}$ -category with two morphisms $\ell: \oplus \rightarrow \oplus$ and $\pi: \oplus \rightarrow \ominus$ s.t. $\ell \dashv \pi$.

IDEA OF THE PROOF OF THE LEMMA 1

Define the *walking adjunction* Adj in the 2-category with two morphisms $\ell: \oplus \rightarrow \oplus$ and $\pi: \oplus \rightarrow \oplus$ s.t. $\ell \dashv \pi$.

Squares in $\mathbb{D}$

Lemma. (1) $(id \times \pi)^*: Fun([1] \times Adj, \mathbb{D}) \rightarrow Fun([1] \times [1], \mathbb{D})$ is a

locally full inclusion with image the horizontally left adjointable squares and horizontally left adjointable natural transformations.

(2) Dually for $(\ell \times id)^*$

(3) $(\ell \times \pi)^*$ similarly with image biadjointable squares and nat. transg.

IDEA OF THE PROOF OF THE LEMMA 1

Define the **walking adjunction** ADJ in the 2 -category with two morphisms $\ell: \oplus \rightarrow \oplus$ and $\pi: \oplus \rightarrow \oplus$ s.t. $\ell \dashv \pi$.

Squares in $\mathcal{D}$

Lemma. (1) $(id \times \pi)^*: FUN([1] \times ADJ, \mathcal{D}) \rightarrow FUN([1] \times [1], \mathcal{D})$ is a

locally full inclusion with image the horizontally left adjointable squares and horizontally left adjointable natural transformations.

(2) Dually for $(\ell \times id)^*$

(3) $(\ell \times \pi)^*$ similarly with image biadjointable squares and nat. transf.

Idea. (1) and (2) follow from the universal property of ADJ .

(3) in particular shows that biadjointability is symmetric.

$$\pi^*: FUN(ADJ, \mathcal{D}) \rightarrow FUN([1], \mathcal{D}) \quad \ell^*: FUN(ADJ, FUN^{lad}([1], \mathcal{D})) \rightarrow FUN([1] \times [1], \mathcal{D})$$

$\rightsquigarrow$ Image $FUN^{lad}([1], \mathcal{D})$ $\rightsquigarrow$ Image biadjointable squares □

IDEA OF THE PROOF OF THE LEMMA 1

$\forall (\mathbb{C}, Q_L, Q_R, Q_B) \in \text{BADJDATA}_2$
 $(-)^{\text{body}}$ admits a left adjoint $\Leftrightarrow \exists \hat{B}(\mathbb{C})$ 2-cat and $h: \mathbb{C} \rightarrow \hat{B}(\mathbb{C})$ s.t.
 $\text{FUN}(\hat{B}(\mathbb{C}), \mathbb{D}) \cong \text{FUN}(\mathbb{C}, Q_L, Q_R, Q_B), \mathbb{D}^{\text{body}})$

IDEA OF THE PROOF OF THE LEMMA 1

$$\forall (\mathbb{C}, Q_L, Q_R, Q_B) \in \mathcal{B}_{\text{ADJ DATA}_2}$$

$(-)^{\text{body}}$ admits a left adjoint $\Leftrightarrow \exists \hat{\mathbb{B}}(\mathbb{C})$ 2-cat and $h: \mathbb{C} \rightarrow \hat{\mathbb{B}}(\mathbb{C})$ s.t.

$$\text{FUN}(\hat{\mathbb{B}}(\mathbb{C}), \mathbb{D}) \cong \text{FUN}(\mathbb{C}, Q_L, Q_R, Q_B), \mathbb{D}^{\text{body}}$$

We construct $\hat{\mathbb{B}}(\mathbb{C})$ and h in three steps:

By previous lemma,

$$(1) \quad \begin{array}{ccc} \bigsqcup_{\pi_0(Q_L)} ([1] \times [1]) & \xrightarrow{U(\text{id} \times \eta)} & \bigsqcup ([1] \times \text{ADJ}) \\ \downarrow & & \downarrow \pi_0(Q_L) \\ \mathbb{C} & \xrightarrow{\quad} & \mathbb{C}_L \end{array} \quad \begin{array}{l} \text{FUN}(-, \mathbb{D}) \text{ FUN}(\mathbb{C}_L, \mathbb{D}) \rightarrow \text{FUN}(\mathbb{C}, \mathbb{D}) \\ \rightsquigarrow \text{in locally full inclusion} \\ \text{with image the functor} \\ \text{mapping } Q \text{ to hor. left adj. sq.} \end{array}$$

(2) Same for $Q_R \rightsquigarrow \mathbb{C}_L \rightarrow \mathbb{C}_{L,R}$

(3) Same for Q_B , with

$$\bigsqcup_{\pi_0(Q_B)} ([1] \times [1]) \xrightarrow{U(\ell \times \eta)} \bigsqcup_{\pi_0(Q_B)} (\text{ADJ} \times \text{ADJ})$$

$$\left. \begin{array}{l} \rightsquigarrow \mathbb{C}_{L,R} \rightarrow \hat{\mathbb{B}}(\mathbb{C}) \\ \text{Then } h: \mathbb{C} \rightarrow \mathbb{C}_L \rightarrow \mathbb{C}_{L,R} \rightarrow \hat{\mathbb{B}}(\mathbb{C}) \end{array} \right\}$$

□

PROOF OF THE RECOGNITION PRINCIPLE

Theorem (Recognition principle). An (I, P) -biadjointable functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}$ is universal $\Leftrightarrow$ it is essentially surjective and for all $a \in \mathcal{C}$
 $\text{Hom}(F(a), F(-)): \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ is a free (I, P) -biadjointable functor on $\text{id}_{F(a)}$

Proof plan:

- (1) $\text{Hom}(F(a), F(-)): \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ is a free on a (I, P) -biadjointable functor
- (2) Any universal (I, P) -biadjointable functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}$ is essentially surj.
- (3) If $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}$ is essentially surjective and for all $a \in \mathcal{C}$
 $\text{Hom}(F(a), F(-)): \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$
is a free (I, P) -biadjointable functor on $\text{id}_{F(a)}$, then F is universal.

PROOF OF THE RECOGNITION PRINCIPLE

(1) $\text{Hom}_{\mathbb{B}}(F(a), F(-)) : \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ is a free on a (Σ, P) -bifree object

It can be written as the composition $\mathcal{C}^{\text{op}} \xrightarrow{F} \mathbb{B} \xrightarrow{\text{Hom}(F(a), -)} \text{CAT}$

$\Rightarrow$ it is (Σ, P) -bifree

$G : \mathcal{C}^{\text{op}} \rightarrow \text{CAT}$ (Σ, P) -bifree $\leadsto \exists \bar{G} : \mathbb{B} \rightarrow \text{CAT}$ and a diagram

$$\begin{array}{ccc} \text{Hom}(\text{Hom}(F(a), -), \bar{G}) & \xrightarrow[\sim]{\text{ev}_{\text{id}_{F(a)}}} & \bar{G}(F(a)) \\ \sim \downarrow \hat{F}^* & & \parallel \\ \text{Hom}(\text{Hom}(F(a), F(-)), G) & \xrightarrow{\text{ev}_{\text{id}_{F(a)}}} & G(a) \end{array} \Rightarrow \text{ev}_{\text{id}_{F(a)}} \text{ is an equivalence}$$

• The top map is an equivalence by 2-cat Yoneda

• $\hat{F}^*$ is an equivalence induced by $F^* : \text{Fun}(\mathbb{B}, \text{CAT}) \xrightarrow{\sim} \text{Fun}_{(\Sigma, P)\text{-bifree}}(\mathcal{C}^{\text{op}}, \text{CAT})$

PROOF OF THE RECOGNITION PRINCIPLE

(2) Any universal $(\mathcal{I}, \mathcal{P})$ -biodjointable functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{B}$ is essentially surj.

$\mathcal{I} \subset \mathcal{B}$ full 2-subcat spanned by objects in the image of F

$\leadsto \hat{F}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{I}$ is $(\mathcal{I}, \mathcal{P})$ -biodjointable

$\leadsto \hat{F}$ admits an extension to $\mathcal{B}$ s.t.

$$\begin{array}{ccccc}
 & & \mathcal{C}^{\text{op}} & & \\
 & \searrow F & \downarrow \hat{F} & \swarrow F & \\
 \mathcal{B} & \dashrightarrow & \mathcal{I} & \hookrightarrow & \mathcal{B}
 \end{array}
 \quad \text{commutes}$$

$\leadsto$ By the universal property of $\mathcal{B}$, $\mathcal{B} \dashrightarrow \mathcal{I} \hookrightarrow \mathcal{B}$ is the identity

$\leadsto \mathcal{I} \hookrightarrow \mathcal{B}$ is essentially surjective $\leadsto F$ is essentially surjective

PROOF OF THE RECOGNITION PRINCIPLE

(3) If $F: C^{\text{op}} \rightarrow B$ essentially surjective and for all $a \in C$

$$\text{Hom}(F(a), F(-)): C^{\text{op}} \rightarrow \text{CAT}$$

is a free (I, P) -biadjointable functor on $\text{id}_{F(a)}$, then F is universal.

By the universal property of $h: C^{\text{op}} \rightarrow B_{I, P}(C)$, F extends to $\bar{F}: B_{I, P}(C) \rightarrow B$

It will suffice to show that $\bar{F}$ is an equivalence

Since $F = \bar{F} \circ h$ is essentially surj., $\bar{F}$ is essentially surj.

We want $\bar{F}: \text{Hom}(X, Y) \rightarrow \text{Hom}(\bar{F}(X), \bar{F}(Y))$ is an equivalence

(2) h essentially surj., it suffices $\bar{F}: \text{Hom}(h(a), h(-)) \rightarrow \text{Hom}(F(a), F(-))$ is an equivalence

$$\left. \begin{array}{l} (1) \text{Hom}(h(a), h(-)) \text{ free on } \text{id}_{h(a)} \\ \text{Hom}(F(a), F(-)) \text{ free on } \text{id}_{F(a)} \end{array} \right\} \begin{array}{l} \leadsto \bar{F} \text{ is an equivalence} \\ \nwarrow \bar{F}(\text{id}_{h(a)}) = \text{id}_{F(a)} \end{array}$$

□