Linear Regression

Pramod Duvvuri 4/10/2019

```
# Load libraries
library(MASS)
library(ISLR)
names(Boston)
## [1] "crim"
                   "zn"
                             "indus"
                                        "chas"
                                                                       "age"
                                                  "nox"
   [8] "dis"
                             "tax"
##
                   "rad"
                                        "ptratio" "black"
                                                             "lstat"
                                                                        "medv"
?Boston
plot(medv~lstat,Boston)
                          00
    4
    30
                                                 00
    20
                                                                                          0
    10
                                                                                        0
                            10
                                                  20
                                                                        30
                                                 Istat
# Simple linear regression
fit1=lm(medv~lstat,data=Boston)
fit1
##
## lm(formula = medv ~ lstat, data = Boston)
## Coefficients:
## (Intercept)
                       lstat
                       -0.95
         34.55
summary(fit1)
##
## lm(formula = medv ~ lstat, data = Boston)
## Residuals:
```

```
10 Median
                                3Q
## -15.168 -3.990 -1.318
                             2.034 24.500
##
## Coefficients:
##
               Estimate Std. Error t value Pr(>|t|)
## (Intercept) 34.55384
                           0.56263
                                     61.41
                                              <2e-16 ***
               -0.95005
                           0.03873 -24.53
                                             <2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 6.216 on 504 degrees of freedom
## Multiple R-squared: 0.5441, Adjusted R-squared: 0.5432
## F-statistic: 601.6 on 1 and 504 DF, p-value: < 2.2e-16
plot(medv~lstat,Boston)
abline(fit1,col="red")
    4
    30
medv
                                               00
                                                                     0
    20
                                                                                       0
    9
                                                                                     0
                           10
                                                 20
                                                                      30
                                                Istat
names(fit1)
    [1] "coefficients" "residuals"
                                         "effects"
                                                         "rank"
    [5] "fitted.values" "assign"
                                         "qr"
                                                         "df.residual"
   [9] "xlevels"
                        "call"
                                                         "model"
                                         "terms"
confint(fit1)
##
                   2.5 %
                             97.5 %
## (Intercept) 33.448457 35.6592247
               -1.026148 -0.8739505
predict(fit1,data.frame(lstat=c(5,10,15)),interval="confidence")
##
          fit
                   lwr
## 1 29.80359 29.00741 30.59978
## 2 25.05335 24.47413 25.63256
## 3 20.30310 19.73159 20.87461
# Multiple linear regression
fit2=lm(medv~lstat+age,data=Boston)
```


```
summary(fit2)
##
## Call:
## lm(formula = medv ~ lstat + age, data = Boston)
## Residuals:
##
      Min
               1Q Median
                               3Q
                                      Max
## -15.981 -3.978 -1.283 1.968
                                  23.158
##
## Coefficients:
              Estimate Std. Error t value Pr(>|t|)
                          0.73085 45.458 < 2e-16 ***
## (Intercept) 33.22276
## lstat
              -1.03207
                          0.04819 -21.416 < 2e-16 ***
              0.03454
                          0.01223
                                    2.826 0.00491 **
## age
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 6.173 on 503 degrees of freedom
## Multiple R-squared: 0.5513, Adjusted R-squared: 0.5495
## F-statistic:
                 309 on 2 and 503 DF, p-value: < 2.2e-16
fit3=lm(medv~.,Boston)
summary(fit3)
##
## Call:
## lm(formula = medv ~ ., data = Boston)
##
## Residuals:
##
      Min
               1Q Median
                               3Q
                                     Max
## -15.595 -2.730 -0.518
                           1.777 26.199
##
## Coefficients:
                Estimate Std. Error t value Pr(>|t|)
##
## (Intercept) 3.646e+01 5.103e+00
                                    7.144 3.28e-12 ***
## crim
              -1.080e-01 3.286e-02 -3.287 0.001087 **
## zn
              4.642e-02 1.373e-02
                                     3.382 0.000778 ***
## indus
               2.056e-02 6.150e-02
                                     0.334 0.738288
## chas
              2.687e+00 8.616e-01
                                    3.118 0.001925 **
## nox
              -1.777e+01 3.820e+00 -4.651 4.25e-06 ***
                                    9.116 < 2e-16 ***
              3.810e+00 4.179e-01
## rm
## age
              6.922e-04 1.321e-02
                                     0.052 0.958229
## dis
              -1.476e+00 1.995e-01 -7.398 6.01e-13 ***
## rad
              3.060e-01 6.635e-02
                                     4.613 5.07e-06 ***
              -1.233e-02 3.760e-03 -3.280 0.001112 **
## tax
              -9.527e-01 1.308e-01 -7.283 1.31e-12 ***
## ptratio
## black
              9.312e-03 2.686e-03
                                    3.467 0.000573 ***
## 1stat
              -5.248e-01 5.072e-02 -10.347 < 2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 4.745 on 492 degrees of freedom
## Multiple R-squared: 0.7406, Adjusted R-squared: 0.7338
```

par(mfrow=c(2,2)) plot(fit3) Standardized residuals Residuals vs Fitted Normal Q-Q 9 03692 20 Residuals 2 0 7 -20 0 10 20 30 40 -3 -2 0 2 3 Theoretical Quantiles Fitted values /IStandardized residuals Scale-Location Standardized residuals Residuals vs Leverage 3899 2.0 1.0 0 Cooksodistance 0.0 4 0 10 20 30 40 0.00 0.05 0.10 0.15 0.20 0.25 0.30 Fitted values Leverage fit4=update(fit3,~.-age-indus) summary(fit4) ## ## Call: ## lm(formula = medv ~ crim + zn + chas + nox + rm + dis + rad + tax + ptratio + black + lstat, data = Boston) ## ## Residuals: ## ## Min Median 3Q Max 1Q -15.5984 -2.7386 -0.5046 ## 1.7273 26.2373 ## ## Coefficients: ## Estimate Std. Error t value Pr(>|t|) 36.341145 5.067492 7.171 2.73e-12 *** ## (Intercept) crim -0.108413 0.032779 -3.307 0.001010 ** ## ## zn 0.045845 0.013523 3.390 0.000754 *** 0.854240 ## chas 2.718716 3.183 0.001551 ** -17.376023 3.535243 -4.915 1.21e-06 *** ## nox 0.406316 < 2e-16 *** 3.801579 9.356 ## rm0.185731 -8.037 6.84e-15 *** ## dis -1.492711 0.063402 0.299608 4.726 3.00e-06 *** ## rad -3.493 0.000521 *** ## tax -0.011778 0.003372 -0.946525 0.129066 -7.334 9.24e-13 *** ## ptratio ## black 0.009291 0.002674 3.475 0.000557 *** -0.522553 0.047424 -11.019 < 2e-16 *** ## 1stat ## 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1 ## Signif. codes: ## ## Residual standard error: 4.736 on 494 degrees of freedom


F-statistic: 108.1 on 13 and 492 DF, p-value: < 2.2e-16

```
## Multiple R-squared: 0.7406, Adjusted R-squared: 0.7348
## F-statistic: 128.2 on 11 and 494 DF, p-value: < 2.2e-16
# Nonlinear terms and Interactions
fit5=lm(medv~lstat*age,Boston)
summary(fit5)
##
## Call:
## lm(formula = medv ~ lstat * age, data = Boston)
## Residuals:
##
      Min
               1Q Median
                              3Q
                                     Max
## -15.806 -4.045 -1.333
                            2.085 27.552
##
## Coefficients:
##
                Estimate Std. Error t value Pr(>|t|)
## (Intercept) 36.0885359 1.4698355 24.553 < 2e-16 ***
              ## age
              -0.0007209 0.0198792 -0.036
                                             0.9711
## lstat:age
              0.0041560 0.0018518
                                     2.244
                                             0.0252 *
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6.149 on 502 degrees of freedom
## Multiple R-squared: 0.5557, Adjusted R-squared: 0.5531
## F-statistic: 209.3 on 3 and 502 DF, p-value: < 2.2e-16
fit6=lm(medv~lstat + I(lstat^2),Boston);
summary(fit6)
##
## lm(formula = medv ~ lstat + I(lstat^2), data = Boston)
##
## Residuals:
##
       Min
                 1Q
                     Median
                                  3Q
                                          Max
## -15.2834 -3.8313 -0.5295
                              2.3095 25.4148
##
## Coefficients:
               Estimate Std. Error t value Pr(>|t|)
## (Intercept) 42.862007
                         0.872084
                                   49.15
                                            <2e-16 ***
                          0.123803 -18.84
## 1stat
              -2.332821
                                            <2e-16 ***
              0.043547
                         0.003745
                                   11.63
## I(lstat^2)
                                            <2e-16 ***
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 5.524 on 503 degrees of freedom
## Multiple R-squared: 0.6407, Adjusted R-squared: 0.6393
## F-statistic: 448.5 on 2 and 503 DF, p-value: < 2.2e-16
attach(Boston)
par(mfrow=c(1,1))
plot(medv~lstat)
points(lstat,fitted(fit6),col="red",pch=20)
fit7=lm(medv~poly(lstat,4))
```

points(lstat,fitted(fit7),col="blue",pch=20)



plot(1:20,1:20,pch=1:20,cex=2)



Qualitative predictors

fix(Carseats)

names(Carseats)

```
## [1] "Sales" "CompPrice" "Income" "Advertising" "Population"
## [6] "Price" "ShelveLoc" "Age" "Education" "Urban"
## [11] "US"
```

summary(Carseats)

Sales CompPrice Income Advertising Min. : 21.00 Min. : 0.000 Min. : 77 Min. : 0.000 ## ## 1st Qu.: 5.390 1st Qu.:115 1st Qu.: 42.75 1st Qu.: 0.000 Median : 7.490 Median:125 Median : 69.00 Median : 5.000

```
Mean : 7.496
                    Mean
                           :125
                                  Mean
                                         : 68.66
                                                          : 6.635
                                                   Mean
##
   3rd Qu.: 9.320
                    3rd Qu.:135
                                  3rd Qu.: 91.00
                                                   3rd Qu.:12.000
                    Max.
                                  Max.
                                                          :29.000
          :16.270
                           :175
                                         :120.00
                                                   Max.
##
     Population
                                    ShelveLoc
                       Price
                                                     Age
##
  Min.
          : 10.0
                   Min.
                          : 24.0
                                   Bad
                                        : 96
                                                Min.
                                                       :25.00
                   1st Qu.:100.0
                                   Good : 85
                                                1st Qu.:39.75
##
   1st Qu.:139.0
  Median :272.0
                   Median :117.0
                                   Medium:219
                                                Median :54.50
##
  Mean
         :264.8
                   Mean :115.8
                                                Mean
                                                      :53.32
##
   3rd Qu.:398.5
                   3rd Qu.:131.0
                                                3rd Qu.:66.00
##
  Max.
          :509.0
                   Max.
                         :191.0
                                                Max. :80.00
     Education
                  Urban
          :10.0
                  No :118
                            No :142
## Min.
## 1st Qu.:12.0
                  Yes:282
                            Yes:258
## Median :14.0
## Mean
          :13.9
## 3rd Qu.:16.0
## Max.
          :18.0
fit1=lm(Sales~.+Income:Advertising+Age:Price,Carseats)
summary(fit1)
##
## Call:
## lm(formula = Sales ~ . + Income:Advertising + Age:Price, data = Carseats)
## Residuals:
                               3Q
      Min
               1Q Median
                                      Max
## -2.9208 -0.7503 0.0177 0.6754 3.3413
##
## Coefficients:
##
                       Estimate Std. Error t value Pr(>|t|)
## (Intercept)
                      6.5755654 1.0087470
                                             6.519 2.22e-10 ***
## CompPrice
                      0.0929371 0.0041183 22.567 < 2e-16 ***
## Income
                      0.0108940 0.0026044
                                             4.183 3.57e-05 ***
## Advertising
                      0.0702462 0.0226091
                                             3.107 0.002030 **
## Population
                      0.0001592 0.0003679
                                             0.433 0.665330
## Price
                     -0.1008064 0.0074399 -13.549 < 2e-16 ***
## ShelveLocGood
                      4.8486762 0.1528378 31.724 < 2e-16 ***
## ShelveLocMedium
                      1.9532620 0.1257682 15.531 < 2e-16 ***
## Age
                     -0.0579466 0.0159506 -3.633 0.000318 ***
## Education
                     -0.0208525 0.0196131
                                           -1.063 0.288361
## UrbanYes
                      0.1401597 0.1124019
                                             1.247 0.213171
## USYes
                     -0.1575571 0.1489234 -1.058 0.290729
## Income: Advertising 0.0007510 0.0002784
                                             2.698 0.007290 **
## Price:Age
                      0.0001068 0.0001333
                                           0.801 0.423812
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
## Residual standard error: 1.011 on 386 degrees of freedom
## Multiple R-squared: 0.8761, Adjusted R-squared: 0.8719
## F-statistic:
                 210 on 13 and 386 DF, p-value: < 2.2e-16
contrasts(Carseats$ShelveLoc)
```

Good Medium

```
## Bad
             0
                    0
## Good
             1
                     0
## Medium
# Writing R functions
regplot=function(x,y){
  fit=lm(y~x)
  plot(x,y)
  abline(fit,col="red")
attach(Carseats)
regplot(Price,Sales)
                                           0
                                 0
    15
                        0
                         0
    10
                      0 0
                               00
                             0 0
    2
                                                      0
    0
                      50
                                             100
                                                                     150
                                                  Х
regplot=function(x,y,...){
  fit=lm(y~x)
  plot(x,y,...)
  abline(fit,col="red")
}
regplot(Price, Sales, xlab="Price", ylab="Sales", col="blue", pch=20)
```

