

---

# GEOMETRY

## Through Algebra

---

D.V.S. NIKHIL





# Contents

1	Triangle	1
1.1	Vectors . . . . .	1



# Chapter 1

## Triangle

Consider a triangle with vertices

$$\mathbf{A} = \begin{pmatrix} -5 \\ -2 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} -6 \\ 4 \end{pmatrix} \quad (1.1)$$

### 1.1. Vectors

1.1.1. the direction vector of  $AB$  is defined as

$$\mathbf{B} - \mathbf{A} \quad (1.1.1.1)$$

Find the direction vectors of  $AB$ ,  $BC$  and  $CA$ .

**Solution:**

(a) The Direction vector of  $AB$  is

$$= \mathbf{B} - \mathbf{A} \quad (1.1.1.2)$$

$$\therefore = \begin{pmatrix} -1 - (-5) \\ -1 - (-2) \end{pmatrix} \quad (1.1.1.3)$$

$$= \begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad (1.1.1.4)$$

(b) The Direction vector of  $BC$

$$= \mathbf{C} - \mathbf{B} \quad (1.1.1.5)$$

$$= \begin{pmatrix} -6 - (-1) \\ 4 - (-1) \end{pmatrix} \quad (1.1.1.6)$$

$$= \begin{pmatrix} -5 \\ 5 \end{pmatrix} \quad (1.1.1.7)$$

(c) The Direction vector of  $CA$

$$= \mathbf{A} - \mathbf{C} \quad (1.1.1.8)$$

$$= \begin{pmatrix} -5 - (-6) \\ -2 - (4) \end{pmatrix} \quad (1.1.1.9)$$

$$= \begin{pmatrix} 1 \\ -6 \end{pmatrix} \quad (1.1.1.10)$$

1.1.2. The length of side  $AB, BC$  and  $AC$  is **Solution:** Given,

$$\mathbf{A} = \begin{pmatrix} -5 \\ -2 \end{pmatrix}, \quad (1.1.2.1)$$

$$\mathbf{B} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}, \quad (1.1.2.2)$$

$$\mathbf{C} = \begin{pmatrix} -6 \\ 4 \end{pmatrix} \quad (1.1.2.3)$$

Now solving for  $AB$ ,

$$\|\mathbf{A} - \mathbf{B}\| = \sqrt{(\mathbf{A} - \mathbf{B})^\top (\mathbf{A} - \mathbf{B})} \quad (1.1.2.4)$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -5 \\ -2 \end{pmatrix} - \begin{pmatrix} -1 \\ -1 \end{pmatrix} = \begin{pmatrix} -4 \\ -1 \end{pmatrix} \quad (1.1.2.5)$$

$$\|\mathbf{A} - \mathbf{B}\| = \sqrt{\begin{pmatrix} -4 & -1 \end{pmatrix} \begin{pmatrix} -4 \\ -1 \end{pmatrix}} \quad (1.1.2.6)$$

$$= \sqrt{(4)^2 + (1)^2} \quad (1.1.2.7)$$

$$\Rightarrow \|\mathbf{A} - \mathbf{B}\| = \sqrt{17} \quad (1.1.2.8)$$

Now solving for  $BC$ ,

$$\|\mathbf{B} - \mathbf{C}\| = \sqrt{(\mathbf{B} - \mathbf{C})^\top (\mathbf{B} - \mathbf{C})} \quad (1.1.2.9)$$

$$\mathbf{B} - \mathbf{C} = \begin{pmatrix} 5 \\ -5 \end{pmatrix} \quad (1.1.2.10)$$

$$\|\mathbf{B} - \mathbf{C}\| = \sqrt{\begin{pmatrix} 5 & -5 \end{pmatrix} \begin{pmatrix} 5 \\ -5 \end{pmatrix}} \quad (1.1.2.11)$$

$$= \sqrt{(5)^2 + (5)^2} \quad (1.1.2.12)$$

$$\Rightarrow \|\mathbf{B} - \mathbf{C}\| = \sqrt{50} \quad (1.1.2.13)$$

Now solving for  $AC$ ,

$$\|\mathbf{A} - \mathbf{C}\| = \sqrt{(\mathbf{A} - \mathbf{C})^\top (\mathbf{A} - \mathbf{C})} \quad (1.1.2.14)$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} 1 \\ -6 \end{pmatrix} \quad (1.1.2.15)$$

$$\|\mathbf{A} - \mathbf{C}\| = \sqrt{\begin{pmatrix} 1 & -6 \end{pmatrix} \begin{pmatrix} 1 \\ -6 \end{pmatrix}} \quad (1.1.2.16)$$

$$= \sqrt{(1)^2 + (6)^2} \quad (1.1.2.17)$$

$$\Rightarrow \|\mathbf{A} - \mathbf{C}\| = \sqrt{37} \quad (1.1.2.18)$$



1.1.3. Points  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are defined to be collinear if

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} = 2 \quad (1.1.3.1)$$

**Solution:**

Given that,

$$\mathbf{A} = \begin{pmatrix} -5 \\ -2 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} -1 \\ -1 \end{pmatrix} \quad \mathbf{C} = \begin{pmatrix} -6 \\ 4 \end{pmatrix} \quad (1.1.3.2)$$

Given that  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are collinear if

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} < 3 \quad (1.1.3.3)$$

Let

$$\mathbf{R} = \begin{pmatrix} 1 & 1 & 1 \\ -5 & -1 & -6 \\ -2 & -1 & 4 \end{pmatrix} \quad (1.1.3.4)$$

$$(1.1.3.5)$$

The matrix  $\mathbf{R}$  can be row reduced as follows,

$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & 4 & -1 \\ -2 & -1 & -4 \end{pmatrix} \xleftrightarrow{R_3 \leftarrow -2R_1 + R_3} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 4 & -1 \\ 0 & 1 & 6 \end{pmatrix} \quad (1.1.3.6)$$

$$\xleftrightarrow{R_3 \leftarrow R_3 - \frac{1}{4}R_2} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 4 & -1 \\ 0 & 0 & \frac{25}{4} \end{pmatrix} \quad (1.1.3.7)$$

$$\xleftrightarrow{R_2 \leftarrow \frac{1}{4}R_2} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & -\frac{1}{4} \\ 0 & 0 & \frac{25}{4} \end{pmatrix} \quad (1.1.3.8)$$

There are no zero rows. So,

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} = 3 \quad (1.1.3.9)$$

Hence, from (??) the points  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are not collinear.

Figure 1.1:  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  plot

From Fig. 1.1, We can see that  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are not collinear .

1.1.4. The parametric form of the equation of  $AB$  is

$$\mathbf{x} = \mathbf{A} + k\mathbf{m} \quad (1.1.4.1)$$

where

$$\mathbf{m} = \mathbf{B} - \mathbf{A} \quad (1.1.4.2)$$

is the direction vector of  $AB$ . Find the parametric equations of  $AB$ ,  $BC$  and  $CA$ . **Solution:**

The parametric equation for  $AB$  is given by

$$\mathbf{x} = \mathbf{A} + k\mathbf{m} \quad (1.1.4.3)$$

$$\text{where, } \mathbf{m} = \mathbf{B} - \mathbf{A} \quad (1.1.4.4)$$

$$= \begin{pmatrix} -1 \\ -1 \end{pmatrix} - \begin{pmatrix} -5 \\ -2 \end{pmatrix} \quad (1.1.4.5)$$

$$= \begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad (1.1.4.6)$$

Hence we get,

$$\mathbf{AB} : \mathbf{x} = \begin{pmatrix} -5 \\ -2 \end{pmatrix} + k \begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad (1.1.4.7)$$

Similarly,

$$\mathbf{BC} : \mathbf{x} = \begin{pmatrix} -1 \\ -1 \end{pmatrix} + k \begin{pmatrix} 5 \\ -5 \end{pmatrix} \quad (1.1.4.8)$$

$$\mathbf{CA} : \mathbf{x} = \begin{pmatrix} -6 \\ 4 \end{pmatrix} + k \begin{pmatrix} 1 \\ -6 \end{pmatrix} \quad (1.1.4.9)$$

1.1.5. The normal form of the equation of  $\mathbf{AB}$  is

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \quad (1.1.5.1)$$

where

$$\mathbf{n}^\top \mathbf{m} = \mathbf{n}^\top (\mathbf{B} - \mathbf{A}) = 0 \quad (1.1.5.2)$$

or,

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.3)$$

then find the normal form of the equations of  $\mathbf{AB}$ ,  $\mathbf{BC}$  and  $\mathbf{CA}$

**Solution:** :

The normal equation for the side  $AB$  is

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \quad (1.1.5.4)$$

$$\implies \mathbf{n}^\top \mathbf{x} = \mathbf{n}^\top \mathbf{A} \quad (1.1.5.5)$$

Now our task is to find the  $\mathbf{n}$  so that we can find  $\mathbf{n}^\top$ . As given.

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.6)$$

Here  $\mathbf{m} = \mathbf{B} - \mathbf{A}$  for side  $\mathbf{AB}$

$$\Rightarrow \mathbf{m} = \begin{pmatrix} -1 \\ -1 \end{pmatrix} - \begin{pmatrix} -5 \\ -2 \end{pmatrix} \quad (1.1.5.7)$$

$$= \begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad (1.1.5.8)$$

Now as we have obtained vector  $\mathbf{m}$ .we can use this to obtain vector  $\mathbf{n}$

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \end{pmatrix} \quad (1.1.5.9)$$

The transpose of  $\mathbf{n}$  is

$$\mathbf{n}^\top = \begin{pmatrix} 1 & 4 \end{pmatrix} \quad (1.1.5.10)$$

Hence the normal equation of side  $AB$  is

$$\begin{pmatrix} 1 & 4 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 1 & 4 \end{pmatrix} \begin{pmatrix} -5 \\ -2 \end{pmatrix} \quad (1.1.5.11)$$

$$\Rightarrow \begin{pmatrix} 1 & -4 \end{pmatrix} \mathbf{x} = -13 \quad (1.1.5.12)$$

Figure 1.2: The line  $\mathbf{AB}$  plotted

Similarly

$$\implies \mathbf{BC} : \begin{pmatrix} 5 & -5 \end{pmatrix} \mathbf{x} = -50 \quad (1.1.5.13)$$

Figure 1.3: The line  $\mathbf{BC}$  plotted

$$\implies \mathbf{CA} : \begin{pmatrix} -6 & 1 \end{pmatrix} \mathbf{x} = -32 \quad (1.1.5.14)$$

Figure 1.4: The line  $\mathbf{CA}$  plotted

1.1.6. Find the area of the  $\triangle ABC$  **Solution:**

Given

$$\mathbf{A} = \begin{pmatrix} -5 \\ -2 \end{pmatrix}; \mathbf{B} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}; \mathbf{C} = \begin{pmatrix} -6 \\ 4 \end{pmatrix} \quad (1.1.6.1)$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -5 \\ -2 \end{pmatrix} - \begin{pmatrix} -1 \\ -1 \end{pmatrix} = \begin{pmatrix} -4 \\ -1 \end{pmatrix} \quad (1.1.6.2)$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} -5 \\ -2 \end{pmatrix} - \begin{pmatrix} -6 \\ 4 \end{pmatrix} = \begin{pmatrix} 1 \\ -6 \end{pmatrix} \quad (1.1.6.3)$$

$$\therefore (\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C}) = \begin{vmatrix} -4 & -1 \\ 1 & -6 \end{vmatrix} \quad (1.1.6.4)$$

$$= -4 \times -6 - 1 \times (1) \quad (1.1.6.5)$$

$$= 24 - 1 \quad (1.1.6.6)$$

$$= 23 \quad (1.1.6.7)$$

$$\Rightarrow \frac{1}{2} \|(\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C})\| = \frac{1}{2} \|23\| = \frac{23}{2} \quad (1.1.6.8)$$

1.1.7. Find the angles  $A, B, C$  if

$$\cos A \triangleq \frac{(\mathbf{B} - \mathbf{A})^\top \mathbf{C} - \mathbf{A}}{\|\mathbf{B} - \mathbf{A}\| \|\mathbf{C} - \mathbf{A}\|} \quad (1.1.7.1)$$

**Solution:**

From the given values of  $\mathbf{A}, \mathbf{B}, \mathbf{C}$ ,

(a) Finding the value of angle A

$$\mathbf{B} - \mathbf{A} = \begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad (1.1.7.2)$$

and

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} -1 \\ 6 \end{pmatrix} \quad (1.1.7.3)$$

also calculating the values of norms

$$\|\mathbf{B} - \mathbf{A}\| = \sqrt{17} \quad (1.1.7.4)$$

$$\|\mathbf{C} - \mathbf{A}\| = \sqrt{37} \quad (1.1.7.5)$$

and by doing matrix multiplication we get,

$$(\mathbf{B} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A}) = \begin{pmatrix} 4 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 6 \end{pmatrix} = 2 \quad (1.1.7.6)$$

So, we get

$$\cos A = \frac{2}{\sqrt{17}\sqrt{37}} \quad (1.1.7.7)$$

$$= \frac{2}{\sqrt{629}} \quad (1.1.7.8)$$

$$\Rightarrow A = \cos^{-1} \frac{2}{\sqrt{629}} \quad (1.1.7.9)$$



(b) Finding the value of angle B

$$\mathbf{C} - \mathbf{B} = \begin{pmatrix} -5 \\ 5 \end{pmatrix} \quad (1.1.7.10)$$

and

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -4 \\ -1 \end{pmatrix} \quad (1.1.7.11)$$

also calculating the values of norms

$$\|\mathbf{C} - \mathbf{B}\| = \sqrt{50} \|\mathbf{A} - \mathbf{B}\| = \sqrt{17} \quad (1.1.7.12)$$

and by doing matrix multiplication we get,

$$(\mathbf{C} - \mathbf{B})^\top (\mathbf{A} - \mathbf{B}) = \begin{pmatrix} -5 & 5 \end{pmatrix} \begin{pmatrix} -4 \\ -1 \end{pmatrix} = -25 \quad (1.1.7.13)$$

So, we get

$$\cos B = \frac{-25}{50\sqrt{17}} \quad (1.1.7.14)$$

$$= \frac{-25}{\sqrt{850}} \quad (1.1.7.15)$$

$$\Rightarrow B = \cos^{-1} \frac{-25}{\sqrt{850}} \quad (1.1.7.16)$$

(c) Finding the value of angle C

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} 1 \\ -6 \end{pmatrix} \quad (1.1.7.17)$$

and

$$\mathbf{B} - \mathbf{C} = \begin{pmatrix} 5 \\ -5 \end{pmatrix} \quad (1.1.7.18)$$

also calculating the values of norms

$$\|\mathbf{A} - \mathbf{C}\| = \sqrt{37} \quad (1.1.7.19)$$

$$\|\mathbf{B} - \mathbf{C}\| = \sqrt{50} \quad (1.1.7.20)$$

and by doing matrix multiplication we get,

$$\begin{aligned} (\mathbf{A} - \mathbf{C})^\top (\mathbf{B} - \mathbf{C}) &= \begin{pmatrix} 1 & -6 \end{pmatrix} \begin{pmatrix} 5 \\ -5 \end{pmatrix} \\ &= 35 \end{aligned} \quad (1.1.7.21)$$

so

$$\cos C = \frac{35}{37\sqrt{50}} \quad (1.1.7.22)$$

$$= \frac{35}{\sqrt{1850}} \quad (1.1.7.23)$$

$$\Rightarrow C = \cos^{-1} \frac{35}{\sqrt{1850}} \quad (1.1.7.24)$$