

$$P(a = heads)$$

$$P(a = heads) = 0.5$$


$$P(a = heads) = 0.5$$
$$= \delta(a = 0.5)$$

Probabilities are **all** distributions.





$$P(a) \geq 0$$
, for all  $a$ 



$$P(a = heads) = 0.5$$

$$P(a = tails) = 0.5$$

$$\int P(a)da = 1$$

$$\int P(a)da = 1$$

P(a) has units of 1/a

$$\int P(a|b)da = 1$$

$$P(a) =$$

$$P(a) = \int P(a|b) P(b)db$$

$$P(a) = \int P(a|b) P(b) db$$


"Marginalization"

(also "Expected Value")

$$\int P(a|b)db = 1$$

$$\int P(a|b)db = 1$$

has units of b/a



has units of b/a

$$P(a,b) = P(a)P(a|b)$$

$$P(a,b) = P(a)P(a|b)$$

$$P(b,a) = P(b)P(b|a)$$

$$P(a,b) = P(a)P(a|b)$$

$$P(b,a) = P(b)P(b|a)$$

$$P(b,a) = P(a,b)$$

$$P(a,b) = P(a)P(a|b)$$

$$P(b,a) = P(b)P(b|a)$$

$$P(b,a) = P(a,b)$$

$$P(a|b) = \frac{P(b|a)P(a)}{P(b)}$$


$$P(a,b) = P(a)P(a|b)$$


$$P(b,a) = P(b)P(b|a)$$


$$P(b,a) = P(a,b)$$


$$P(a|b) = \frac{P(b|a)P(a)}{P(b)}$$


Bayes's Theorem













P(B) is our new 'Universe'



P(B) is our new 'Universe'


2.  $P(B|A) = P(B \cap A) / P(A)$ 



P(B) is our new 'Universe'

2.  $P(B|A) = P(B \cap A) / P(A)$ 

3.  $P(B \cap A) = P(A \cap B)$ 




P(B) is our new 'Universe'

2.  $P(B|A) = P(B \cap A) / P(A)$ 

3.  $P(B \cap A) = P(A \cap B)$ 

4.  $P(A|B) = P(A \cap B) / P(B)$ = P(B|A) P(A) / P(B)



P(B) is our new 'Universe'

2.  $P(B|A) = P(B \cap A) / P(A)$ 

3.  $P(B \cap A) = P(A \cap B)$ 

4.  $P(A|B) = P(A \cap B) / P(B)$ = P(B|A) P(A) / P(B)

5. P(A|B) = P(B|A) P(A) / P(B)

# The Monty Hall Problem



(1963 - present)

#### Rules



3 doors

3 prizes — 1 behind each door (goat, hair extensions, new car)


#### Rules




You select **door 1**.

Monty opens one of the remaining doors...

#### Your door





Monty asks: stay with door 1 or switch to door 2?

# Your door

Re-phrase: is it more probable that the car is behind door 1 or door 2? Or doesn't it matter?

How does Bayes's Theorem help?

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

#### Likelihood:

1. If car behind door 1, probability of Monty choosing door 3 is 1/2.

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

- 1. If car behind door 1, probability of Monty choosing door 3 is 1/2.
- 2. If car is behind door 3, Monty has to open door 2. So probability he opens 3 and finds no car is 1.

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

- 1. If car behind door 1, probability of Monty choosing door 3 is 1/2.
- 2. If car is behind door 3, Monty has to open door 2. So probability he opens 3 and finds no car is 1.
  - 3. If car is behind door 2, Monty opens door 3 with a probability of 1, and finds no car w/ probability of 1.

- 1. You chose door 1.
- 2. Monty opened door 3 and there was no car.

- 1. If car behind door 1, probability of Monty choosing door 3 is 1/2.
- 2. If car is behind door 3, Monty has to open door 2. So probability he opens 3 and finds no car is 1.
  - 3. If car is behind door 2, Monty opens door 3 with a probability of 1, and finds no car w/ probability of 1.



I have data.

I have a model.\*

<sup>\*</sup> often based on physics

I have data.

I have a model.\*

I want to learn about my parameters of my model (and/or physics) based on my data.

\* often based on physics

# Model Fitting

I have data.

I have a model.\*

I want to learn about my parameters of my model (and/or physics) based on my data.

\* often based on physics

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

D = Data
Theta = Model parameters

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

D = Data
Theta = Model parameters

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

Probability of Theta conditioned on (or given) on Data "Posterior Probability Distribution Function" aka "The Posterior"

D = Data
Theta = Model parameters

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

Probability of Data conditioned on (or given) Theta 
"Likelihood Function" or "Data Model" 
aka 
"The Likelihood"

D = Data
Theta = Model parameters

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

Marginal or Prior Probability(ies) for Theta aka

"The Prior(s)"

D = Data
Theta = Model parameters

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

Probability of the Data or "The Evidence"

$$\frac{P(\theta|D)}{P(D|\theta)P(\theta)} = \frac{P(D|\theta)P(\theta)}{P(D)}$$
 Posterior Evidence

D = Data
Theta = Model parameters

# Updating Beliefs: Graphically

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)}$$

