

OPEN

Dilution of whisky – the molecular perspective

Björn C. G. Karlsson (5)1,3 & Ran Friedman (5)2,3

Received: 13 March 2017

Accepted: 12 June 2017

Published online: 17 August 2017

Whisky is distilled to around 70% alcohol by volume (vol-%) then diluted to about 40 vol-%, and often drunk after further slight dilution to enhance its taste. The taste of whisky is primarily associated with amphipathic molecules, such as guaiacol, but why and how dilution enhances the taste is not well understood. We carried out computer simulations of water-ethanol mixtures in the presence of guaiacol, providing atomistic details on the structure of the liquid mixture. We found that guaiacol is preferentially associated with ethanol, and, therefore, primarily found at the liquid-air interface in mixtures that contain up to 45 vol-% of ethanol. At ethanol concentrations of 59 vol-% or higher, guaiacol is increasingly surrounded by ethanol molecules and is driven to the bulk. This indicates that the taste of guaiacol in the whisky would be enhanced upon dilution prior to bottling. Our findings may apply to other flavour-giving amphipathic molecules and could contribute to optimising the production of spirits for desired tastes. Furthermore, it sheds light on the molecular structure of water-alcohol mixtures that contain small solutes, and reveals that interactions with the water may be negligible already at 89 vol-% of ethanol.

SCIENTIFIC REPORTS | 7:6489 | DOI:10.1038/s41598-017-06423-5

GIZMODO Popular Hack Used by Whisky Snobs Actually Works, Says Science TECH SCIENCE CULTURE CARS REVIEWS LONGFORM VIDEO Rae Paoletta 8/17/17 10:40am • Filed to: BOOZE NEWZ > Here's the scientific reason it's better to dilute whiskey with a little water MAXIM STYLE GEAR RIDES ENTERTAINMENT SPORTS LUXURY MAXIM MAN VIDEO MAXIM STORE MAXIM'S FINEST HERE'S THE ABSOLUTE BEST WAY TO DRINK WHISKEY,

ACCORDING TO SCIENCE Have you been doing it all wrong?

BRANDON FRIEDERICH - AUG 22, 2017