

Other Regression Models

Prof. Daniel A. Menasce
Dept. of Computer Science
George Mason University

1

© 2001. D. A. Menascé. All Rights Reserved.

Multiple Linear Regression

- Use to predict the value of the response variable as function of k predictor variables X_1, \dots, X_n .

$$\hat{Y}_i = b_0 + b_1 X_{1i} + b_2 X_{2i} + \dots + b_x X_{ki}$$

- Similar to simple linear regression.
- MS Excel can be used to do multiple linear regression.

2

© 2001. D. A. Menascé. All Rights Reserved.

CPU Time (y _i)	I/O Time (x _{1i})	Memory Requirement (x _{2i})
2	14	70
5	16	75
7	27	144
9	42	190
10	39	210
13	50	235
20	83	400

Want to find:

$$\text{CPU Time} = b_0 + b_1 * \text{I/O Time} + b_2 * \text{Memory Requirement}$$

3


© 2001. D. A. Menascé. All Rights Reserved.

SUMMARY OUTPUT

Regression Statistics

Multiple R	0.9870	R
R Square	0.9742	
Adjusted R Square	0.9614	
Standard Error	1.1511	
Observations	7	

	Coefficients	Standard Error	t Stat	Lower 95%	Upper 95%	Lower 90.0%	Upper 90.0%
Intercept (b ₀)	-0.16145	0.91345	-0.17674	-2.69759	2.37470	-2.10878	1.78589
X Variable 1 (b ₁)	0.11824	0.19260	0.61389	-0.41652	0.65299	-0.29236	0.52884
X Variable 2 (b ₂)	0.02650	0.04045	0.65519	-0.08580	0.13881	-0.05973	0.11273



4

© 2001. D. A. Menascé. All Rights Reserved.

Curvilinear Regression

- Approach: plot a scatter plot. If it does not look linear, try non-linear models:

Non-linear	Linear
$y = a + b/x$	$y = a + b(1/x)$
$y = 1/(a + bx)$	$(1/y) = a + bx$
$y = x/(a + bx)$	$(x/y) = a + bx$
$y = a \times b^x$	$\ln y = \ln a + x \ln b$
$y = a + bx^n$	$y = a + b(x^n)$