


Estimation Procedures

Daniel A. Menascé. Ph.D.
Dept. of Computer Science
George Mason University

1

© 2002 D. A. Menascé. All Rights Reserved.


Statistical Inference



2

© 2002 D. A. Menascé. All Rights Reserved.

Interval Estimate



The interval estimate of the population parameter will have a specified confidence or probability of correctly estimating the population parameter.

3

© 2002 D. A. Menascé. All Rights Reserved.

Properties of Point Estimators

- Example of point estimator: sample mean.
- Properties:
 - Unbiasedness: the expected value of all possible sample statistics (of given size n) is equal to the population parameter.
$$E[\bar{X}] = \mu$$
$$E[s^2] = \sigma^2$$
 - Efficiency: precision as estimator of the population parameter.
 - Consistency: as the sample size increases the sample statistic becomes a better estimator of the population parameter.

4

© 2002 D. A. Menascé. All Rights Reserved.

Unbiasedness of the Mean

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$$

$$E[\bar{X}] = \frac{E\left[\sum_{i=1}^n X_i\right]}{n} = \frac{\sum_{i=1}^n E[X_i]}{n} =$$

$$\frac{\sum_{i=1}^n \mu}{n} = \frac{n\mu}{n} = \mu$$

© 2002 D. A. Menascé. All Rights Reserved.

5

Sample size= 15 1.7% of population

	Sample 1	Sample 2	Sample 3			
	0.0739	0.0202	0.2918			
	0.1407	0.1089	0.4696			
	0.1257	0.0242	0.8644			
	0.0432	0.4253	0.1494			
	0.1784	0.1584	0.4242			
	0.4106	0.8948	0.0051			
	0.1514	0.0352	1.1706			
	0.4542	0.1752	0.0084			
	0.0485	0.3287	0.0600			
	0.1705	0.1697	0.7820			
	0.3335	0.0920	0.4985			
	0.1772	0.1488	0.0988			
	0.0242	0.2486	0.4896			
	0.2183	0.4627	0.1892			
	0.0274	0.4079	0.1142			
	E[sample]			Population Error		
Sample Average	0.1718	0.2467	0.3744	0.2643	0.2083	26.9%
Sample Variance	0.0180	0.0534	0.1204	0.0639	0.0440	45.3%
Efficiency (average)	18%	18%	80%			
Efficiency (variance)	59%	21%	173%			

© 2002 D. A. Menascé. All Rights Reserved.

6

Sample size = 87

10% of population

Sample 1	Sample 2	Sample 3
0.5725	0.3864	0.4627
0.0701	0.0488	0.2317
0.2165	0.0611	0.1138
0.6581	0.0881	0.0047
0.0440	0.5866	0.2438
0.1777	0.3419	0.0819
0.2380	0.1923	0.6581

	0.0102	0.9460	0.0714			
	0.4325	0.0445	0.2959		Population	% Rel. Error
Sample Average	0.2239	0.2203	0.2178	0.2206	0.2083	5.9%
Sample Variance	0.0452688	0.0484057	0.0440444	0.0459	0.0440	4.3%
Efficiency (average)	7.5%	5.7%	4.5%			
Efficiency (variance)	2.9%	10.0%	0.1%			

7

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval Estimation of the Mean

- Known population standard deviation.
- Unknown population standard deviation:
 - Large samples: sample standard deviation is a good estimate for population standard deviation. OK to use normal distribution.
 - Small samples and original variable is normally distributed: use t distribution with $n-1$ degrees of freedom.

8

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval Estimation of the Mean

$$\Pr[c_1 \leq \mu \leq c_2] = 1 - \alpha$$

(c_1, c_2) : confidence interval


α : significance level (e.g., 0.05)

$1 - \alpha$: confidence coefficient (e.g., 0.95)

$100(1 - \alpha)$: confidence level (e.g., 95%)

9

© 2002 D. A. Menascé. All Rights Reserved.



Sample		Interval include μ ?
1		YES
2		YES
3		NO
...		
100		YES

$100(1 - \alpha)$ of the 100 samples include the population mean μ .

10

© 2002 D. A. Menascé. All Rights Reserved.

Central Limit Theorem


- If the observations in a sample are independent and come from the same population that has mean μ and standard deviation σ then the sample mean for **large** samples has a normal distribution with mean μ and standard deviation σ/\sqrt{n} .

$$\bar{x} \sim N(\mu, \sigma / \sqrt{n})$$

- The standard deviation of the sample mean is called the *standard error*.

11

© 2002 D. A. Menascé. All Rights Reserved.



Confidence Interval (large (n>30) samples)

- 100 (1-α)% confidence interval for the population mean:

$$\left(\bar{x} - z_{1-\alpha/2} \frac{s}{\sqrt{n}}, \bar{x} + z_{1-\alpha/2} \frac{s}{\sqrt{n}} \right)$$

\bar{x} : sample mean

s: sample standard deviation

n: sample size

$z_{1-\alpha/2}$: (1-α/2)-quantile of a unit normal variate (N(0,1)).

13

© 2002 D. A. Menascé. All Rights Reserved.

	0.4325	0.0445	0.2959		Population
Sample Average	0.2239	0.2203	0.2178	0.2206	0.2083
Sample Variance	0.0452688	0.0484057	0.0440444	0.0459	0.0440
Efficiency (average)	7.5%	5.7%	4.5%		
Efficiency (variance)	2.9%	10.0%	0.1%		
95% interval lower	0.1792	0.1740	0.1737		
95% interval upper	0.2686	0.2665	0.2619	0.0894	
Mean in interval	YES	YES	YES		
99% interval lower	0.1651	0.1595	0.1598		
99% interval upper	0.2826	0.2810	0.2757	0.1175	
Mean in interval	YES	YES	YES		
90% interval lower	0.1864	0.1815	0.1807		
90% interval upper	0.2614	0.2591	0.2548	0.0750	
Mean in interval	YES	YES	YES		

In Excel:

½ interval = CONFIDENCE(1-0.95,s,n)

α

interval size

14

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval (small samples, normally distributed population)

- 100 (1- α)% confidence interval for the population mean:

$$\left(\bar{x} - t_{[1-\alpha/2; n-1]} \frac{s}{\sqrt{n}}, \bar{x} + t_{[1-\alpha/2; n-1]} \frac{s}{\sqrt{n}} \right)$$

\bar{x} : sample mean

s: sample standard deviation

n: sample size

$t_{[1-\alpha/2; n-1]}$: critical value of the t distribution with $n-1$ degrees of freedom for an area of $\alpha/2$ for the upper tail.

15

© 2002 D. A. Menascé. All Rights Reserved.

Student's t distribution

$$t(\nu) \sim \frac{N(0,1)}{\sqrt{\chi^2(\nu)/\nu}}$$

ν : number of degree of freedom.

$\chi^2(\nu)$: chi-square distribution with ν degrees of freedom. Equal to the sum of squares of ν unit normal variates.

- the pdf of a t-variate is similar to that of a $N(0,1)$.
- for $\nu > 30$ a t distribution can be approximated by $N(0,1)$.

16

© 2002 D. A. Menascé. All Rights Reserved.

Using the t Distribution. Sample size= 15.

	0.0274	0.4079	0.1142	E[sample]	Population	Error
Sample Average	0.1718	0.2467	0.3744	0.2643	0.2083	26.9%
Sample Variance	0.0180	0.0534	0.1204	0.0639	0.0440	45.3%
Efficiency (average)	18%	18%	80%			
Efficiency (variance)	59%	21%	173%			
95% interval lower	0.0975	0.1187	0.1823			
95% interval upper	0.2462	0.3747	0.5665			
Mean in interval	YES	YES	YES			

95%, n-1
critical value

2.145

In Excel: TINV(1-0.95,15-1)

α

© 2002 D. A. Menascé. All Rights Reserved.

17

Confidence Interval for the Variance

- If the original variable is normally distributed then the chi-square distribution can be used to develop a confidence interval estimate of the population variance.
- The $(1-\alpha)\%$ confidence interval for σ^2 is

$$\frac{(n-1)s^2}{\chi_U^2} \leq \sigma^2 \leq \frac{(n-1)s^2}{\chi_L^2}$$


χ_L^2 : lower critical value of χ^2

χ_U^2 : upper critical value of χ^2

© 2002 D. A. Menascé. All Rights Reserved.

18

Chi-square distribution



19

© 2002 D. A. Menascé. All Rights Reserved.

95% confidence interval for the population variance
for a sample of size 100 for a $N(3,2)$ population.

2.91903	average
4.71435	variance
2.17126	std deviation

73.36110 lower critical value of chi-square for 95%

128.42193 upper critical value of chi-square for 95%

lower bound for confidence interval for the variance 3.634277

upper bound for confidence interval for the variance 6.361966

1- $\alpha/2$
↓
In Excel: CHIINV (0.975, 99)
↑
CHIINV (0.025, 99)
↑
 $\alpha/2$

The population variance (4 in this case) is in the interval
(3.6343, 6.362) with 95% confidence.

20

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval for the Variance

If the population is not normally distributed, the confidence interval, especially for small samples, is not very accurate.

21

© 2002 D. A. Menascé. All Rights Reserved.

Prediction Interval for a Future Value

- Interval in which a future value will lie with a degree of confidence.

$$\bar{X} - t_{[1-\alpha/2; n-1]} s \sqrt{1 + 1/n} \leq X_f \leq \bar{X} + t_{[1-\alpha/2; n-1]} s \sqrt{1 + 1/n}$$

$(1-\alpha/2)$ -quantile of t -variate with $n-1$ degrees of freedom.

3.0028	average
2.1487	std deviation
t [1-0.05/2;24]	2.0639
Lower bound	-1.5197
Upper bound	7.5254

In Excel: TINV(α ,24)

A future value will lie in the interval (-1.519,7.525) with 95% confidence.

22

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval for Proportions

- For categorical data:
 - E.g. file types
 $\{\text{html}, \text{html}, \text{gif}, \text{jpg}, \text{html}, \text{pdf}, \text{ps}, \text{html}, \text{pdf} \dots\}$
 - If n_I of n observations are of type html, then the sample proportion of html files is $p = n_I/n$.
- The population proportion is π .
- Goal: provide confidence interval for the population proportion π .

23

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval for Proportions

- The sampling distribution of the proportion formed by computing p from all possible samples of size n from a population of size N with replacement tends to a normal with mean π and standard error $\sigma_p = \sqrt{\frac{\pi(1-\pi)}{n}}$.
- The normal distribution is being used to approximate the binomial. So, $n\pi \geq 10$.

24

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval for Proportions

- The $(1-\alpha)\%$ confidence interval for π is

$$\left(p - z_{1-\alpha/2} \sqrt{\frac{p(1-p)}{n}}, p + z_{1-\alpha/2} \sqrt{\frac{p(1-p)}{n}} \right)$$

p: sample proportion.

n: sample size

$z_{1-\alpha/2}$: $(1-\alpha/2)$ -quantile of a unit normal variate ($N(0,1)$).

25

© 2002 D. A. Menascé. All Rights Reserved.

Confidence Interval for Proportions

- One thousand entries are selected from a Web log. Six hundred and fifty correspond to gif files. Find 90% and 95% confidence intervals for the proportion of files that are gif files.

Sample size (n)	1000
No. gif files in sample	650
Sample proportion (p)	0.65
$n \cdot p$	650

> 10 OK

90% confidence interval

alpha	0.1
$1-\alpha/2$	0.95
$z_{0.95}$	1.645
Lower bound	0.625
Upper bound	0.675

In Excel:

NORMSINV(1-0.1/2)

95% confidence interval

alpha	0.05
$1-\alpha/2$	0.975
$z_{0.975}$	1.960
Lower bound	0.620
Upper bound	0.680

NORMSINV(1-0.05/2)

26

© 2002 D. A. Menascé. All Rights Reserved.


Comparing Alternatives

- Suppose you want to compare two cache replacement policies under similar workloads.
- Metric of interest: cache hit ratio.
- Types of comparisons:
 - Paired observations
 - Unpaired observations.

27

© 2002 D. A. Menascé. All Rights Reserved.

Paired Observations



28

© 2002 D. A. Menascé. All Rights Reserved.

Example of Paired Observations

- Six similar workloads were used to compare the cache hit ratio obtained under object replacement policies A and B on a Web server. Is A better than B?

Workload	Cache Hit Ratio		A-B
	Policy A	Policy B	
1	0.35	0.28	0.07
2	0.46	0.37	0.09
3	0.29	0.34	-0.05
4	0.54	0.60	-0.06
5	0.32	0.22	0.10
6	0.15	0.18	-0.03
Sample mean			0.02000
Sample variance			0.00552
Sample standard dev.			0.07430

29

© 2002 D. A. Menascé. All Rights Reserved.

Example of Paired Observations

Sample mean	0.02000
Sample variance	0.00552
Sample standard dev.	0.07430

In Excel:
TINV(1-0.9,5)

0.95 quantile of t-variable with 5 degrees of freedom

90% confidence interval

lower bound

upper bound

2.015

-0.0411

0.0811

$$\left(\bar{x} - t_{[1-\alpha/2;n-1]} \frac{s}{\sqrt{n}}, \bar{x} + t_{[1-\alpha/2;n-1]} \frac{s}{\sqrt{n}} \right)$$

Diagram illustrating the components of the confidence interval formula:

- \bar{x} (Sample mean) is 0.02
- $t_{[1-\alpha/2;n-1]}$ (t-value) is 2.015
- s (Sample standard deviation) is 0.0743
- n (Sample size) is 6

30

© 2002 D. A. Menascé. All Rights Reserved.

Example of Paired Observations

Sample mean	0.02000
Sample variance	0.00552
Sample standard dev.	0.07430

In Excel:
TINV(1-0.9,5)

0.95 quantile of t-variable with 5 degrees of freedom

90% confidence interval

lower bound

upper bound

2.015

-0.0411


0.0811

The interval includes zero, so we cannot say that policy A is better than policy B.

31

© 2002 D. A. Menascé. All Rights Reserved.

Unpaired Observations



32

© 2002 D. A. Menascé. All Rights Reserved.

Unpaired Observations (t-test)

1. Size of samples for A and B: n_A and n_B
2. Compute sample means:

$$\bar{x}_A = \frac{1}{n_A} \sum_{i=1}^{n_A} x_{iA}$$

$$\bar{x}_B = \frac{1}{n_B} \sum_{i=1}^{n_B} x_{iB}$$

33

© 2002 D. A. Menascé. All Rights Reserved.

Unpaired Observations (t-test)

3. Compute the sample standard deviations:

$$s_A = \sqrt{\frac{\left(\sum_{i=1}^{n_A} x_{iA}^2 \right) - n_A (\bar{x}_A)^2}{n_A - 1}}$$

$$s_B = \sqrt{\frac{\left(\sum_{i=1}^{n_B} x_{iB}^2 \right) - n_B (\bar{x}_B)^2}{n_B - 1}}$$

34

© 2002 D. A. Menascé. All Rights Reserved.

Unpaired Observations (t-test)

4. Compute the mean difference: $\bar{x}_a - \bar{x}_b$
5. Compute the standard deviation of the mean difference:
6. Compute the effective number of degrees of freedom.

$$s = \sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}$$

$$v = \frac{(s_a^2/n_a + s_b^2/n_b)^2}{\frac{1}{n_a - 1} \left(\frac{s_a^2}{n_a}\right)^2 + \frac{1}{n_b - 1} \left(\frac{s_b^2}{n_b}\right)^2} - 2$$

35

© 2002 D. A. Menascé. All Rights Reserved.

Unpaired Observations (t-test)

7. Compute the confidence interval for the mean difference:

$$(\bar{x}_a - \bar{x}_b) \pm t_{[1-\alpha/2;v]} \times s$$

8. If the confidence interval includes zero, the difference is not significant at 100(1- α)% confidence level.

36

© 2002 D. A. Menascé. All Rights Reserved.

Example of Unpaired Observations

- Two cache replacement policies A and B are compared under similar workloads. Is A better than B?

Workload	Cache Hit Ratio	
	Policy A	Policy B
1	0.35	0.49
2	0.23	0.33
3	0.29	0.33
4	0.21	0.55
5	0.21	0.65
6	0.15	0.18
7	0.42	0.29
8		0.35
9		0.44
Mean	0.2657	0.4011
St. Dev	0.0934	0.1447

37

© 2002 D. A. Menascé. All Rights Reserved.

Example of Unpaired Observations

na	7
nb	9
mean diff	-0.135
st. dev. Diff	0.059776
Eff. Degr. Freedom	12
alpha	0.1
1-alpha/2	0.95
t[1-alpha/2,v]	1.796
90% confidence interval	
lower bound	-0.243
upper bound	-0.028


← In Excel: TINV(1-0.9,12)

At a 90% confidence level the two policies are not identical since zero is not in the interval. With 90% confidence, the cache hit ratio for policy A is smaller than that for policy B. So, policy B is better at that confidence level.

38

© 2002 D. A. Menascé. All Rights Reserved.

Approximate Visual Test



CIs do not overlap:
A is higher than B

CIs overlap and mean
of A is in B's CI:
A and B are similar

CIs overlap and mean
of A is not in B's CI:
need to do t-test

39

© 2002 D. A. Menascé. All Rights Reserved.

Example of Visual Test

Workload	Cache Hit Ratio	
	Policy A	Policy B
1	0.35	0.49
2	0.23	0.33
3	0.29	0.33
4	0.21	0.55
5	0.21	0.65
6	0.15	0.18
7	0.42	0.29
8		0.35
9		0.44
Mean	0.2657	0.4011
St. Dev	0.0934	0.1447

na 7
 nb 9
 alpha 0.1 for
 1-alpha/2 0.95
 Policy A Policy B
 t[1-alpha/2,v] 1.9432 1.8595
 90% Confidence Interval
 lower bound 0.197 0.311
 upper bound 0.334 0.491

90% confidence interval

CIs overlap but mean of A is
 not in CI of B and vice-versa.
 Need to do a t-test.

40

© 2002 D. A. Menascé. All Rights Reserved.

One-sided Confidence Intervals



- Useful to test the hypothesis that the mean is greater (or smaller) than a certain value.

$$\Pr[\mu \geq c_1] = 1 - \alpha$$

$$\Pr[\mu \leq c_2] = 1 - \alpha$$

41

© 2002 D. A. Menascé. All Rights Reserved.



In general:

$$c_1 = \bar{x} - \frac{s}{\sqrt{n}} t_{[1-\alpha; n-1]}$$

42

© 2002 D. A. Menascé. All Rights Reserved.

One-sided Confidence Intervals

$$\begin{aligned} &(-\infty, \bar{x} + t_{[1-\alpha; n-1]} s / \sqrt{n}) \\ &(\bar{x} - t_{[1-\alpha; n-1]} s / \sqrt{n}, \infty) \end{aligned}$$

43

© 2002 D. A. Menascé. All Rights Reserved.

Determining Sample Size

- Large samples imply high confidence.
- Large samples require more data collection effort.
- How to determine the sample size n to estimate the population parameter with accuracy $r\%$ and confidence level of $100(1-\alpha)\%$?

44

© 2002 D. A. Menascé. All Rights Reserved.

Determining the Sample Size for the Mean

- Perform a set of measurements to estimate the sample mean and the sample variance.
- Determine the sample size to obtain proper accuracy as follows:

$$\bar{x} \pm z \frac{s}{\sqrt{n}} = \bar{x} \pm \frac{\bar{x}r}{100}$$

$$\Rightarrow n = \left(\frac{100zs}{r\bar{x}} \right)^2$$

45

© 2002 D. A. Menascé. All Rights Reserved.

Determining the Sample Size for the Mean

- A preliminary test shows that the sample mean of the response time is 5 sec and the sample standard deviation is 1.5. How many repetitions are needed to get the response time within 2% accuracy at 95% confidence level?

$$r = 2 \quad \bar{x} = 5 \quad s = 1.5$$

$$z = 1.96$$

865 repetitions would be Needed!


$$n = \left(\frac{100 \times 1.96 \times 1.5}{2 \times 5} \right)^2 = 864.36$$

46

© 2002 D. A. Menascé. All Rights Reserved.

Determining the Sample Size for the Mean

Accuracy (r)	Confidence Level (1-alpha)	X	S	Sample size
1	0.95	5	0.8	984
2	0.95	5	0.8	246
5	0.95	5	0.8	40
1	0.9	5	0.8	693
2	0.9	5	0.8	174
5	0.9	5	0.8	28



47

© 2002 D. A. Menascé. All Rights Reserved.

Computing Important Quantiles in Excel

$z_{1-\alpha/2}$ = (1- $\alpha/2$)-quantile of a unit normal variate (N(0,1)):
 = NORMINV (1- $\alpha/2$,0,1) = NORMSINV(1- $\alpha/2$)
 Half-interval = CONFIDENCE (α , σ ,n)

$t_{[1-\alpha/2;n-1]}$ = (1- $\alpha/2$)-quantile of t -variate with $n-1$ degrees of freedom = TINV(α ,n-1)

χ_L^2 : lower critical value of χ^2 = CHIINV (1- $\alpha/2$,n-1)
 χ_U^2 : upper critical value of χ^2 = CHIINV ($\alpha/2$, n-1)

48

© 2002 D. A. Menascé. All Rights Reserved.