

Summarizing Data

Daniel A. Menascé, Ph.D.
Dept of Computer Science
George Mason University


Major Properties of Numerical Data

- Central Tendency: arithmetic mean, geometric mean, median, mode.
- Variability: range, interquartile range, variance, standard deviation, coefficient of variation, mean absolute deviation.
- Skewness: coefficient of skewness.
- Kurtosis

Measures of Central Tendency

- Arithmetic Mean

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$$


- Based on all observations  greatly affected by extreme values.

© 2001 D. A. Menascé. All Rights Reserved.

3

Effect of Outliers on Average

	1.1	1.1
	1.4	1.4
	1.8	1.8
	1.9	1.9
	2.3	2.3
	2.4	2.4
	2.8	2.8
	3.1	3.1
	3.4	3.4
	3.8	3.8
	10.3	3.5
Average	3.1	2.5



© 2001 D. A. Menascé. All Rights Reserved.

4

Geometric Mean

- Geometric Mean: $\left(\prod_{i=1}^n X_i \right)^{1/n}$
- Used when the product of the observations is of interest.
- Important when multiplicative effects are at play:
 - Cache hit ratios at several levels of cache
 - Percentage performance improvements between successive versions.
 - Performance improvements across protocol layers.

© 2001 D. A. Menascé. All Rights Reserved.

5

Example of Geometric Mean

Test Number	Performance Improvement			Avg. Performance Improvement per Layer
	Operating System	Middleware	Application	
1	1.18	1.23	1.10	1.17
2	1.25	1.19	1.25	1.23
3	1.20	1.12	1.20	1.17
4	1.21	1.18	1.12	1.17
5	1.30	1.23	1.15	1.23
6	1.24	1.17	1.21	1.21
7	1.22	1.18	1.14	1.18
8	1.29	1.19	1.13	1.20
9	1.30	1.21	1.15	1.22
10	1.22	1.15	1.18	1.18
Average Performance Improvement per Layer				1.20

© 2001 D. A. Menascé. All Rights Reserved.

6

Properties of the Geometric Mean

$$gm\left(\frac{x_1}{y_1}, \dots, \frac{x_n}{y_n}\right) = \frac{gm(x_1, \dots, x_n)}{gm(y_1, \dots, y_n)} = \frac{1}{gm(y_1/x_1, \dots, y_n/x_n)}$$

- The choice of the base does not change the conclusion.
- Useful for benchmarks
 - x: throughput on target system.
 - y: throughput on base system.

© 2001 D. A. Menascé. All Rights Reserved.

7

Median

- Middle Value in an Ordered Set of Data.
- If there are no ties, 50% of the values are smaller than the median and 50% are larger.

	1.1	1.1
	1.4	1.4
	1.8	1.8
	1.9	1.9
	2.3	2.3
	2.4	2.4
	2.8	2.8
	3.1	3.1
	3.4	3.4
	3.8	3.8
	10.3	3.5
Median	2.4	2.4

© 2001 D. A. Menascé. All Rights Reserved.


8

Median


- The median is unaffected by extreme values.
- Obtaining the median:
 - Odd-sized samples: $X_{(n+1)/2}$
 - Even-sized samples: $\frac{X_{n/2} + X_{(n/2)+1}}{2}$

Mode

- Most frequently occurring value.
- Mode may not exist.
- Single mode distributions: unimodal.
- Distributions with two modes: bimodal.



unimodal



bimodal

Quantiles (quartiles, percentiles) and midhinge

- Quartiles: split the data into quarters.
 - First quartile (Q1): value of X_i such that 25% of the observations are smaller than X_i .
 - Second quartile (Q2): value of X_i such that 50% of the observations are smaller than X_i .
 - Third quartile (Q3): value of X_i such that 75% of the observations are smaller than X_i .
- Percentiles: split the data into hundredths.
- Midhinge: $Midhinge = \frac{Q_3 + Q_1}{2}$

© 2001 D. A. Menascé. All Rights Reserved.

11

Example of Quartiles

1.05
1.06
1.09
1.19
1.21
1.28
1.34
1.34
1.77
1.80
1.83
2.15
2.21
2.27
2.61
2.67
2.77
2.83
3.51
3.77
5.76
5.78
32.07
144.91

Q1	1.32
Q2	2.18
Q3	3.00
Midhinge	2.16

In Excel:

Q1=PERCENTILE(<array>,0.25)

Q2=PERCENTILE(<array>,0.5)

Q3=PERCENTILE(<array>,0.75)

© 2001 D. A. Menascé. All Rights Reserved.

12

Example of Percentile

1.05
1.06
1.09
1.19
1.21
1.28
1.34
1.34
1.77
1.80
1.83
2.15
2.21
2.27
2.61
2.67
2.77
2.83
3.51
3.77
5.76
5.78
32.07
144.91

80-percentile 3.613002

In Excel:

p-th percentile=PERCENTILE(<array>,p)
($0 \leq p \leq 1$)

© 2001 D. A. Menascé. All Rights Reserved.

13

Range, Interquartile Range, Variance, and Standard Deviation

- Range: $X_{\max} - X_{\min}$
- Interquartile Range: $Q_3 - Q_1$
– not affected by extreme values.

- Variance:

$$s^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}$$

In Excel:

$s^2 = \text{VAR}(<\text{array}>)$

- Standard Deviation:

In Excel:

$s = \text{STDEV}(<\text{array}>)$

$$s = \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}}$$

© 2001 D. A. Menascé. All Rights Reserved.

14

Meanings of the Variance and Standard Deviation

- The larger the spread of the data around the mean, the larger the variance and standard deviation.
- If all observations are the same, the variance and standard deviation are zero.
- The variance and standard deviation cannot be negative.
- Variance is measured in the square of the units of the data.
- Standard deviation is measured in the same units as the data.

© 2001 D. A. Menascé. All Rights Reserved.

15

Coefficient of Variation

- Coefficient of variation (COV) : s / \bar{X}

– no units

1.05
1.06
1.09
1.19
1.21
1.28
1.34
1.34
1.77
1.80
1.83
2.15
2.21
2.27
2.61
2.67
2.77
2.83
3.51
3.77
5.76
5.78
32.07
144.91

S	29.50
Average	9.51
COV	3.10

© 2001 D. A. Menascé. All Rights Reserved.

16

Coefficient of Skewness

- Coefficient of skewness: $\frac{1}{ns^3} \sum_{i=1}^n (X_i - \bar{X})^3$

	(X-Xi)^3
1.05	-606.1
1.06	-602.9
1.09	-596.1
1.19	-575.2
1.21	-571.8
1.28	-557.9
1.34	-546.4
1.34	-544.8
1.77	-464.5
1.80	-458.1
1.83	-453.1
2.15	-398.9
2.21	-388.8
2.27	-379.0
2.61	-328.5
2.67	-320.5
2.77	-306.6
2.83	-298.7
3.51	-215.9
3.77	-189.6
5.76	-52.9
5.78	-52.1
32.07	11476.6
144.91	2482007.1

4.033

© 2001 D. A. Menascé. All Rights Reserved.

17

Mean Absolute Deviation

- Mean absolute deviation: $\frac{1}{n} \sum_{i=1}^n |X_i - \bar{X}|$


	abs(Xi-Xbar)
1.05	8.46
1.06	8.45
1.09	8.42
1.19	8.32
1.21	8.30
1.28	8.23
1.34	8.18
1.34	8.17
1.77	7.74
1.80	7.71
1.83	7.68
2.15	7.36
2.21	7.30
2.27	7.24
2.61	6.90
2.67	6.84
2.77	6.74
2.83	6.68
3.51	6.00
3.77	5.74
5.76	3.75
5.78	3.73
32.07	22.56
144.91	135.39
	315.90

Average	9.51
Mean absolute deviation	13.16


© 2001 D. A. Menascé. All Rights Reserved.

18


Shapes of Distributions



Right-skewed distribution



Symmetric distribution



Left-skewed distribution

© 2001 D. A. Menascé. All Rights Reserved.

19

Confidence Interval for the Mean

- The sample mean is an estimate of the population mean.
- Problem: given k samples of the population (with k sample means), get a single estimate of the population mean.
- Only probabilistic statements can be made:

© 2001 D. A. Menascé. All Rights Reserved.

20

Confidence Interval for the Mean

$$\Pr[c_1 \leq \mu \leq c_2] = 1 - \alpha$$

where,

(c_1, c_2) : confidence interval

$100(1 - \alpha)$: confidence level (usually 90 or 95%)

$1 - \alpha$: confidence coefficient.


Central Limit Theorem

- If the observations in a sample are independent and come from the same population that has mean μ and standard deviation σ then the sample mean for large samples has a normal distribution with mean μ and standard deviation σ/\sqrt{n} .
- The standard deviation of the sample mean is called the *standard error*.

Central Limit Theorem

Population mean = μ

Population std deviation = σ



Average of $x_1, \dots, x_M = \mu$

Standard deviation of $x_1, \dots, x_M = \sigma / \sqrt{n}$

© 2001 D. A. Menascé. All Rights Reserved.

23

Confidence Interval

- 100 (1- α)% confidence interval for the population mean:

$$(\bar{x} - z_{1-\alpha/2} s / \sqrt{n}, \bar{x} + z_{1-\alpha/2} s / \sqrt{n})$$

\bar{x} : sample mean

s: sample standard deviation

n: sample size

$z_{1-\alpha/2}$: (1- $\alpha/2$)-quantile of a unit normal variate (N(0,1)).

© 2001 D. A. Menascé. All Rights Reserved.

24

Example of Confidence Interval Computation

CPU Time (msec)
5.76
2.67
3.77
2.27
2.83
1.05
2.61
1.06
5.78
3.51
2.77
1.83
1.77
1.19
2.21
24.80
1.80
1.34
1.28
1.21
2.15
1.09
1.34
32.07

n	24
sample mean	4.51
sample std	7.56
alpha	0.1
conf level	90
1-(alpha/2)	0.95
z0.95	1.645
c1	1.97
c2	7.04

from a Normal Table

With 90% confidence the population mean is in the interval 1.97 7.04

© 2001 D. A. Menascé. All Rights Reserved.

25

From Excel: Tools > Data Analysis > Descriptive Statistics

Descriptive Statistics (from Excel Analysis Pack)

Mean	9.510589
Standard Error	6.021322
Median	2.180555
Mode	#N/A
Standard Deviation	29.49833
Sample Variance	870.1515
Kurtosis	21.65021
Skewness	4.59114
Range	143.8572
Minimum	1.047923
Maximum	144.9051
Sum	228.2541
Count	24
Confidence Level(95.0%)	12.45604

$$\leftarrow \frac{s}{\sqrt{n}}$$

© 2001 D. A. Menascé. All Rights Reserved.

26

Box-and-Whisker Plot

- Graphical representation of data through a five-number summary.

I/O Time (msec)
8.04
9.96
5.68
6.95
8.81
10.84
4.26
4.82
8.33
7.58
7.24
7.46
8.84
5.73
6.77
7.11
8.15
5.39
6.42
7.81
12.74
6.08

Five-number Summary	
Minimum	4.26
First Quartile	6.08
Median	7.35
Third Quartile	8.33
Maximum	12.74

