Equations of State for Realistic Water Models

1 Introduction

Water molecules are remarkably simple — consisting only of two hydrogen atoms and one oxygen atom — yet liquid water is highly complex, with well-known thermodynamic anomalies in its density,[1–4] heat capacity,[5–8] thermal conductivity,[9–11] diffusivity,[12–15] and surface tension,[16–19] amongst other properties. Many of these anomalies occur in either the supercooled, high-pressure, or negative-pressure regimes, which are experimentally difficult to access, yet have direct technological applications and consequences.[20–23] Nanoconfinement only amps up liquid water's bizarre array of oddities, such as strongly altered dielectric properties,[24–26] reduced friction,[27,28] interfacial-modulated reactivity,[29] and a potential superionic state.[30,31]

Molecular dynamics (MD) has become an increasingly popular technique to study liquid water systems, especially under nanoconfinement or at interfaces, for nanotechnological applications.[32–35] MD simulations are also naturally able to access experimentally-challenging regimes, such as deeply-supercooled liquid states and exotic high-pressure ices, without special difficulty.[36–39] In particular, the rise of machine learning potentials (MLPs) trained on *ab initio* reference data such as DFT allow highly accurate and realistic modelling of homogenous and heterogenous systems from first principles, capturing effects such as chemical reactivity and many-body interactivity without needing *post-hoc* adjustments.[40–45] MLP-driven MD simulations promise to accelerate materials design for technological applications involving water, by enabling key insights into water's behaviours at the nanoscale[46–48] and under extreme conditions.[4,30] Yet, conclusions drawn from MD simulations of liquid water, whether done through MLPs, computationally cheaper force fields (FFs), or even directly at an *ab initio* level, can only be as accurate as the underlying model.[49–53] This raises the critical question: how can we benchmark the performance and realism of different water models, especially across the extreme supercooled and high-pressure regimes?

In principle, the properties of liquid water under a given model can be measured using MD simulations across a wide range of thermodynamic conditions, and compared against experimental data pointwise. This is already a common practice, for example, when parametrizing FFs.[54–57] However, such a 'pointwise' approach is computationally wasteful: each state point requires a fresh simulation, and the resulting comparisons offer no general method of interpolating between sampled conditions, nor extrapolating towards extreme regimes. A more powerful

approach is to reconstruct the full Equation of State (EoS) of a given water model directly from a comparatively sparse set of simulations, by measuring fundamental thermodynamic quantities — such as the pressure and internal energy — as functions of volume and temperature. These can then be used to build a continuous representation of the model's free energy surface, allowing further properties of the model, at any state point within the domain of validity, to be obtained by direct evaluation without further simulation.

Our proposed approach offers two distinct advantages. Firstly, it allows the "realism" of a given water model to be assessed efficiently and comprehensively, by comparing its reconstructed EoS against experimental reference data across the full thermodynamic plane, rather than at a handful of isolated conditions. Secondly, and more importantly, it provides a natural framework for comparing models of foundationally different character — from simple, computationally cheap FFs such as SPC/E and TIP4P, to MLPs trained on high-accuracy DFT or coupled-cluster reference data — on equal footing, since the free energy surface construction makes no assumptions about the underlying interaction model, beyond requiring access to pressures and internal energies. This is particularly valuable in the supercooled and high-pressure regimes, where experimental measurements are difficult to obtain, but where accurate modelling is most needed to resolve long-standing questions about the microscopic origins of water's thermodynamic anomalies.

2 Theory

2.1 The Equation of State (EoS) in classical thermodynamics

Consider the canonical ensemble, i.e. a system of constant particle number N and volume V freely exchanging energy with a thermal reservoir of temperature T. The partition function is defined as:

$$Z(N,V,T) \;:=\; \frac{1}{{\mathcal{N}_0}^N}\int_{\boldsymbol{\omega}\in\Omega_N(V)} \exp\left[-\frac{1}{k_B T}\mathcal{H}(\boldsymbol{\omega})\right]\,\mathrm{d}^{6N}\boldsymbol{\omega} \tag{1}$$

where k_B is the Boltzmann constant, $\Omega_N(V)$ is the set of all available microstates at the given volume, $\mathcal{H}(\boldsymbol{\omega})$ is the Hamiltonian at microstate $\boldsymbol{\omega}$, and $\mathcal{N}_0$ is an arbitrary normalization constant which nondimensionalizes the expression. In general, the partition function is a smooth analytic function of V and T, except at phase transitions. The Helmholtz free energy is then defined as:

$$F(N,V,T) \;:=\; -k_B T \ln Z(N,V,T)\,. \tag{2}$$

Suppose there is some thermodynamic observable $A(\boldsymbol{\omega})$ which is a function of the microstate, and a corresponding conjugate variable λ such that $\partial_\lambda \mathcal{H} = A$. The thermal expectation of the observable is given by the Boltzmann distribution:

$$\langle A \rangle_{NVT} \quad = \quad \frac{\int_{\boldsymbol{\omega} \in \Omega_N(V)} A(\boldsymbol{\omega}) \exp\left[-\frac{1}{k_B T} \mathcal{H}(\boldsymbol{\omega})\right] \, \mathrm{d}^{6N}\boldsymbol{\omega}}{\int_{\boldsymbol{\omega} \in \Omega_N(V)} \exp\left[-\frac{1}{k_B T} \mathcal{H}(\boldsymbol{\omega})\right] \, \mathrm{d}^{6N}\boldsymbol{\omega}} \tag{3}$$

which, upon substituting eqs. (1) and (2), can be expressed in terms of the partition function and Helmholtz free energy respectively:

$$\langle A \rangle_{NVT} \quad = \quad -\frac{k_B T}{Z} \frac{\partial Z}{\partial \lambda} \quad = \quad \frac{\partial F}{\partial \lambda} \, . \tag{4}$$

As such, knowledge of the exact analytical form of the Helmholtz free energy $F(N, V, T)$ as a function of volume, temperature, and other parameters of interest λ is sufficient to derive all macroscopic equilibrium properties $\langle A \rangle_{NVT}$ in the canonical ensemble. Furthermore, by inverting the relevant variables, equilibrium properties in other ensembles can also be obtained; for example, the isobaric expectation $\langle A \rangle_{NpT}$ can be extracted by evaluating $\langle A \rangle_{NVT}$ at the inverse $V = V(p)$ of the $p(V)$ function. For this reason, especially where a single continuous thermodynamic phase of a given substance is concerned, the functional form of $F(N, V, T)$ is often known as the *Equation of State*.

2.2 EoSs for bulk phases

Now let us focus only on bulk condensed-phase substances, e.g. liquid water. In the thermodynamic limit $N \to \infty$, interfacial and edge effects vanish, and statistical fluctuations of bulk observables are suppressed relative to their expectation values. As such, all thermodynamic variables are either *extensive* and scale proportionally to N, or *intensive* and scale independently from N. It is useful to work only with intensive quantities; this can be achieved by dividing all extensive properties by the particle number N, or in other words we consider only quantities like the *volume per particle* or *internal energy per particle*, which for this subsection we mark with a circumflex:

$$\hat{V} \; := \; \frac{V}{N} \qquad ; \qquad \hat{U} \; := \; \frac{U}{N} \qquad ; \qquad \text{etc.} \tag{5}$$

In this chapter, we use the convention that each "particle" of liquid water refers to the entire water molecule, i.e. three atoms, or hydronium or hydroxide ions. This is a sensible choice, as the energy scale of the chemical bindings holding water molecules together far exceeds the thermodynamic energy scales which determine liquid water's bulk properties,[58–61] but any other convention would not affect results beyond a numerical factor.

Most importantly, we have the *free energy per particle*:

$$\hat{F}(\hat{V}, T) \quad := \quad \frac{F(N, V, T)}{N} \quad = \quad -\frac{k_B T}{N} \ln Z(N, V, T) \tag{6}$$

and eq. (4) can be applied without modification for intensive quantities if F is replaced by $\hat{F}$. For example, the pressure p and internal energy per particle $\hat{U}$ can be obtained from the derivatives along volume and temperature respectively:

$$p(\hat{V},T) = -\frac{\partial \hat{F}}{\partial \hat{V}} , \tag{7}$$

$$\hat{U}(\hat{V},T) = -T^2 \frac{\partial}{\partial T}\left[\frac{\hat{F}}{T} \right] . \tag{8}$$

Other physically-meaningful derivatives are given in Table 1. The free energy per particle $\hat{F}$ therefore serves as a more useful Equation of State for bulk condensed-phase substances, where intensive properties are of greater interest than extensive ones due to trivial scaling with particle number. As such, for the rest of this chapter, we refer only to intensified quantities as per eq. (5) implicitly; that is, the circumflex symbol $\hat{}$, and the phrase "per particle", will not be used explicitly but should be kept in mind.

Table 1: **Physically-meaningful derivatives of the free energy per particle,** corresponding to macroscopic expectations of bulk thermodynamic observables in the canonical ensemble. All of these observables therefore can be calculated, if the form of $\hat{F}(\hat{V}, T)$ is known.

Name	**Symbol**	**Definition**	**Equivalent Identity**
Pressure	p	$-\left(\frac{\partial \hat{F}}{\partial \hat{V}}\right)_T$	
Entropy per particle	$\hat{S}$	$-\left(\frac{\partial \hat{F}}{\partial T}\right)_{\hat{V}}$	
Internal energy per particle	$\hat{U}$	$\frac{1}{N}\langle \mathcal{H} \rangle_{NVT}$	$\hat{F} + T\hat{S}$
Isochoric heat capacity per particle	$\hat{C}_V$	$\left(\frac{\partial \hat{U}}{\partial T}\right)_{\hat{V}}$	$-T\left(\frac{\partial^2 \hat{F}}{\partial T^2}\right)_{\hat{V}}$
Isobaric heat capacity per particle	$\hat{C}_p$	$\left(\frac{\partial \hat{H}}{\partial T}\right)_p$	$\hat{C}_V + \hat{V}T\beta_T{\gamma_v}^2$
Thermal expansion coefficient	α	$\frac{1}{V}\left(\frac{\partial V}{\partial T}\right)_p$	$\beta_T \gamma_v$
Isothermal compressibility	β_T	$-\frac{1}{V}\left(\frac{\partial V}{\partial p}\right)_T$	$\left[\hat{V}\left(\frac{\partial^2 \hat{F}}{\partial \hat{V}^2}\right)_T\right]^{-1}$
Thermal pressure coefficient	γ_v	$\left(\frac{\partial p}{\partial T}\right)_V$	$-\frac{\partial^2 \hat{F}}{\partial \hat{V} \partial T}$
Isentropic bulk modulus	κ_s	$-V\left(\frac{\partial p}{\partial V}\right)_S$	$\frac{1}{\beta_T} + \frac{\hat{V}T{\gamma_v}^2}{\hat{C}_V}$

2.3 The dimensionless free energy

For the purposes of computer-assisted numerical optimization and fitting, it is often easier to work with dimensionless, purely numerical functions. A further simplification, as often performed in experimentally-fitted multiparameter EoSs,[62–66] can be made by factorizing $k_B T$ out from the free energy to leave a nondimensional quantity:

$$f \quad := \quad \frac{1}{k_B T} F \quad = \quad -\frac{1}{N} \ln Z \tag{9}$$

which we term the *dimensionless free energy*. Similarly, the volume and temperature variables can be nondimensionalized by dividing by some reference scale. In particular, the IAPWS-95[62] formulation, which serves as a standard reference for experimentally-fitted wide-range liquid and gaseous water EoSs, is expressed in terms of a *reduced density* and *inverse reduced temperature*:

$$\delta \quad := \quad \frac{V_c}{V} \,, \tag{10}$$

$$\tau \quad := \quad \frac{T_c}{T} \tag{11}$$

where $V_c = 92.9\,\text{Å}^3$ and $T_c = 647\,\text{K}$ are the volume and temperature at water's liquid-vapour critical point. The reason for this choice of variables is that it naturally bounds the low-pressure dilute-gas limit to a finite numerical range $0 < \delta, \tau < 1$, where the fluid behaves closely to an ideal gas and therefore does not require sophisticated description, while also expanding the high-pressure strongly-interacting dense-liquid regime to a much broader numerical range $1 < \delta, \tau < \infty$ which can be more easily adjusted by functional optimization. Scaling the variables based on V_c and T_c also fixes the critical point at $\delta = \tau = 1$, allowing non-analytic divergent terms to be inserted by hand at this specific point in the EoS.

Moreover, this choice of variables $f(\delta, \tau)$ simplifies many of the physically-meaningful free energy derivatives in Table 1, perhaps by coincidence. The pressure and internal energy from eqs. (7) and (8), for example, become:

$$\frac{pV}{k_B T} \quad = \quad \delta \frac{\partial f}{\partial \delta} \,, \tag{12}$$

$$\frac{U}{k_B T} \quad = \quad \tau \frac{\partial f}{\partial \tau} \tag{13}$$

where the LHS terms are known as the *reduced pressure* and *reduced internal energy*, noting their similarity to the expected expressions for the ideal gas. In fact, it can be easily shown — just from the above two equations — that the dimensionless free energy for the monatomic gas is given by $f(\delta, \tau) = \ln \delta + \frac{3}{2} \ln \tau$, showing the elegance of this representation. Other derivatives are shown in Table 2. We thus proceed with developing liquid water EoSs using the $f(\delta, \tau)$ form.

Table 2: **Dimensionally reduced thermodynamic observables from Table 1**, expressed in terms of the dimensionless free energy $f(\delta, \tau)$ as a function of reduced density $\delta = V_c/V$ and inverse reduced temperature $\tau = T_c/T$. Here subscripts denote partial derivatives, such as $f_\tau = \partial f/\partial \tau$.

Name	**Definition**	**Equivalent Identity**
Reduced pressure	$\frac{pV}{k_B T}$	δf_δ
Reduced entropy	$\frac{S}{k_B}$	$\tau f_\tau - f$
Reduced internal energy	$\frac{U}{k_B T}$	τf_τ
Reduced isochoric heat capacity	$\frac{C_V}{k_B}$	$-\tau^2 f_{\tau\tau}$
Reduced isobaric heat capacity	$\frac{C_p}{k_B}$	$-\tau^2 f_{\tau\tau} + \frac{\delta\left(f_\delta - \tau f_{\delta\tau}\right)^2}{2f_\delta + \delta f_{\delta\delta}}$
Reduced thermal expansion coeff.	αT	$\frac{f_\delta - \tau f_{\delta\tau}}{2f_\delta + \delta f_{\delta\delta}}$
Reduced isothermal compressibility	$\frac{\beta_T k_B T}{V}$	$\left(2\delta f_\delta + \delta^2 f_{\delta\delta}\right)^{-1}$
Reduced thermal pressure coeff.	$\frac{\gamma_v V}{k_B}$	$\delta f_\delta - \delta\tau f_{\delta\tau}$
Reduced isentropic bulk modulus	$\frac{\kappa_s V}{k_B T}$	$2\delta f_\delta + \delta^2 f_{\delta\delta} - \frac{\left(\delta f_\delta - \delta\tau f_{\delta\tau}\right)^2}{\tau^2 f_{\tau\tau}}$

3 Constructing the EoS

3.1 Problem statement

If both pressure and internal energy as functions of volume and temperature are known, then $f(\delta, \tau)$ can determined up to an additive constant by integrating eqs. (12) and (13). Fortunately, both of these are easily accessed in MD simulations. The EoS for a given water model can therefore be constructed by performing a sequence of canonical ensemble MD simulations of liquid water, fixed at varying volumes and temperatures, and obtaining a set of data points for $p(V, T)$ and $U(V, T)$; these measurements can then in principle be numerically integrated using e.g. the trapezoidal rule. In practice, however, a very fine grid of measurements is required to resolve curvatures in the integrand, resulting in prohibitively high cost for computationally expensive models. Furthermore, the sampled pressures and internal energies from each simulation are subject to statistical noise, and integrating them directly would result in an accumulation of correlated errors.

A preferable approach would be to instead construct a general flexible approximation or "model" for the dimensionless free energy function, by prescribing a fixed composition of elementary mathematical operations, and best-fitting the derivatives of the model onto eqs. (12) and (13) using standard regression techniques. The resulting best-fit model is therefore a targeted approximator for the true underlying $f(\delta, \tau)$, and is known as a *representation* for the Equation of State.

To be precise: suppose we have performed N_{sim} many simulations at the volumes $\{V_1, V_2, \ldots\}$ and temperatures $\{T_1, T_2, \ldots\}$, with the corresponding pressure results $\{p_1, p_2, \ldots\}$ and internal energy results $\{U_1, U_2, \ldots\}$. We wish to construct a model $\tilde{f}(\delta, \tau; \boldsymbol{\theta})$, which has some set of parameters $\boldsymbol{\theta} = \{\theta_1, \theta_2, \ldots\}$ controlling the behaviour of $\tilde{f}$. We may make $\tilde{f}(\delta, \tau; \boldsymbol{\theta})$ a good approximation of the true $f(\delta, \tau)$, by minimizing the *loss function*:

$$\begin{aligned} \mathcal{L}(\boldsymbol{\theta}) &= \sum_{i=1}^{N_{\text{sim}}} \left| \nabla \tilde{f}(\delta_i, \tau_i; \boldsymbol{\theta}) - \nabla f(\delta_i, \tau_i) \right|^2 \\ &= \sum_{i=1}^{N_{\text{sim}}} \left[\left(\frac{\partial \tilde{f}(\delta_i, \tau_i; \boldsymbol{\theta})}{\partial \delta} - \frac{p_i V_i}{k_B T_i} \right)^2 + \left(\frac{\partial \tilde{f}(\delta_i, \tau_i; \boldsymbol{\theta})}{\partial \tau} - \frac{U_i}{k_B T_i} \right)^2 \right] \end{aligned} \tag{14}$$

so that the resulting best-fit $\tilde{f}(\delta, \tau; \boldsymbol{\theta}_{\text{min}})$ serves as a good representation of the EoS, and can be used to extract the thermodynamic observables of Table 2 efficiently and accurately.

The expressions of eqs. (1) and (9) guarantee that the target $f(\delta, \tau)$ is analytic almost everywhere within the phase space for a continuous state of matter, but the true function may be arbitrarily complex. The difficulty of modelling liquid water is shown, for example, by the plethora of functional forms and fitting parameters used in experimentally-fitted EoSs like IAPWS-95[62] and GERG-2008,[63] which employ 249 and 68 free fitting parameters respectively. Ultimately, there is no guarantee that a "simple" EoS can be constructed for any water model,

especially if water's thermodynamic anomalies are to be captured faithfully. Simple polynomials, for example, may encounter Runge's phenomenon at high polynomial orders,[67,68] leading to unphysical model "blow-ups" even within the interpolation region. Experimentally-fitted EoSs choose a mixture of complex mathematical forms, such as logarithmic, exponential, and trigonometric functions, which are often hand-selected directly or indirectly,[62–66] but such ad-hoc models are inherently difficult to extend to new regimes, as shown by the variety of forms used in literature.

We therefore turn to a method inspired by the work of Brown (ref. 69), using local basis functions to build complete and stable approximators across arbitrary domains.

3.1.1 Local basis functions

For the context of this chapter, we define a family of C^k-smooth *local basis functions* to be a set of functions $\{B_1(x,y), B_2(x,y), \ldots\}$ satisfying:

- Each $B_i(x,y)$ is itself a C^k-smooth function everywhere on $\mathbb{R}^2$;
- For each $B_i(x,y)$, there is a finite simply-connected domain $\mathcal{D}_i \subset \mathbb{R}^2$ such that $B_i(x,y) = 0$ for all points outside $\mathcal{D}_i$;
- For any point (x,y), there exists at least one i such that (x,y) is in $\mathcal{D}_i$;
- The inner product is $\iint_{\mathbb{R}^2} B_i(x,y)B_j(x,y)\,\mathrm{d}x\,\mathrm{d}y = 0$ for all $i \neq j$;
- The norm $\iint_{\mathbb{R}^2} (B_i(x,y))^2\,\mathrm{d}x\,\mathrm{d}y$ is positive and finite (i.e. exists) for all i.

Note that the first two conditions also imply that $B_i(x,y)$, as well as all of its derivatives up to k^{th} order, go smoothly to zero at the boundary of $\mathcal{D}_i$. The first two conditions together thus mean that each B_i acts only as a localized "perturbation" within a small region of $\mathbb{R}^2$, while the third condition means that the domains of all the $\{B_i\}$ together "patch" the full space of inputs, and the fourth and fifth conditions allow the set of $\{B_i\}$ to form a complete orthonormal basis for the space of square-integrable functions.

We can therefore write any function as a linear combination of these local basis functions:

$$\tilde{f}(x,y;\boldsymbol{\theta}) \quad = \quad \sum_i c_i B_i(x,y) \tag{15}$$

where the coefficients $\{c_1, c_2, \ldots\}$ serve as the model parameters $\boldsymbol{\theta}$.

The use of local basis functions has a two-fold advantage. Since they form a complete orthonormal basis, the model is guaranteed to systematically converge towards the true underlying function as more terms are included in the expression of eq. (15), as long as there is sufficient data to calculate the coefficients c_i. More importantly, the use of localized functions allow independent additive perturbations to be made to the EoS representation; parameters

in one thermodynamic regime need not change if new measurements in another regime require modification of the EoS. This makes the model *stable* against small perturbations of the fitting data.

Furthermore, the linear form of eq. (15) reduces the loss function in eq. (14) to a standard linear least-squares problem, which can be expressed as:

$$\mathcal{L}(\boldsymbol{\theta}) \;=\; \left| \begin{pmatrix} \partial_\delta B_1(\delta_1,\tau_1) & \partial_\delta B_2(\delta_1,\tau_1) & \partial_\delta B_3(\delta_1,\tau_1) & \dots \\ \partial_\delta B_1(\delta_2,\tau_2) & \partial_\delta B_2(\delta_2,\tau_2) & \partial_\delta B_3(\delta_2,\tau_2) & \dots \\ \vdots & \vdots & \vdots & \ddots \\ \partial_\tau B_1(\delta_1,\tau_1) & \partial_\tau B_2(\delta_1,\tau_1) & \partial_\tau B_3(\delta_1,\tau_1) & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \end{pmatrix} - \begin{pmatrix} p_1V_1/k_BT_1 \\ p_2V_2/k_BT_2 \\ \vdots \\ U_1/k_BT_1 \\ \vdots \end{pmatrix} \right|^2 \tag{16}$$

noting that the linear least squares problem $\mathcal{L}(\boldsymbol{\theta}) = |A\boldsymbol{\theta} - \mathbf{b}|^2$ has the well-known solution $\boldsymbol{\theta}_{\min} = (A^TA)^{-1}A^T\mathbf{b}$. This least-squares problem can also be further modified to include inverse-variance weighting, e.g. in the case of heteroscedastic data, or to propagate data uncertainties into parameter covariances.

3.1.2 Cardinal B-splines

The notation used in this section differs from the standard notation (e.g. as used in ref. 70). We begin by reviewing B-splines in the 1D case, before elaborating the 2D case.

A B-spline of order $p+1$ is a collection of local basis functions $B_i^{(p)}(x)$, which are each piecewise polynomials of degree p, defined by a sequence of m many "knots" demarcating the domains of each $B_i^{(p)}(x)$. We denote these knots as $\{t_1, t_2, \dots, t_m\}$, which must be sorted into nondecreasing order $t_1 \le t_2 \le \dots \le t_m$. The B-spline is said to be "cardinal" if its knots are spaced equally; that is to say, the interval $\delta t = t_{i+1} - t_i$ is the same for all i. For our methodology, we use only cardinal B-splines.

B-splines may be constructed recursively via the Cox–de Boor formula. This starts with the $p = 0$ case, which are piecewise constant functions:

$$B_i^{(0)}(x) \;:=\; \begin{cases} 1 & \text{if } t_i \le x < t_{i+1}\,, \\ 0 & \text{otherwise.} \end{cases} \tag{17}$$

The higher order B-splines are then defined by recursion:

$$B_i^{(p)}(x) \;:=\; \frac{x - t_i}{p\,\delta t}\, B_i^{(p-1)}(x) \;+\; \frac{t_{i+p+1} - x}{p\,\delta t}\, B_{i+1}^{(p-1)}(x)\,. \tag{18}$$

It can be checked that the resulting $B_i^{(p)}(x)$ is zero everywhere outside of $t_i \le x < t_{i+p+1}$, and is C^{p-1}-smooth. It can also be checked that the set of $\{B_1^{(p)}, B_2^{(p)}, \dots, B_{m-p-1}^{(p)}\}$ forms

a complete orthonormal basis within the space of $t_1 \leq x < t_m$, and both the derivative and integral of a B-spline are themselves B-splines.

Visualizing these B-splines, especially from the expression of eq. (18), may be challenging; the author finds it easier to visualize B-splines using the fact that their derivatives follow a similar recursion:

$$\frac{\mathrm{d}}{\mathrm{d}x} B_i^{(p)}(x) \;=\; \frac{1}{\delta t}\, B_i^{(p-1)}(x) \;-\; \frac{1}{\delta t}\, B_{i+1}^{(p-1)}(x) \tag{19}$$

i.e. the derivative of $B_i^{(p)}$ is simply "two copies" of $B_i^{(p-1)}$ added together, with one copy translated rightwards by δt and inverted, and the value of $B_i^{(p)}$ is obtained by integrating its derivative. An example is shown in Figure 1.

B-splines are highly useful for constructing smooth approximations of single-variable functions $f(x)$ within the range of $t_1 \leq x < t_m$, as they form a complete basis, and feature the same general ease of representation and computation associated with simple polynomials, but do not suffer from Runge's phenomenon. The usual expression is to use one specific value of p, which is typically not very large, and sum over the indices i representing translations of the basis functions over the knots:

$$f(x) \;\approx\; \sum_{i=1}^{m-p-1} c_i B_i^{(p)}(x) \tag{20}$$

with this being indicated as a "B-spline approximation of order $p+1$". Loosely speaking, the B-splines are able to approximate $f(x)$ by cutting the domain up into short intervals $t_i \leq x < t_{i+1}$, and fitting degree p polynomials independently within each interval with the constraint that all of the polynomials must connect smoothly. The local error of the approximation, at any point, is therefore on the order of $\mathcal{O}(\delta t^p)$. The finite truncation of the polynomials to a low degree p and short interval δt prevents spurious local divergences; as such, the accuracy of the approximation is usually improved by increasing the number of knots m, while leaving p small.

For our context, we may extend the theory of 1D B-splines into 2D by taking the tensor product of the basis functions over x and y:

$$\tilde{f}(x, y; \boldsymbol{\theta}) \;=\; \sum_{i=1}^{m_x-p_x-1} \sum_{j=1}^{m_y-p_y-1} c_{ij} B_i^{(p_x)}(x) B_j^{(p_y)}(y) \tag{21}$$

noting that the number of knots m_y, knot sequence $\{t_1, t_2, \ldots, t_{m_y}\}$, and order p_y along the y-axis may not necessarily be the same as for the x-axis.




Figure 1: **Construction of 1D cardinal B-splines,** starting from $p = 0$ in the top row to $p = 3$ in the bottom row, using the derivative recursion of eq. (19).

3.1.3 Regularized B-splines for EoS models

For simplicity, we construct our EoS model using 2D cardinal B-splines with matching order $p_x = p_y = 4$. The choice of using degree 4 polynomials to represent the dimensionless free energy $f(\delta, \tau)$ ensures that its second derivatives, such as heat capacity, are piecewise quadratic. The knot sequences are determined by the input fitting data, chosen such that the endpoints t_1 and t_m are equal to the extremal values in the fitting data, and the total number of knot intervals in both dimensions is as close to the number of data points as possible. This means that the domain of (δ, τ) space covered by the model is the bounding box of the input fitting data, and is divided into subdomains spanned by smoothly-connected polynomials of degree 4.

This construction is already suitable for the linear least-squares form presented in eq. (16), if the tensorial coefficients c_{ij} are rearranged into a single vector of parameters $\boldsymbol{\theta}$. However, in practice the number of model parameters may exceed the number of independent data points — or, one or more subdomains may contain no data points, causing columns of zeros to occur in the coefficient matrix — leading to an underdetermined least-squares problem whose solution would be overfitted.

The solution to this is to introduce a *regularization*, which penalizes undesirable behaviours in the model fitting. In our case, we expect that the heat capacity, compressibility, and thermal pressure coefficient of any continuous phase of matter should be a smooth surface; these observables are all second derivatives of the free energy, hence a sensible regularization would be to minimize their curvatures by minimizing the fourth derivatives $\partial_\delta^4 \tilde{f}$ and $\partial_\tau^4 \tilde{f}$ of the model. The choice of using B-spline order $p = 4$ guarantees that these fourth derivatives are piecewise constant within each subdomain, so there are an integer number of values of $\partial_\delta^4 \tilde{f}$ and $\partial_\tau^4 \tilde{f}$ which allows for easy construction of a *regularization matrix*.

Writing $\boldsymbol{\delta}$, $\boldsymbol{\tau}$, $\mathbf{V}$, $\mathbf{T}$, $\mathbf{p}$, $\mathbf{U}$ as the data vectors (i.e. observations from simulation data) of length N_{sim}, and $\boldsymbol{\theta}$ as the vector of model parameters, the regularized least-squares problem therefore has loss function given by:

$$\mathcal{L}(\boldsymbol{\theta}) = \left| \begin{pmatrix} \nabla_{\boldsymbol{\theta}}^{\mathrm{T}} \left[\partial_\delta \tilde{f}(\boldsymbol{\delta}, \boldsymbol{\tau}) \right] \\ \nabla_{\boldsymbol{\theta}}^{\mathrm{T}} \left[\partial_\tau \tilde{f}(\boldsymbol{\delta}, \boldsymbol{\tau}) \right] \\ \lambda \nabla_{\boldsymbol{\theta}}^{\mathrm{T}} \left[\partial_\delta^4 \tilde{f}(\boldsymbol{\delta}, \boldsymbol{\tau}) \right] \\ \lambda \nabla_{\boldsymbol{\theta}}^{\mathrm{T}} \left[\partial_\tau^4 \tilde{f}(\boldsymbol{\delta}, \boldsymbol{\tau}) \right] \end{pmatrix} \boldsymbol{\theta} - \begin{pmatrix} \mathbf{pV}/k_B\mathbf{T} \\ \mathbf{U}/k_B\mathbf{T} \\ \mathbf{0} \\ \mathbf{0} \end{pmatrix} \right| \tag{22}$$

where λ is a positive real number known as the *regularization parameter*; setting $\lambda \to 0$ returns the fitting problem to the unregularized case, whereas setting $\lambda \to \infty$ causes the model to prioritize "flattening" the curve everywhere instead of fitting the curve through the data points. A good choice of λ is to set it high enough such that the reduced χ^2 of the fit, i.e. fitting residuals normalized by data uncertainty, is on the order of unity. This can be achieved by performing a series of trial fits over varying λ, and selecting the fit with reduced χ^2 closest to one. An example is shown in Figure 2.




Figure 2: **Regularization of a 1D B-spline;** increasing the regularization parameter λ leads to a straight-line fit (top left), while decreasing λ leads to overfitting (bottom right). The best-fit is achieved when the reduced χ^2 is close to one (middle right). This figure is reproduced from ref. 69.

3.1.4 Summary: fitting EoSs

We choose to model the dimensionless free energy using cardinal B-splines in 2D, constructed by taking the tensor product of 1D B-splines of order $p = 4$:

$$\tilde{f}(\delta, \tau; \boldsymbol{\theta}) \;=\; \sum_{i=1}^{m_x - 5} \sum_{j=1}^{m_y - 5} c_{ij} B_i^{(4)}(\delta) B_j^{(4)}(\tau) \tag{23}$$

where the knot sequences are chosen to span the input fitting data's bounding box, with roughly as many intervals as data points. This model is then fitted to the input data using the regularized least-squares loss function given by eq. (22), with the regularization parameter λ chosen to bring the reduced χ^2 of the fit to unity.

In this way, a single set of simulation data, carried out at a finite number of volume and temperature state points V_i and T_i with corresponding results for pressure $p_i = p(V_i, T_i)$ and internal energy $U_i = U(V_i, T_i)$, allows for the construction of an approximate Equation of State $f(\delta, \tau)$, and therefore obtainment of all thermodynamic observables given in Table 2.

Uncertainty estimates can then be performed using *bootstrapping*, a generic technique which makes no assumption on the fitting parameters' covariances. Specifically, we may construct an *ensemble* of EoS representations $\tilde{f}(\delta, \tau; \boldsymbol{\theta})$, where each model is fitted to randomly-perturbed copies of the original data with one point randomly removed. Hence, for any thermodynamic observable, we may "query" the entire ensemble of models and use the ensemble spread as a good estimator for the inherent uncertainty of the predicted observable.

4 Preliminary Results and Discussion

For testing purposes, we fitted an EoS to synthetic data for liquid water, generated by sampling 60 points from the GERG-2008 formulation.[63] We demonstrate that the model is able to replicate the internal energy and pressure surfaces $U(V, T)$ and $p(V, T)$, and thus extract thermodynamic observables along isochores $V = const$. More importantly, we also demonstrate that numerical inversion of the $V(p)$ curve at any given temperature is possible, allowing thermodynamic observables to be extracted along isotherms $T = const$ as a function of pressure, or along isobars $p = const$ as a function of temperature. That said, we observe that model bootstrapping can cause localized noise in the pressure surface, causing the $p(V)$ curve to be non-monotonic in a small area and thus generating a large uncertainty in the inversion of $V(p)$. See figures below.




Figure 3: Internal energy, predicted by the EoS $f(\delta, \tau)$ fitted on synthetic data. Note that synthetic data was supplied with nominal uncertainties of 5 meV on every point, so the fitting errors being of similar magnitude is consistent with the $\chi^2 = 1$ regularization.




Figure 4: Pressure, predicted by the EoS $f(\delta, \tau)$ fitted on synthetic data. Note that synthetic data was supplied with nominal uncertainties of 5 bar on every point, so the fitting errors being of similar magnitude is consistent with the $\chi^2 = 1$ regularization.




Figure 5: Isochores, predicted by the EoS $f(\delta, \tau)$ fitted on synthetic data. All curves are mostly smooth, except for the heat capacity C_V which has poor extrapolation to low temperatures, and the model reports unusually large uncertainty for low densities as the data only extends to $900\,\mathrm{kg/m^3}$.




Figure 6: Isotherms, predicted by the EoS $f(\delta, \tau)$ fitted on synthetic data; localized uncertainty spikes correspond to the model predicting a weakly noisy $V(p)$ curve at that pressure. The vertical lines in the negative-pressure range for the thermal expansion coefficient α and compressibility β is inherent to the source data, generated from the GERG-2008 formulation (see below), and is thus not a concern.




Figure 7: Isobars, predicted by the EoS $f(\delta, \tau)$ fitted on synthetic data; same comment as above wrt localized uncertainty spikes.




Figure 8: Pressures from the reference GERG-2008 formulation, which the above model was targeting.




Figure 9: Isochores from the reference GERG-2008 formulation, which the above model was targeting.




Figure 10: Isotherms from the reference GERG-2008 formulation, which the above model was targeting. The vertical lines in the negative-pressure range for the thermal expansion coefficient α and compressibility β is an inherent limitation of the GERG-2008 formulation, which was only designed for use in near-ambient conditions.




Figure 11: Isobars from the reference GERG-2008 formulation, which the above model was targeting.

References

[1] Helena L Pi, Juan L Aragones, Carlos Vega, Eva G Noya, Jose L F Abascal, Miguel A Gonzalez, and Carl McBride. Anomalies in water as obtained from computer simulations of the TIP4P/2005 model: density maxima, and density, isothermal compressibility and heat capacity minima. *Mol. Phys.*, 107(4-6):365–374, February 2009. doi: 10.1080/00268970902784926.

[2] C H Cho, S Singh, and G W Robinson. An explanation of the density maximum in water. *Phys. Rev. Lett.*, 76(10):1651–1654, March 1996. doi: 10.1103/PhysRevLett.76.1651.

[3] Gaël Pallares, Miguel A Gonzalez, Jose Luis F Abascal, Chantal Valeriani, and Frédéric Caupin. Equation of state for water and its line of density maxima down to -120 MPa. *Phys. Chem. Chem. Phys.*, 18(8):5896–5900, February 2016. doi: 10.1039/c5cp07580g.

[4] Yizhi Song, Renxi Liu, Chunyi Zhang, Yifan Li, Biswajit Santra, Mohan Chen, Michael L Klein, and Xifan Wu. Understanding the density maximum of water with machine-learned potentials. *Sci. Adv.*, 12(22):eaec6748, May 2026. doi: 10.1126/sciadv.aec6748.

[5] C A Angell and J C Tucker. Anomalous heat capacities of supercooled water and heavy water. *Science*, 181(4097):342–344, July 1973. doi: 10.1126/science.181.4097.342.

[6] C A Angell, J Shuppert, and J C Tucker. Anomalous properties of supercooled water. heat capacity, expansivity, and proton magnetic resonance chemical shift from 0 to -38%. *J. Phys. Chem.*, 77(26):3092–3099, December 1973. doi: 10.1021/j100644a014.

[7] Kam-Tung Chan, Dylan A Folkner, Zekun Chen, Margaret L Berrens, Alexei A Stuchebrukhov, Lee-Ping Wang, and Davide Donadio. Structural origin of water heat capacity anomaly from classical and quantum simulations. *arXiv preprint 2607.08957*, July 2026. doi: 10.48550/arXiv.2607.08957.

[8] E Tombari, C Ferrari, and G Salvetti. Heat capacity anomaly in a large sample of supercooled water. *Chem. Phys. Lett.*, 300(5-6):749–751, February 1999. doi: 10.1016/S0009-2614(98)01392-X.

[9] Oliver R Gittus and Fernando Bresme. Unravelling the origin of water's thermal conductivity maximum: Compressibility, tetrahedrality and nuclear quantum effects. *J. Am. Chem. Soc.*, 147(46):42175–42183, November 2025. doi: 10.1021/jacs.4c12898.

[10] M L Huber, R A Perkins, D G Friend, J V Sengers, M J Assael, I N Metaxa, K Miyagawa, R Hellmann, and E Vogel. New international formulation for the thermal conductivity of H2O. *J. Phys. Chem. Ref. Data*, 41(3):033102, September 2012. doi: 10.1063/1.4738955.

[11] Pradeep Kumar and H Eugene Stanley. Thermal conductivity minimum: a new water anomaly. *J. Phys. Chem. B*, 115(48):14269–14273, December 2011. doi: 10.1021/jp2051867.

[12] C A Angell, E D Finch, and P Bach. Spin–echo diffusion coefficients of water to 2380 bar and -20°c. *J. Chem. Phys.*, 65(8):3063–3066, October 1976. doi: 10.1063/1.433518.

[13] Sungho Han, Pradeep Kumar, and H Eugene Stanley. Absence of a diffusion anomaly of water in the direction perpendicular to hydrophobic nanoconfining walls. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.*, 77(3 Pt 1):030201, March 2008. doi: 10.1103/PhysRevE.77.030201.

[14] F X Prielmeier, E W Lang, R J Speedy, and H Lüdemann. Diffusion in supercooled water to 300 MPa. *Phys. Rev. Lett.*, 59(10):1128–1131, September 1987. doi: 10.1103/PhysRevLett.59.1128.

[15] Xiaojing Teng, Bailang Liu, and Toshiko Ichiye. Understanding how water models affect the anomalous pressure dependence of their diffusion coefficients. *J. Chem. Phys.*, 153(10):104510, September 2020. doi: 10.1063/5.0021472.

[16] Alexander Gorfer, Christoph Dellago, and Marcello Sega. High-density liquid (HDL) adsorption at the supercooled water/vapor interface and its possible relation to the second surface tension inflection point. *J. Chem. Phys.*, 158(5):054503, February 2023. doi: 10.1063/5.0132985.

[17] Marcello Sega, George Horvai, and Pál Jedlovszky. Microscopic origin of the surface tension anomaly of water. *Langmuir*, 30(11):2969–2972, March 2014. doi: 10.1021/la404868e.

[18] Václav Vinš, Jiří Hykl, Jan Hrubý, Aleš Blahut, David Celný, Miroslav Čenský, and Olga Prokopová. Possible anomaly in the surface tension of supercooled water: New experiments at extreme supercooling down to -31.4 °c. *J. Phys. Chem. Lett.*, 11(11):4443–4447, June 2020. doi: 10.1021/acs.jpclett.0c01163.

[19] Jiaxing Yuan, Kun Qiu, Gang Sun, and Hajime Tanaka. Competing hydrogen-bond orders drive water's anomalous surface tension. *Nat. Commun.*, 17(1), February 2026. doi: 10.1038/s41467-026-69356-6.

[20] Pablo G Debenedetti and H Eugene Stanley. Supercooled and glassy water. *Phys. Today*, 56(6):40–46, June 2003. doi: 10.1063/1.1595053.

[21] Pablo G Debenedetti. Supercooled and glassy water. *J. Phys. Condens. Matter*, 15(45):R1669–R1726, November 2003. doi: 10.1088/0953-8984/15/45/R01.

[22] Philip H Handle, Thomas Loerting, and Francesco Sciortino. Supercooled and glassy water: Metastable liquid(s), amorphous solid(s), and a no-man's land. *Proc. Natl. Acad. Sci. U. S. A.*, 114(51):13336–13344, December 2017. doi: 10.1073/pnas.1700103114.

[23] Gaël Pallares, Mouna El Mekki Azouzi, Miguel A González, Juan L Aragones, José L F Abascal, Chantal Valeriani, and Frédéric Caupin. Anomalies in bulk supercooled water at negative pressure. *Proc. Natl. Acad. Sci. U. S. A.*, 111(22):7936–7941, June 2014. doi: 10.1073/pnas.1323366111.

[24] L Fumagalli, A Esfandiar, R Fabregas, S Hu, P Ares, A Janardanan, Q Yang, B Radha, T Taniguchi, K Watanabe, G Gomila, K S Novoselov, and A K Geim. Anomalously low dielectric constant of confined water. *Science*, 360(6395):1339–1342, June 2018. doi: 10.1126/science.aat4191.

[25] R Wang, M Souilamas, A Esfandiar, R Fabregas, S Benaglia, H Nevison-Andrews, Q Yang, J Normansell, P Ares, G Ferrari, A Principi, A K Geim, and L Fumagalli. In-plane dielectric constant and conductivity of confined water. *Nature*, 646(8085):606–610, October 2025. doi: 10.1038/s41586-025-09558-y.

[26] T Dufils, C Schran, J Chen, A K Geim, L Fumagalli, and A Michaelides. Origin of dielectric polarization suppression in confined water from first principles. *Chem. Sci.*, 15(2):516–527, January 2024. doi: 10.1039/D3SC04740G.

[27] B Radha, A Esfandiar, F C Wang, A P Rooney, K Gopinadhan, A Keerthi, A Mishchenko, A Janardanan, P Blake, L Fumagalli, M Lozada-Hidalgo, S Garaj, S J Haigh, I V Grigorieva, H A Wu, and A K Geim. Molecular transport through capillaries made with atomic-scale precision. *Nature*, 538(7624):222–225, October 2016. doi: 10.1038/nature19363.

[28] Eleonora Secchi, Sophie Marbach, Antoine Niguès, Derek Stein, Alessandro Siria, and Lydéric Bocquet. Massive radius-dependent flow slippage in carbon nanotubes. *Nature*, 537(7619):210–213, September 2016. doi: 10.1038/nature19315.

[29] Xavier R Advincula, Yair Litman, Kara D Fong, William C Witt, Christoph Schran, and Angelos Michaelides. How reactive is water at the nanoscale and how to control it? *Sci. Adv.*, 12(26):eaeb5772, June 2026. doi: 10.1126/sciadv.aeb5772.

[30] Venkat Kapil, Christoph Schran, Andrea Zen, Ji Chen, Chris J Pickard, and Angelos Michaelides. The first-principles phase diagram of monolayer nanoconfined water. *Nature*, 609(7927):512–516, September 2022. doi: 10.1038/s41586-022-05036-x.

[31] Jian Jiang, Yurui Gao, Lei Li, Yuan Liu, Weiduo Zhu, Chongqin Zhu, Joseph S Francisco, and Xiao Cheng Zeng. Rich proton dynamics and phase behaviours of nanoconfined ices. *Nat. Phys.*, 20(3):456–464, March 2024. doi: 10.1038/s41567-023-02341-8.

[32] Masumeh Foroutan, S Mahmood Fatemi, and Farshad Esmaeilian. A review of the structure and dynamics of nanoconfined water and ionic liquids via molecular dynamics simulation. *Eur. Phys. J. E Soft Matter*, 40(2):19, February 2017. doi: 10.1140/epje/i2017-11507-7.

[33] Claudia Goy, Gregory Kimmel, Ying Jiang, and Chantal Valeriani. Advances in unveiling water's molecular mysteries. *J. Chem. Phys.*, 163(11):110401, September 2025. doi: 10.1063/5.0292407.

[34] Charlotte I Lynch, Shanlin Rao, and Mark S P Sansom. Water in nanopores and biological channels: A molecular simulation perspective. *Chem. Rev.*, 120(18):10298–10335, September 2020. doi: 10.1021/acs.chemrev.9b00830.

[35] Yuki Nagata, Tatsuhiko Ohto, Ellen H G Backus, and Mischa Bonn. Molecular modeling of water interfaces: From molecular spectroscopy to thermodynamics. *J. Phys. Chem. B*, 120(16):3785–3796, April 2016. doi: 10.1021/acs.jpcb.6b01012.

[36] Paolo La Francesca and Paola Gallo. Molecular dynamics simulations of supercooled aqueous solutions of calcium perchlorate: Thermodynamics and structure of Martian solutes in TIP4P/2005 water. *J. Phys. Chem. B*, 129(34):8776–8787, August 2025. doi: 10.1021/acs.jpcb.5c03712.

[37] Anders Nilsson and Lars G M Pettersson. The structural origin of anomalous properties of liquid water. *Nat. Commun.*, 6(1):8998, December 2015. doi: 10.1038/ncomms9998.

[38] Jeremy C Palmer, Fausto Martelli, Yang Liu, Roberto Car, Athanassios Z Panagiotopoulos, and Pablo G Debenedetti. Metastable liquid-liquid transition in a molecular model of water. *Nature*, 510(7505):385–388, June 2014. doi: 10.1038/nature13405.

[39] Aleks Reinhardt, Mandy Bethkenhagen, Federica Coppari, Marius Millot, Sebastien Hamel, and Bingqing Cheng. Thermodynamics of high-pressure ice phases explored with atomistic simulations. *Nat. Commun.*, 13(1):4707, August 2022. doi: 10.1038/s41467-022-32374-1.

[40] Albert P Bartók, Mike C Payne, Risi Kondor, and Gábor Csányi. Gaussian approximation potentials: the accuracy of quantum mechanics, without the electrons. *Phys. Rev. Lett.*, 104(13):136403, April 2010. doi: 10.1103/PhysRevLett.104.136403.

[41] Jörg Behler and Michele Parrinello. Generalized neural-network representation of high-dimensional potential-energy surfaces. *Phys. Rev. Lett.*, 98(14):146401, April 2007. doi: 10.1103/PhysRevLett.98.146401.

[42] Volker L Deringer, Miguel A Caro, and Gábor Csányi. Machine learning interatomic potentials as emerging tools for materials science. *Adv. Mater.*, 31(46):e1902765, November 2019. doi: 10.1002/adma.201902765.

[43] Pei-Lin Kang, Cheng Shang, and Zhi-Pan Liu. Large-scale atomic simulation via machine learning potentials constructed by global potential energy surface exploration. *Acc. Chem. Res.*, 53(10):2119–2129, October 2020. doi: 10.1021/acs.accounts.0c00472.

[44] Fabian L Thiemann, Niamh O'Neill, Venkat Kapil, Angelos Michaelides, and Christoph Schran. Introduction to machine learning potentials for atomistic simulations. *J. Phys. Condens. Matter*, 37(7):073002, December 2024. doi: 10.1088/1361-648X/ad9657.

[45] Oliver T Unke, Stefan Chmiela, Huziel E Sauceda, Michael Gastegger, Igor Poltavsky, Kristof T Schütt, Alexandre Tkatchenko, and Klaus-Robert Müller. Machine learning force fields. *Chem. Rev.*, 121(16):10142–10186, August 2021. doi: 10.1021/acs.chemrev.0c01111.

[46] Pascal Friederich, Florian Häse, Jonny Proppe, and Alán Aspuru-Guzik. Machine-learned potentials for next-generation matter simulations. *Nat. Mater.*, 20(6):750–761, June 2021. doi: 10.1038/s41563-020-0777-6.

[47] Hyuna Kwon, Marcos F Calegari Andrade, Shane Ardo, Daniel V Esposito, Tuan Anh Pham, and Tadashi Ogitsu. Confinement effects on proton transfer in TiO2 nanopores from machine learning potential molecular dynamics simulations. *ACS Appl. Mater. Interfaces*, 16(24):31687–31695, June 2024. doi: 10.1021/acsami.4c02339.

[48] M S Sansom and P C Biggin. Water at the nanoscale. *Nature*, 414(6860):156–159, November 2001. doi: 10.1038/35102651.

[49] Stefan Chmiela, Huziel E Sauceda, Klaus-Robert Müller, and Alexandre Tkatchenko. Towards exact molecular dynamics simulations with machine-learned force fields. *Nat. Commun.*, 9(1):3887, September 2018. doi: 10.1038/s41467-018-06169-2.

[50] Anna-Lena M Fischer, Anna Tichy, Janik Kokot, Valentin J Hoerschinger, Robert F Wild, Jakob R Riccabona, Johannes R Loeffler, Franz Waibl, Patrick K Quoika, Philipp Gschwandtner, Stefano Forli, Andrew B Ward, Klaus R Liedl, Martin Zacharias, and Monica L Fernández-Quintero. The role of force fields and water models in protein folding and unfolding dynamics. *J. Chem. Theory Comput.*, 20(5):2321–2333, March 2024. doi: 10.1021/acs.jctc.3c01106.

[51] Andreas Köster, Thomas Spura, Gábor Rutkai, Jan Kessler, Hendrik Wiebeler, Jadran Vrabec, and Thomas D Kühne. Assessing the accuracy of improved force-matched water models derived from ab initio molecular dynamics simulations. *J. Comput. Chem.*, 37(19): 1828–1838, July 2016. doi: 10.1002/jcc.24398.

[52] C Vega, J L F Abascal, M M Conde, and J L Aragones. What ice can teach us about water interactions: a critical comparison of the performance of different water models. *Faraday Discuss.*, 141:251–76; discussion 309–46, 2009. doi: 10.1039/b805531a.

[53] Carlos Vega and Jose L F Abascal. Simulating water with rigid non-polarizable models: a general perspective. *Phys. Chem. Chem. Phys.*, 13(44):19663–19688, November 2011. doi: 10.1039/c1cp22168j.

[54] Valeria Molinero and Emily B Moore. Water modeled as an intermediate element between carbon and silicon. *J. Phys. Chem. B*, 113(13):4008–4016, April 2009. doi: 10.1021/jp805227c.

[55] J L F Abascal and C Vega. A general purpose model for the condensed phases of water: TIP4P/2005. *J. Chem. Phys.*, 123(23):234505, December 2005. doi: 10.1063/1.2121687.

[56] Saeed Izadi, Ramu Anandakrishnan, and Alexey V Onufriev. Building water models: A different approach. *J. Phys. Chem. Lett.*, 5(21):3863–3871, November 2014. doi: 10.1021/jz501780a.

[57] Sergei Izvekov and Gregory A Voth. Multiscale coarse graining of liquid-state systems. *J. Chem. Phys.*, 123(13):134105, October 2005. doi: 10.1063/1.2038787.

[58] David Eisenberg and Walter Kauzmann. *The structure and properties of water*. Oxford Classic Texts in the Physical Sciences. Oxford University Press, London, England, October 2005.

[59] Felix Franks, editor. *The physics and physical chemistry of water*. Water. Springer, New York, NY, 1972 edition, December 2012.

[60] Oleksii Khorolskyi and Nikolay P Malomuzh. pH and H-bonding energy for pure water. *Chem. Phys. Lett.*, 828(140713):140713, October 2023. doi: 10.1016/j.cplett.2023.140713.

[61] A Morozenko, I V Leontyev, and A A Stuchebrukhov. Dipole moment and binding energy of water in proteins from crystallographic analysis. *J. Chem. Theory Comput.*, 10(10): 4618–4623, October 2014. doi: 10.1021/ct500358r.

[62] W Wagner and A Pruß. The IAPWS formulation 1995 for the thermodynamic properties of ordinary water substance for general and scientific use. *J. Phys. Chem. Ref. Data*, 31 (2):387–535, June 2002. doi: 10.1063/1.1461829.

[63] O Kunz and W Wagner. The GERG-2008 wide-range equation of state for natural gases and other mixtures: An expansion of GERG-2004. *J. Chem. Eng. Data*, 57(11):3032–3091, November 2012. doi: 10.1021/je300655b.

[64] U Setzmann and W Wagner. A new equation of state and tables of thermodynamic properties for methane covering the range from the melting line to 625 K at pressures up to 1000 MPa. *J. Phys. Chem. Ref. Data*, 20(6):1061–1155, November 1991. doi: 10.1063/1.555898.

[65] Roland Span and Wolfgang Wagner. A new equation of state for carbon dioxide covering the fluid region from the triple-point temperature to 1100 K at pressures up to 800 MPa. *J. Phys. Chem. Ref. Data*, 25(6):1509–1596, November 1996. doi: 10.1063/1.555991.

[66] Roland Span, Eric W Lemmon, Richard T Jacobsen, Wolfgang Wagner, and Akimichi Yokozeki. A reference equation of state for the thermodynamic properties of nitrogen for temperatures from 63.151 to 1000 K and pressures to 2200 MPa. *J. Phys. Chem. Ref. Data*, 29(6):1361–1433, November 2000. doi: 10.1063/1.1349047.

[67] James F Epperson. On the Runge example. *Am. Math. Mon.*, 94(4):329–341, April 1987. doi: 10.1080/00029890.1987.12000642.

[68] Carl D T Runge. Über empirische Funktionen und die Interpolation zwischen äquidistanten Ordinaten. *Z. Math. Phys.*, 46:224–243, 1901.

[69] J Michael Brown. Local basis function representations of thermodynamic surfaces: Water at high pressure and temperature as an example. *Fluid Phase Equilib.*, 463:18–31, May 2018. doi: 10.1016/j.fluid.2018.02.001.

[70] C de Boor. *A practical guide to splines*. Springer, Berlin, Germany, December 1978.