

SOLUTIONS TO AHLFORS' COMPLEX ANALYSIS

By:

DUSTIN SMITH

Contents

| | | |
|----------|--------------------------------------------------------------|-----------|
| 1 | Complex Numbers | 5 |
| 1.1 | The Algebra of Complex Numbers | 5 |
| 1.1.1 | Arithmetic Operations | 5 |
| 1.1.2 | Square Roots | 6 |
| 1.1.3 | Justification | 7 |
| 1.1.4 | Conjugation, Absolute Value | 7 |
| 1.1.5 | Inequalities | 10 |
| 1.2 | The Geometric Representation of Complex Numbers | 11 |
| 1.2.1 | Geometric Addition and Multiplication | 11 |
| 1.2.2 | The Binomial Equation | 12 |
| 1.2.3 | Analytic Geometry | 14 |
| 1.2.4 | The Spherical Representation | 15 |
| 2 | Complex Functions | 17 |
| 2.1 | Introduction to the Concept of Analytical Function | 17 |
| 2.1.1 | Limits and Continuity | 17 |
| 2.1.2 | Analytic Functions | 17 |
| 2.1.3 | Polynomials | 19 |
| 2.1.4 | Rational Functions | 19 |
| 2.2 | Elementary Theory of Power Series | 20 |
| 2.2.1 | Sequences | 20 |
| 2.2.2 | Series | 20 |
| 2.2.3 | Uniform Convergence | 20 |
| 2.2.4 | Power Series | 22 |
| 2.2.5 | Abel's Limit Theorem | 26 |
| 2.3 | The Exponential and Trigonometric Functions | 26 |
| 2.3.1 | The Exponential | 26 |
| 2.3.2 | The Trigonometric Functions | 26 |
| 2.3.3 | Periodicity | 28 |
| 2.3.4 | The Logarithm | 28 |
| 3 | Analytic Functions as Mappings | 31 |
| 3.1 | Elementary Point Set Topology | 31 |
| 3.1.1 | Sets and Elements | 31 |
| 3.1.2 | Metric Spaces | 31 |

1 Complex Numbers

1.1 The Algebra of Complex Numbers

1.1.1 Arithmetic Operations

1. Find the values of

$$(1 + 2i)^3, \quad \frac{5}{-3 + 4i}, \quad \left(\frac{2 + i}{3 - 2i}\right)^2, \quad (1 + i)^n + (1 - i)^n$$

For the first problem, we have $(1 + 2i)^3 = (-3 + 4i)(1 + 2i) = -11 - 2i$. For the second problem, we should multiply by the conjugate $\bar{z} = -3 - 4i$.

$$\frac{5}{-3 + 4i} \frac{-3 - 4i}{-3 - 4i} = \frac{-15 - 20i}{25} = \frac{-3}{5} - \frac{4}{5}i$$

For the third problem, we should first multiply by $\bar{z} = 3 + 2i$.

$$\frac{2 + i}{3 - 2i} \frac{3 + 2i}{3 + 2i} = \frac{8 + i}{13}$$

Now we need to just square the result.

$$\frac{1}{169}(8 + i)^2 = \frac{63 + 16i}{169}$$

For the last problem, we will need to find the polar form of the complex numbers. Let $z_1 = 1 + i$ and $z_2 = 1 - i$. Then the modulus of $z_1 = \sqrt{2} = z_2$. Let ϕ_1 and ϕ_2 be the angles associated with z_1 and z_2 , respectively. Then $\phi_1 = \arctan(1) = \frac{\pi}{4}$ and $\phi_2 = \arctan(-1) = \frac{-\pi}{4}$. Then $z_1 = \sqrt{2}e^{\pi i/4}$ and $z_2 = \sqrt{2}e^{-\pi i/4}$.

$$\begin{aligned} z_1^n + z_2^n &= 2^{n/2} [e^{n\pi i/4} + e^{-n\pi i/4}] \\ &= 2^{n/2+1} \left[\frac{e^{n\pi i/4} + e^{-n\pi i/4}}{2} \right] \\ &= 2^{n/2+1} \cos\left(\frac{n\pi}{4}\right) \end{aligned}$$

2. If $z = x + iy$ (x and y real), find the real and imaginary parts of

$$z^4, \quad \frac{1}{z}, \quad \frac{z-1}{z+1}, \quad \frac{1}{z^2}$$

For z^4 , we can use the binomial theorem since $(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$. Therefore,

$$(x + iy)^4 = \binom{4}{0}(iy)^4 + \binom{4}{1}x(iy)^3 + \binom{4}{2}x^2(iy)^2 + \binom{4}{3}x^3(iy) + \binom{4}{4}x^4 = y^4 - 4xy^3i - 6x^2y^2 + 4x^3yi + x^4$$

Then the real and imaginary parts are

$$\begin{aligned} u(x, y) &= x^4 + y^4 - 6x^2y^2 \\ v(x, y) &= 4x^3y - 4xy^3 \end{aligned}$$

For second problem, we need to multiply by the conjugate \bar{z} .

$$\frac{1}{x + iy} \frac{x - iy}{x - iy} = \frac{x - iy}{x^2 + y^2}$$

so the real and imaginary parts are

$$u(x, y) = \frac{x}{x^2 + y^2}$$

$$v(x, y) = \frac{-y}{x^2 + y^2}$$

For the third problem, we have $\frac{x-1+iy}{x+1-iy}$. Then $\bar{z} = x + 1 + iy$.

$$\frac{x-1+iy}{x+1-iy} \frac{x+1+iy}{x+1+iy} = \frac{x^2-1+2xyi}{(x+1)^2+y^2}$$

Then real and imaginary parts are

$$u(x, y) = \frac{x^2-1}{(x+1)^2+y^2}$$

$$v(x, y) = \frac{2xy}{(x+1)^2+y^2}$$

For the last problem, we have

$$\frac{1}{z^2} = \frac{x^2-y^2-2xyi}{x^4+2x^2y^2+y^4}$$

so the real and imaginary parts are

$$u(x, y) = \frac{x^2-y^2}{x^4+2x^2y^2+y^4}$$

$$v(x, y) = \frac{-2xy}{x^4+2x^2y^2+y^4}$$

3. Show that $(\frac{-1 \pm i\sqrt{3}}{2})^3 = 1$ and $(\frac{\pm 1 \pm i\sqrt{3}}{2})^6 = 1$.

Both problems will can be handled easily by converting to polar form. Let $z_1 = \frac{-1 \pm i\sqrt{3}}{2}$. Then $|z_1| = 1$. Let ϕ_+ be the angle for the positive z_1 and ϕ_- for the negative. Then $\phi_+ = \arctan(-\sqrt{3}) = \frac{2\pi}{3}$ and $\phi_- = \arctan(\sqrt{3}) = \frac{4\pi}{3}$. We can write $z_{1+} = e^{2i\pi/3}$ and $z_{1-} = e^{4i\pi/3}$.

$$z_{1+}^3 = e^{2i\pi}$$

$$= 1$$

$$z_{1-}^3 = e^{4i\pi}$$

$$= 1$$

Therefore, $z_1^3 = 1$. For the second problem, $\phi_{ij} = \pm \frac{\pi}{3}$ and $\pm \frac{2\pi}{3}$ for $i, j = +, -$ and the $|z_2| = 1$. When we raise z to the sixth poewr, the argument becomes $\pm 2\pi$ and $\pm 4\pi$.

$$e^{\pm 2i\pi} = e^{\pm 4i\pi} = z^6 = 1$$

1.1.2 Square Roots

1. Compute

$$\sqrt{i}, \quad \sqrt{-i}, \quad \sqrt{1+i}, \quad \sqrt{\frac{1-i\sqrt{3}}{2}}$$

For \sqrt{i} , we are looking for x and y such that

$$\begin{aligned} \sqrt{i} &= x + iy \\ i &= x^2 - y^2 + 2xyi \\ x^2 - y^2 &= 0 \\ 2xy &= 1 \end{aligned} \tag{1.1}$$

$$\tag{1.2}$$

From equation (1.1), we see that $x^2 = y^2$ or $\pm x = \pm y$. Also, note that i is the upper half plane (UHP). That is, the angle is positive so $x = y$ and $2x^2 = 1$ from equation (1.1). Therefore, $\sqrt{i} = \frac{1}{\sqrt{2}}(1 + i)$. We also could have done this problem using the polar form of z . Let $z = i$. Then $z = e^{i\pi/2}$ so $\sqrt{z} = e^{i\pi/4}$ which is exactly what we obtained. For $\sqrt{-i}$, let $z = -i$. Then z in polar form is $z = e^{-i\pi/2}$ so $\sqrt{z} = e^{-i\pi/4} = \frac{1}{\sqrt{2}}(1 - i)$. For $\sqrt{1+i}$, let $z = 1 + i$. Then $z = \sqrt{2}e^{i\pi/4}$ so $\sqrt{z} = 2^{1/4}e^{i\pi/8}$. Finally, for $\sqrt{\frac{1-i\sqrt{3}}{2}}$, let $z = \frac{1-i\sqrt{3}}{2}$. Then $z = e^{-i\pi/3}$ so $\sqrt{z} = e^{-i\pi/6} = \frac{1}{2}(\sqrt{3} - i)$.

2. Find the four values of $\sqrt[4]{-1}$.

Let $z = \sqrt[4]{-1}$ so $z^4 = -1$. Let $z = re^{i\theta}$ so $r^4 e^{4i\theta} = -1 = e^{i\pi(1+2k)}$.

$$\begin{aligned} r^4 &= 1 \\ \theta &= \frac{\pi}{4}(1 + 2k) \end{aligned}$$

where $k = 0, 1, 2, 3$. Since when $k = 4$, we have $k = 0$. Then $\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4},$ and $\frac{7\pi}{4}$.

$$z = e^{i\pi/4}, e^{3i\pi/4}, e^{5i\pi/4}, e^{7i\pi/4}$$

3. Compute $\sqrt[4]{i}$ and $\sqrt[4]{-i}$.

Let $z = \sqrt[4]{i}$ and $z = re^{i\theta}$. Then $r^4 e^{4i\theta} = i = e^{i\pi/2}$.

$$\begin{aligned} r^4 &= 1 \\ \theta &= \frac{\pi}{8} \end{aligned}$$

so $z = e^{i\pi/8}$. Now, let $z = \sqrt[4]{-i}$. Then $r^4 e^{4i\theta} = e^{-i\pi/2}$ so $z = e^{-i\pi/8}$.

4. Solve the quadratic equation

$$z^2 + (\alpha + i\beta)z + \gamma + i\delta = 0.$$

The quadratic equation is $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2}$. For the complex polynomial, we have

$$z = \frac{-\alpha - \beta i \pm \sqrt{\alpha^2 - \beta^2 - 4\gamma + i(2\alpha\beta - 4\delta)}}{2}$$

Let $a + bi = \sqrt{\alpha^2 - \beta^2 - 4\gamma + i(2\alpha\beta - 4\delta)}$. Then

$$z = \frac{-\alpha - \beta \pm (a + bi)}{2}$$

1.1.3 Justification

1. Show that the system of all matrices of the special form

$$\begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix},$$

combined by matrix addition and matrix multiplication, is isomorphic to the field of complex numbers.

2. Show that the complex number system can be thought of as the field of all polynomials with real coefficients modulo the irreducible polynomial $x^2 + 1$.

1.1.4 Conjugation, Absolute Value

1. Verify by calculation the values of

$$\frac{z}{z^2 + 1}$$

for $z = x + iy$ and $\bar{z} = x - iy$ are conjugate.

For z , we have that $z^2 = x^2 - y^2 + 2xyi$.

$$\begin{aligned}\frac{z}{z^2 + 1} &= \frac{x + iy}{x^2 - y^2 + 1 + 2xyi} \\ &= \frac{x + iy}{x^2 - y^2 + 1 + 2xyi} \cdot \frac{x^2 - y^2 + 1 - 2xyi}{x^2 - y^2 + 1 - 2xyi} \\ &= \frac{x(x^2 - y^2 + 1) + 2xy^2 + iy(x^2 - y^2 + 1 - 2x^2)}{(x^2 - y^2 + 1)^2 + 4x^2y^2}\end{aligned}\quad (1.3)$$

For \bar{z} , we have that $\bar{z}^2 = x^2 - y^2 - 2xyi$.

$$\begin{aligned}\frac{\bar{z}}{\bar{z}^2 + 1} &= \frac{x - iy}{x^2 - y^2 + 1 - 2xyi} \\ &= \frac{x - iy}{x^2 - y^2 + 1 - 2xyi} \cdot \frac{x^2 - y^2 + 1 + 2xyi}{x^2 - y^2 + 1 + 2xyi} \\ &= \frac{x(x^2 - y^2 + 1) + 2xy^2 - iy(x^2 - y^2 + 1 - 2x^2)}{(x^2 - y^2 + 1)^2 + 4x^2y^2}\end{aligned}\quad (1.4)$$

Therefore, we have that equations (1.3) and (1.4) are conjugates.

2. Find the absolute value (modulus) of

$$-2i(3 + i)(2 + 4i)(1 + i) \quad \text{and} \quad \frac{(3 + 4i)(-1 + 2i)}{(-1 - i)(3 - i)}.$$

When we expand the first problem, we have that

$$z_1 = -2i(3 + i)(2 + 4i)(1 + i) = 32 + 24i$$

so

$$|z_1| = \sqrt{32^2 + 24^2} = 40.$$

For the second problem, we have that

$$z_2 = \frac{(3 + 4i)(-1 + 2i)}{(-1 - i)(3 - i)} = 2 - \frac{3}{2}i$$

so

$$|z_2| = \sqrt{4 + 9/4} = \frac{5}{2}.$$

3. Prove that

$$\left| \frac{a - b}{1 - \bar{a}b} \right| = 1$$

if either $|a| = 1$ or $|b| = 1$. What exception must be made if $|a| = |b| = 1$?

Recall that $|z|^2 = z\bar{z}$.

$$\begin{aligned}1^2 &= \left| \frac{a - b}{1 - \bar{a}b} \right|^2 \\ 1 &= \left(\frac{a - b}{1 - \bar{a}b} \right) \left(\overline{\frac{a - b}{1 - \bar{a}b}} \right) \\ &= \left(\frac{a - b}{1 - \bar{a}b} \right) \left(\frac{\bar{a} - \bar{b}}{1 - a\bar{b}} \right) \\ &= \frac{a\bar{a} - a\bar{b} - \bar{a}b + b\bar{b}}{1 - \bar{a}b - a\bar{b} + a\bar{a}b\bar{b}}\end{aligned}\quad (1.5)$$

If $|a| = 1$, then $|a|^2 = a\bar{a} = 1$ and similarly for $|b|^2 = 1$. Then equation (1.5) becomes

$$\frac{1 - a\bar{b} - \bar{a}b + b\bar{b}}{1 - \bar{a}b - a\bar{b} + b\bar{b}} \quad \text{and} \quad \frac{1 - a\bar{b} - \bar{a}b + a\bar{a}}{1 - \bar{a}b - a\bar{b} + a\bar{a}}$$

respectively which is one. If $|a| = |b| = 1$, then $|a|^2 = |b|^2 = 1$ so equation (1.5) can be written as

$$\frac{2 - a\bar{b} - \bar{a}b}{2 - \bar{a}b - a\bar{b}}.$$

Therefore, we must have that $a\bar{b} + \bar{a}b \neq 2$.

4. Find the conditions under which the equation $az + b\bar{z} + c = 0$ in one complex unknown has exactly one solution, and compute that solution.

Let $z = x + iy$. Then $az + b\bar{z} + c = a(x + iy) + b(x - iy) + c = 0$.

$$(a + b)x + c = 0 \quad (1.6a)$$

$$(a - b)y = 0 \quad (1.6b)$$

Lets consider equation (1.6b). We either have that $a = b$ or $y = 0$. If $a = b$, then WLOG equation (1.6a) can be written as

$$x = \frac{-c}{2a}$$

and $y \in \mathbb{R}$. For fixed a, b, c , we have infinitely many solutions when $a = b$ since $z = \frac{-c}{2a} + iy$ for $y \in \mathbb{R}$. If $y = 0$, then equation (1.6a) can be written as

$$x = \frac{-c}{a + b}.$$

Therefore, $z = x$ and we have only one solution.

5. Prove that Lagrange's identity in the complex form

$$\left| \sum_{i=1}^n a_i b_i \right|^2 = \sum_{i=1}^n |a_i|^2 \sum_{i=1}^n |b_i|^2 - \sum_{1 \leq i < j \leq n} |a_i \bar{b}_j - a_j \bar{b}_i|^2.$$

Let's consider

$$\left| \sum_{i=1}^n a_i b_i \right|^2 + \sum_{1 \leq i < j \leq n} |a_i \bar{b}_j - a_j \bar{b}_i|^2 = \sum_{i=1}^n |a_i|^2 \sum_{i=1}^n |b_i|^2.$$

Then we can write the lefthand side as

$$\begin{aligned} \left| \sum_{i=1}^n a_i b_i \right|^2 + \sum_{1 \leq i < j \leq n} |a_i \bar{b}_j - a_j \bar{b}_i|^2 &= \sum_{i=1}^n a_i b_i \sum_{j=1}^n \bar{a}_j \bar{b}_j + \sum_{1 \leq i < j \leq n} (a_i \bar{b}_j - a_j \bar{b}_i)(\bar{a}_i b_j - \bar{a}_j b_i) \\ &= \sum_{i,j=1}^n a_i b_i \bar{a}_j \bar{b}_j + \sum_{i < j} (|a_i|^2 |b_j|^2 + |a_j|^2 |b_i|^2) - \sum_{i < j} (a_i \bar{a}_j b_i \bar{b}_j + \bar{a}_i a_j \bar{b}_i b_j) \\ &= \sum_{i=j=1}^n |a_i|^2 |b_i|^2 + \sum_{i \neq j} a_i b_i \bar{a}_j \bar{b}_j + \sum_{i < j} (|a_i|^2 |b_j|^2 + |a_j|^2 |b_i|^2) \\ &\quad - \sum_{i < j} (a_i \bar{a}_j b_i \bar{b}_j + \bar{a}_i a_j \bar{b}_i b_j) \end{aligned}$$

For $i \neq j$, $\sum_{i \neq j} a_i b_i \bar{a}_j \bar{b}_j - \sum_{i < j} (a_i \bar{a}_j b_i \bar{b}_j + \bar{a}_i a_j \bar{b}_i b_j) = 0$. Thus, we now have

$$\sum_{i=1}^n |a_i|^2 |b_i|^2 + \sum_{i < j} (|a_i|^2 |b_j|^2 + |a_j|^2 |b_i|^2).$$

When the indicies of both series on the right hand side coincide,

$$\sum_{i=1}^n |a_i|^2 \sum_{i=1}^n |b_i|^2 = \sum_{i=1}^n |a_i|^2 |b_i|^2. \quad (1.7)$$

That is, both a_i and b_i index together on the left of side equation (1.7). When a_i and b_i dont index together on the left side of equation (1.7),

$$\sum_{i=1}^n |a_i|^2 \sum_{i=1}^n |b_i|^2 = \sum_{i < j} (|a_i|^2 |b_j|^2 + |a_j|^2 |b_i|^2)$$

as was needed to be shown.

1.1.5 Inequalities

1. Prove that

$$\left| \frac{a-b}{1-\bar{a}b} \right| < 1$$

if $|a| < 1$ and $|b| < 1$.

From the properties of the modulus, we have that

$$\begin{aligned} \left| \frac{a-b}{1-\bar{a}b} \right| &= \frac{|a-b|}{|1-\bar{a}b|} \\ &= \frac{|a-b|^2}{|1-\bar{a}b|^2} \end{aligned} \tag{1.8}$$

$$\begin{aligned} &= \frac{(a-b)(\bar{a}-\bar{b})}{(1-\bar{a}b)(1-a\bar{b})} \\ &= \frac{|a|^2 + |b|^2 - a\bar{b} - \bar{a}b}{1 + |a|^2|b|^2 - \bar{a}b - a\bar{b}} \\ &< \frac{2 - a\bar{b} - \bar{a}b}{2 - \bar{a}b - a\bar{b}} \\ &= 1 \end{aligned} \tag{1.9}$$

From equations (1.8) and (1.9), we have

$$\begin{aligned} \frac{|a-b|^2}{|1-\bar{a}b|^2} &< 1 \\ \frac{|a-b|}{|1-\bar{a}b|} &< 1 \end{aligned}$$

2. Prove Cauchy's inequality by induction.

Cauchy's inequality is

$$|a_1b_1 + \cdots + a_nb_n|^2 \leq (|a_1|^2 + \cdots + |a_n|^2)(|b_1|^2 + \cdots + |b_n|^2)$$

which can be written more compactly as

$$\left| \sum_{i=1}^n a_i b_i \right|^2 \leq \sum_{i=1}^n |a_i|^2 \sum_{i=1}^n |b_i|^2.$$

For the base case, $i = 1$, we have

$$|a_1b_1|^2 = (a_1b_1)(\bar{a}_1\bar{b}_1) = a_1\bar{a}_1b_1\bar{b}_1 = |a_1|^2|b_1|^2$$

so the base case is true. Now let the equality hold for all $k-1 \in \mathbb{Z}$ where $k-1 \leq n$. That is, we assume that

$$\left| \sum_{i=1}^{k-1} a_i b_i \right|^2 \leq \sum_{i=1}^{k-1} |a_i|^2 \sum_{i=1}^{k-1} |b_i|^2$$

to be true.

$$\begin{aligned} \left| \sum_{i=1}^{k-1} a_i b_i \right|^2 + |a_k b_k|^2 &\leq \sum_{i=1}^{k-1} |a_i|^2 \sum_{i=1}^{k-1} |b_i|^2 + |a_k b_k|^2 \\ \left| \sum_{i=1}^k a_i b_i \right|^2 &\leq \sum_{i=1}^{k-1} |a_i|^2 \sum_{i=1}^{k-1} |b_i|^2 + (a_k b_k)(\bar{a}_k \bar{b}_k) \\ &= \sum_{i=1}^{k-1} |a_i|^2 \sum_{i=1}^{k-1} |b_i|^2 + |a_k|^2 |b_k|^2 \\ &= \sum_{i=1}^k |a_i|^2 \sum_{i=1}^k |b_i|^2 \end{aligned}$$

Therefore, by the principal of mathematical induction, Cauchy's inequality is true for all $n \geq 1$ for $n \in \mathbb{Z}^+$.

3. If $|a_i| < 1$, $\lambda_i \geq 0$ for $i = 1, \dots, n$ and $\lambda_1 + \lambda_2 + \dots + \lambda_n = 1$, show that

$$|\lambda_1 a_1 + \lambda_2 a_2 + \dots + \lambda_n a_n| < 1.$$

Since $\sum_{i=1}^n \lambda_i = 1$ and $\lambda_i \geq 0$, $0 \leq \lambda_i < 1$. By the triangle inequality,

$$\begin{aligned} |\lambda_1 a_1 + \lambda_2 a_2 + \dots + \lambda_n a_n| &\leq |\lambda_1| |a_1| + \dots + |\lambda_n| |a_n| \\ &< \sum_{i=1}^n \lambda_i \\ &= 1 \end{aligned}$$

4. Show that there are complex numbers z satisfying

$$|z - a| + |z + a| = 2|c|$$

if and only if $|a| \leq |c|$. If this condition is fulfilled, what are the smallest and largest values $|z|$?

By the triangle inequality,

$$|z - a| + |z + a| \geq |(z - a) - (z + a)| = 2|a|$$

so

$$\begin{aligned} 2|c| &= |z - a| + |z + a| \\ &\geq |(z - a) - (z + a)| \\ &= 2|a| \end{aligned}$$

Thus, $|c| \geq |a|$. If $|a| \leq |c|$, then let $z = |c| \frac{a}{|a|}$.

$$\begin{aligned} |z - a| + |z + a| &\geq |(z - a) + (z + a)| \\ &= 2|c| \end{aligned}$$

since $\frac{a}{|a|}$ is a unit vector.

$$= 2|c|$$

Thus, $2|c| \geq 2|c|$ which is equality.

$$\begin{aligned} 2|c| &= |z + a| + |z - a| \\ 4|c|^2 &= (|z + a| + |z - a|)^2 \\ &= 2(|z|^2 + |a|^2) \\ &\leq 4(|z|^2 + |a|^2) \\ |c|^2 &\leq |z|^2 + |a|^2 \\ \sqrt{|c|^2 - |a|^2} &\leq |z| \end{aligned}$$

1.2 The Geometric Representation of Complex Numbers

1.2.1 Geometric Addition and Multiplication

- Find the symmetric points of a with respect to the lines which bisect the angles between the coordinate axes.
- Prove that the points a_1, a_2, a_3 are vertices of an equilateral triangle if and only if $a_1^2 + a_2^2 + a_3^2 = a_1 a_2 + a_2 a_3 + a_1 a_3$.
- Suppose that a and b are two vertices of a square. Find the two other vertices in all possible cases.
- Find the center and the radius of the circle which circumscribes the triangle with vertices a_1, a_2, a_3 . Express the result in symmetric form.

1.2.2 The Binomial Equation

1. Express $\cos(3\varphi)$, $\cos(4\varphi)$, and $\sin(5\varphi)$ in terms of $\cos(\varphi)$ and $\sin(\varphi)$.

For these problems, the sum addition identities will be employed; that is,

$$\begin{aligned}\cos(\alpha \pm \beta) &= \cos(\alpha) \cos(\beta) \mp \sin(\alpha) \sin(\beta) \\ \sin(\alpha \pm \beta) &= \sin(\alpha) \cos(\beta) \pm \sin(\beta) \cos(\alpha)\end{aligned}$$

We can write $\cos(3\varphi)$ as $\cos(2\varphi + \varphi)$ so

$$\begin{aligned}\cos(3\varphi) &= \cos(2\varphi + \varphi) \\ &= \cos(2\varphi) \cos(\varphi) - \sin(2\varphi) \sin(\varphi) \\ &= [\cos^2(\varphi) - \sin^2(\varphi)] \cos(\varphi) - 2 \sin(\varphi) \cos(\varphi) \sin(\varphi) \\ &= \cos^3(\varphi) - 3 \sin^2(\varphi) \cos(\varphi)\end{aligned}$$

For $\cos(4\varphi)$, we have

$$\begin{aligned}\cos(4\varphi) &= \cos(2\varphi) \cos(2\varphi) - \sin(2\varphi) \sin(2\varphi) \\ &= [\cos^2(\varphi) - \sin^2(\varphi)]^2 - 4 \sin^2(\varphi) \cos^2(\varphi) \\ &= \cos^4(\varphi) + \sin^4(\varphi) - 6 \sin^2(\varphi) \cos^2(\varphi)\end{aligned}$$

For $\sin(5\varphi)$, we have

$$\begin{aligned}\sin(5\varphi) &= \sin(4\varphi) \cos(\varphi) + \sin(\varphi) \cos(4\varphi) \\ &= 2 \sin(2\varphi) \cos(2\varphi) [\cos^2(\varphi) - \sin^2(\varphi)] \cos(\varphi) + \sin^5(\varphi) + \sin(\varphi) \cos^4(\varphi) - 6 \sin^3(\varphi) \cos^2(\varphi) \\ &= 5 \sin(\varphi) \cos^4(\varphi) - 10 \sin^3(\varphi) \cos^2(\varphi) + \sin^5(\varphi)\end{aligned}$$

2. Simplify $1 + \cos(\varphi) + \cos(2\varphi) + \cdots + \cos(n\varphi)$ and $\sin(\varphi) + \cdots + \sin(n\varphi)$.

Instead of considering the two separate series, we will consider the series

$$\begin{aligned}1 + \cos(\varphi) + i \sin(\varphi) + \cdots + \cos(n\varphi) + i \sin(n\varphi) &= 1 + e^{i\varphi} + e^{2i\varphi} + \cdots + e^{ni\varphi} \\ &= \sum_{k=0}^n e^{ki\varphi}\end{aligned}$$

Recall that $\sum_{k=0}^{n-1} r^k = \frac{1-r^n}{1-r}$. So

$$\begin{aligned}&= \frac{1 - e^{i\varphi(n+1)}}{1 - e^{i\varphi}} \\ &= \frac{e^{i\varphi(n+1)} - 1}{e^{i\varphi} - 1}\end{aligned}\tag{1.10}$$

Note that $\sin(\frac{\varphi}{2}) = \frac{e^{i\varphi/2} - e^{-i\varphi/2}}{2i}$ so $2ie^{i\varphi/2} \sin(\frac{\varphi}{2}) = e^{i\varphi} - 1$. We can now write equation (1.10) as

$$\begin{aligned}\sum_{k=0}^n e^{ki\varphi} &= \frac{e^{i\varphi(n+1)/2} \sin(\frac{\varphi(n+1)}{2})}{e^{i\varphi/2} \sin(\frac{\varphi}{2})} \\ &= \frac{\sin(\frac{\varphi(n+1)}{2})}{\sin(\frac{\varphi}{2})} e^{in\varphi/2}\end{aligned}\tag{1.11}$$

By taking the real and imaginary parts of equation (1.11), we get the series for $\sum_{k=0}^n \cos(k\varphi)$ and $\sum_{k=0}^n \sin(k\varphi)$, respectively.

$$\begin{aligned}\sum_{k=0}^n \cos(k\varphi) &= \frac{\sin(\frac{\varphi(n+1)}{2})}{\sin(\frac{\varphi}{2})} \cos(\frac{n\varphi}{2}) \\ \sum_{k=0}^n \sin(k\varphi) &= \frac{\sin(\frac{\varphi(n+1)}{2})}{\sin(\frac{\varphi}{2})} \sin(\frac{n\varphi}{2})\end{aligned}$$

3. Express the fifth and tenth roots of unity in algebraic form.

To find the roots of unity, we are looking to solve $z^n = 1$. Let $z = e^{i\theta}$ and $1 = e^{2ik\pi}$. Then $\theta = \frac{2k\pi}{n}$. For the fifth roots of unity, $n = 5$ and $k = 0, 1, \dots, 4$ so we have

$$\begin{aligned}\omega_0 &= e^0 = \cos(0) + i \sin(0) \\ &= 1 \\ \omega_1 &= e^{2\pi/5} = \cos\left(\frac{2\pi}{5}\right) + i \sin\left(\frac{2\pi}{5}\right) \\ \omega_2 &= e^{4\pi/5} = \cos\left(\frac{4\pi}{5}\right) + i \sin\left(\frac{4\pi}{5}\right) \\ \omega_3 &= e^{6\pi/5} = \cos\left(\frac{6\pi}{5}\right) + i \sin\left(\frac{6\pi}{5}\right) \\ \omega_4 &= e^{8\pi/5} = \cos\left(\frac{8\pi}{5}\right) + i \sin\left(\frac{8\pi}{5}\right)\end{aligned}$$

Now we can plot the roots of unity on the unit circle.




Figure 1.1: The fifth roots of unity.

For the tenth roots of unity, $n = 10$ and $k = 0, 1, \dots, 9$ so we have

$$\begin{aligned}\omega_0 &= e^0 = \cos(0) + i \sin(0) \\ &= 1 \\ \omega_1 &= e^{2\pi/10} = \cos\left(\frac{\pi}{5}\right) + i \sin\left(\frac{\pi}{5}\right) \\ \omega_2 &= e^{4\pi/10} = \cos\left(\frac{2\pi}{5}\right) + i \sin\left(\frac{2\pi}{5}\right) \\ \omega_3 &= e^{6\pi/10} = \cos\left(\frac{3\pi}{5}\right) + i \sin\left(\frac{3\pi}{5}\right) \\ \omega_4 &= e^{8\pi/10} = \cos\left(\frac{4\pi}{5}\right) + i \sin\left(\frac{4\pi}{5}\right) \\ \omega_5 &= e^{10\pi/10} = \cos(\pi) + i \sin(\pi) \\ &= -1 \\ \omega_6 &= e^{12\pi/10} = \cos\left(\frac{6\pi}{5}\right) + i \sin\left(\frac{6\pi}{5}\right) \\ \omega_7 &= e^{14\pi/10} = \cos\left(\frac{7\pi}{5}\right) + i \sin\left(\frac{7\pi}{5}\right) \\ \omega_8 &= e^{16\pi/10} = \cos\left(\frac{8\pi}{5}\right) + i \sin\left(\frac{8\pi}{5}\right) \\ \omega_9 &= e^{18\pi/10} = \cos\left(\frac{9\pi}{5}\right) + i \sin\left(\frac{9\pi}{5}\right)\end{aligned}$$

Now we can plot the roots of unity on the unit circle.




Figure 1.2: The tenth roots of unity.

4. If ω is given by $\omega = \cos\left(\frac{2\pi}{n}\right) + i \sin\left(\frac{2\pi}{n}\right)$, prove that

$$1 + \omega^h + \omega^{2h} + \dots + \omega^{(n-1)h} = 0$$

for any integer h which is not a multiple of n .

Let $\omega = \cos\left(\frac{2\pi}{n}\right) + i \sin\left(\frac{2\pi}{n}\right)$ be written in exponential form as $\omega = e^{2\pi i/n}$. Then the series can be written as

$$\sum_{k=0}^{n-1} (e^{2\pi i h/n})^k = \frac{e^{2i h \pi} - 1}{e^{2h i \pi/n} - 1}.$$

Since h is an integer, $e^{2i h \pi} = 1$; therefore, the series zero.

5. What is the value of

$$1 - \omega^h + \omega^{2h} - \dots + (-1)^{n-1} \omega^{(n-1)h}?$$

We can represent this series similarly as

$$\sum_{k=0}^{n-1} (-e^{2\pi i h/n})^k = \frac{(-1)^n e^{2i h \pi} - 1}{-e^{2h i \pi/n} - 1} = \frac{1 + (-1)^{n+1} e^{2i h \pi}}{1 + e^{2h i \pi/n}}.$$

Again, since h is an integer, we have that $e^{2i h \pi} = 1$ which leaves us with

$$\frac{1 + (-1)^{n+1}}{1 + e^{2h i \pi/n}} = \begin{cases} 0, & \text{if } n \text{ is even} \\ \frac{2}{1 + e^{2h i \pi/n}}, & \text{if } n \text{ is odd} \end{cases}$$

1.2.3 Analytic Geometry

1. When does $az + b\bar{z} + c = 0$ represent a line?
2. Write the equation of an ellipse, hyperbola, parabola in complex form.

For $x, y, h, k, a, b \in \mathbb{R}$ such that $a, b \neq 0$, we define a real ellipse as

$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1.$$

Let $z = \frac{x}{a} + i \frac{y}{b}$ and $z_0 = \frac{h}{a} + i \frac{k}{b}$. If we expand the equation for an ellipse, we have

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{h^2}{a^2} + \frac{k^2}{b^2} - \frac{2xh}{a^2} - \frac{2yk}{b^2} = 1.$$

Notice that $|z|^2 = \frac{x^2}{a^2} + \frac{y^2}{b^2}$ and $|z_0|^2 = \frac{h^2}{a^2} + \frac{k^2}{b^2}$. Now, let's write the ellipse as

$$|z|^2 + |z_0|^2 - \frac{2xh}{a^2} - \frac{2yk}{b^2} + \frac{yh}{ab}i - \frac{yh}{ab}i + \frac{xk}{ab}i - \frac{xk}{ab}i = |z|^2 + |z_0|^2 - \bar{z}z_0 - z\bar{z}_0 = 1.$$

Thus, the equation of an ellipse in the complex plane is

$$(z - z_0)(\bar{z} - \bar{z}_0) = |z - z_0|^2 = 1 \Rightarrow |z - z_0| = 1$$

where z and z_0 are defined above. Additionally, the standard form of an ellipse in the complex plane is of the form

$$|z - a| + |z - b| = c$$

where $c > |a - b|$. Let a and b be the foci of a hyperbola. Then when the magnitude of the difference of z and the foci is a constant, we will have a hyperbola.

$$||z - a| - |z - b|| = c$$

3. Prove that the diagonals of a parallelogram bisect each other and that the diagonals of a rhombus are orthogonal.
4. Prove analytically that the midpoints of parallel chords to a circle lie on a diameter perpendicular to the chords.
5. Show that all circles that pass through a and $1/\bar{a}$ intersect the circle $|z| = 1$ at right angles.

1.2.4 The Spherical Representation

1. Show that z and z' correspond to diametrically opposite points on the Riemann sphere if and only if $z\bar{z}' = -1$.
2. A cube has its vertices on the sphere S and its edges parallel to the coordinate axes. Find the stereographic projections of the vertices.
3. Same problem for a regular tetrahedron in general position.
4. Let Z, Z' denote the stereographic projections of z, z' , and let N be the north pole. Show that the triangles NZZ' and Nzz' are similar, and use this to derive

$$d(z, z') = \frac{2|z - z'|}{\sqrt{1 + |z|^2}\sqrt{1 + |z'|^2}}.$$

5. Find the radius of the spherical image of the circle in the plane whose center is a and radius R .

2 Complex Functions

2.1 Introduction to the Concept of Analytical Function

2.1.1 Limits and Continuity

2.1.2 Analytic Functions

1. If $g(w)$ and $f(z)$ are analytic functions, show that $g(f(z))$ is also analytic.
2. Verify Cauchy-Riemann's equations for the function z^2 and z^3 .

Let $z = x + iy$. Then $z^2 = x^2 - y^2 + 2xyi$ and $z^3 = x^3 - 3xy^2 + i(3x^2y - y^3)$. For $f(z) = z^2$, the Cauchy-Riemann equations are

$$\begin{aligned}u_x &= 2x & v_y &= 2x \\u_y &= -2y & -v_x &= -2y\end{aligned}$$

Thus, the Cauchy-Riemann equation satisfied for $f(z) = z^2$. For $f(z) = z^3$, the Cauchy-Riemann equations are

$$\begin{aligned}u_x &= 3x^2 - 3y^2 & v_y &= 3x^2 - 3y^2 \\u_y &= -6xy & -v_x &= -6xy\end{aligned}$$

Thus, the Cauchy-Riemann equation satisfied for $f(z) = z^3$.

3. Find the most general harmonic polynomial of the form $ax^3 + bx^2y + cxy^2 + dy^3$. Determine the conjugate harmonic function and the corresponding analytic function by integration and by the formal method.

In order to be harmonic, $u(x, y) = ax^3 + bx^2y + cxy^2 + dy^3$ has to satisfy $\nabla^2 u = 0$ so

$$u_{xx} + u_{yy} = (3a + c)x + (3d + b)y = 0.$$

Thus, $3a = -c$ and $3d = -b$ so

$$u(x, y) = ax^3 - 3axy^2 - 3dx^2y + dy^3.$$

To find the harmonic conjugate $v(x, y)$, we need to look at the Cauchy-Riemann equations. By the Cauchy-Riemann equations,

$$u_x = 3ax^2 - 3ay^2 - 6dxy = v_y.$$

Then we can integrate with respect to y to find $v(x, y)$.

$$v(x, y) = \int (3ax^2 - 3ay^2 - 6dxy) dy = 3ax^2y - ay^3 - 3dxy^2 + g(x)$$

Using the second Cauchy-Riemann, we have

$$v_x = 6axy - 3dy^2 + g'(x) = -u_y = 3dx^2 + 6axy - 3dy^2$$

so $g'(x) = 3dx^2$. Then $g(x) = dx^3 + C$ and

$$v(x, y) = 3ax^2y - ay^3 - 3dxy^2 + dx^3 + C.$$

4. Show that an analytic function cannot have a constant absolute value without reducing to a constant.

Let $f = u(x, y) + iv(x, y)$. Then the modulus of f is $|f| = \sqrt{u^2 + v^2}$. If the modulus of f is constant, then $u^2 + v^2 = c$ for some constant c . If $c = 0$, then $f = 0$ which is constant. Suppose $c \neq 0$. By taking the derivative with respect to x and y , we have

$$\begin{aligned} 0 &= \frac{\partial}{\partial x}(u^2 + v^2) \\ &= 2uu_x + 2vv_x \\ &= uu_x + vv_x \\ 0 &= \frac{\partial}{\partial y}(u^2 + v^2) \\ &= uu_y + vv_y \end{aligned}$$

Since f is analytic, f satisfies the Cauchy-Riemann. That is, $u_x = v_y$ and $u_y = -v_x$.

$$uv_y + vv_x = 0 \quad (2.1a)$$

$$-uv_x + vv_y = 0 \quad (2.1b)$$

Setting equation (2.1a) equal to equation (2.1b), we have

$$v_x(u + v) + v_y(u - v) = 0.$$

Now, either v_x and v_y are zero, v_x and $u - v$ are zero, v_y and $u + v$ are zero, or $u + v$ and $u - v$ are zero. If $v_x = v_y = 0$, then f is constant. If $v_x = 0$ and $u - v = 0$, then $u_y = 0$ and $u = v$. Since $u = v$ and $v_x = 0$, then so does $u_x = 0$ and it also follows that $v_y = 0$; thus, f is a constant. By the same argument, f is a constant when $v_y = 0$ and $u + v = 0$. If $u + v = 0$ and $u - v = 0$, then $u = \pm v$ so $u = v = 0$ and f is a constant.

5. Prove rigorously that the functions $f(z)$ and $\overline{f(\bar{z})}$ are simultaneously analytic.

Let $g(z) = \overline{f(\bar{z})}$ and suppose f is analytic. Then $g'(z)$ is

$$\begin{aligned} g'(z) &= \lim_{\Delta z \rightarrow 0} \frac{g(z + \Delta z) - g(z)}{\Delta z} \\ &= \lim_{\Delta z \rightarrow 0} \frac{\overline{f(\bar{z} + \overline{\Delta z})} - \overline{f(\bar{z})}}{\Delta z} \\ &= \lim_{\Delta z \rightarrow 0} \left[\overline{\frac{f(\bar{z} + \overline{\Delta z}) - f(\bar{z})}{\overline{\Delta z}}} \right] \end{aligned}$$

Since conjugation is continuous, we can move the limit inside the conjugation.

$$\begin{aligned} &= \overline{\lim_{\Delta z \rightarrow 0} \frac{f(\bar{z} + \overline{\Delta z}) - f(\bar{z})}{\overline{\Delta z}}} \\ &= \overline{f'(\bar{z})} \end{aligned}$$

Thus, g is differentiable with derivative $\overline{f'(\bar{z})}$. Suppose $\overline{f(\bar{z})}$ is analytic and let $\overline{g(\bar{z})} = f(z)$. Then by the same argument, f is differentiable with derivative $\overline{g'(\bar{z})}$. Therefore, $f(z)$ and $\overline{f(\bar{z})}$ are simultaneously analytic.

We could also use the Cauchy-Riemann equations. Let $f(z) = u(x, y) + iv(x, y)$ where $z = x + iy$ so $\bar{z} = x - iy$. Then $\overline{f(\bar{z})} = \alpha(x, y) - i\beta(x, y)$ where $\alpha(x, y) = u(x, -y)$ and $\beta(x, y) = v(x, -y)$. In order for both to be analytic, they both need to satisfy the Cauchy-Riemann equations. That is, $u_x = v_y$, $u_y = -v_x$, $\alpha_x = \beta_y$ and $\alpha_y = -\beta_x$.

$$\begin{aligned} u_x(x, y) &= v_y(x, y) \\ u_y(x, y) &= -v_x(x, y) \\ \alpha_x(x, y) &= u_x(x, -y) \\ \alpha_y(x, y) &= -u_y(x, -y) \end{aligned}$$

$$-\beta_x(x, y) = v_x(x, -y)$$

$$\beta_y(x, y) = v_y(x, -y)$$

Suppose that $\overline{f(\bar{z})}$ satisfies the Cauchy-Riemann equations. Then $\alpha_x = u_x(x, -y) = v_y(x, -y) = \beta_y$ and $\alpha_y = -u_y(x, -y) = v_x(x, -y) = -\beta_x$. Therefore,

$$u_x(x, -y) = v_y(x, -y)$$

$$u_y(x, -y) = -v_x(x, -y)$$

which means $f(\bar{z})$ satisfies the Cauchy-Riemann equations. Now, recall that $|z| = |\bar{z}|$. Since $f(\bar{z})$ satisfies the Cauchy-Riemann equations, for an $\epsilon > 0$ there exists a $\delta > 0$ such that when $0 < |\Delta z| < \delta$, $|f(\bar{z}) - \bar{z}_0| = |f(z) - z_0| < \epsilon$. Thus, $\lim_{\Delta z \rightarrow 0} f(z) = z_0$ so $f(z)$ is analytic if $\overline{f(\bar{z})}$ is analytic.

6. Prove that the functions $u(z)$ and $u(\bar{z})$ are simultaneously harmonic.

Since u is the real part of $f(z)$, $u(z) = u(x, y)$ where $z = x + iy$. Suppose $u(z)$ is harmonic. Then $u(z)$ satisfies Laplace equation.

$$\nabla^2 u(z) = u_{xx} + u_{yy} = 0$$

Now, $u(\bar{z}) = u(x, -y)$ where $\frac{\partial^2}{\partial x^2} u(\bar{z}) = u_{xx}$ and $\frac{\partial^2}{\partial y^2} u(\bar{z}) = u_{yy}$ so

$$\nabla^2 u(\bar{z}) = u_{xx} + u_{yy} = 0.$$

Since $u(z)$ is harmonic, $u_{xx} + u_{yy} = 0$ so it follows that $u(\bar{z})$ is harmonic as well.

7. Show that a harmonic function satisfies the formal differential equation

$$\frac{\partial^2 u}{\partial z \partial \bar{z}} = 0.$$

Let u be a harmonic. Then $\nabla^2 u = 0$.

$$\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) \quad (2.2a)$$

$$\frac{\partial}{\partial z} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \quad (2.2b)$$

From equation (2.2a), we have

$$\frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) u = \frac{1}{2} (u_x + i u_y).$$

Then we have

$$\frac{\partial^2 u}{\partial z \partial \bar{z}} = \frac{1}{4} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (u_x + i u_y) = \frac{1}{4} [u_{xx} + u_{yy} + i(u_{yx} - u_{xy})]$$

Since u is a solution to the Laplace equation, u has continuous first and second derivatives. That is, $u \in C^2$ at a minimum. By Schwarz's theorem, $u_{xy} = u_{yx}$ so

$$\frac{\partial^2 u}{\partial z \partial \bar{z}} = 0.$$

Schwarz's theorem states that if f is a function of two variables such that f_{xy} and f_{yx} both exist and are continuous at some point (x_0, y_0) , then $f_{xy} = f_{yx}$.

2.1.3 Polynomials

2.1.4 Rational Functions

1. Use the method of the text to develop

$$\frac{z^4}{z^3 - 1} \quad \text{and} \quad \frac{1}{z(z+1)^2(z+2)^3}$$

in partial fractions.

Let $R(z) = \frac{z^4}{z^3-1}$. Then we need to find $G(z)$ and $H(z)$ such that $G(z) + H(z) = R(z)$. $G(z)$ is the singular part of $R(z)$ and the degree of $G(z)$ is order of the pole at infinity. Dividing z^4 by $z^3 - 1$, we get

$$R(z) = z + \frac{z}{z^3-1}$$

so $G(z) = z$ and $H(z) = \frac{z}{z^3-1}$. The order of the pole at infinity is one.

2. If Q is a polynomial with distinct roots $\alpha_1, \dots, \alpha_n$, and if P is a polynomial of degree $< n$, show that

$$\frac{P(z)}{Q(z)} = \sum_{k=1}^n \frac{P(\alpha_k)}{Q'(\alpha_k)(z - \alpha_k)}.$$

Let's multiply by $Q(z)$. We then have

$$P(z) = \sum_{k=1}^n \frac{P(\alpha_k)}{Q'(\alpha_k)(z - \alpha_k)} Q(z)$$

which are both polynomials of degree less than n and agreeing at $z = \alpha_k$.

3. Use the formula in the preceding exercise to prove that there exists a unique polynomial P of degree $< n$ with given values c_k at the points α_k (Lagrange's interpolation polynomial).
4. What is the general form of a rational function which has absolute value 1 on the circle $|z| = 1$? In particular, how are the zeros and poles related to each other?
5. If a rational function is real on $|z| = 1$, how are the zeros and poles situated?
6. If $R(z)$ is a rational function of order n , how large and how small can the order of $R'(z)$ be?

2.2 Elementary Theory of Power Series

2.2.1 Sequences

2.2.2 Series

2.2.3 Uniform Convergence

1. Prove that a convergent sequence is bounded.

Let $\{a_n\}$ be a convergent sequence and $\lim_{n \rightarrow \infty} a_n = a$. Let $\epsilon = 1$. Then there exists an $n > N$ such that $|a_n - a| < 1$.

$$|a_n| = |a_n - a + a|$$

By the triangle inequality, we have

$$\begin{aligned} &\leq |a_n - a| + |a| \\ |a_n| - |a| &\leq |a_n - a| \end{aligned}$$

Therefore, we have that

$$\begin{aligned} |a_n| - |a| &\leq |a_n - a| < 1 \\ |a_n| &< 1 + |a| \end{aligned}$$

For all $n > N$, $|a_n| < 1 + |a|$ so let $A = \max\{1 + |a|, |a_1|, \dots, |a_N|\}$. Thus, $|a_n| < A$ for some finite A and hence $\{a_n\}$ is bounded by A .

2. If $\lim_{n \rightarrow \infty} z_n = A$, prove that

$$\lim_{n \rightarrow \infty} \frac{1}{n} (z_1 + z_2 + \dots + z_n) = A.$$

Given $\epsilon > 0$ there exists some $n > N$ such that

$$|z_n - A| < \frac{N\epsilon}{2}.$$

Now, since $\lim_{n \rightarrow \infty} z_n = A$ converges, it is Cauchy. Therefore, there exists $n, m > N$ such that

$$|z_m - z_n| < \frac{N\epsilon}{2}.$$

Repeating this we have that $|z_1 + \dots + z_n - nA|$ or $|1/n(z_1 + \dots + z_{n-1}) - A + (z_n - A)/n|$. For a fixed N , we can find n such that

$$\sum_{i=1}^{n-1} |z_i - A| < \frac{N\epsilon}{2}$$

We now have that

$$\begin{aligned} |1/n(z_1 + \dots + z_{n-1}) - A + (z_n - A)/n| &\leq \left| 1/n \sum_{i=1}^{n-1} (z_i - A) \right| + 1/n|z_n - A| \\ &\leq 1/n \sum_{i=1}^{n-1} |z_i - A| + 1/n|z_n - A| \\ &< 1/n \frac{N\epsilon}{2} + 1/n \frac{N\epsilon}{2} \\ &< \epsilon \end{aligned} \quad (2.3)$$

Equation (2.3) can be written as $|1/n(z_1 + \dots + z_n) - A| < \epsilon$ so

$$\lim_{n \rightarrow \infty} 1/n(z_1 + \dots + z_n) = A.$$

3. Show that the sum of an absolutely convergent series does not change if the terms are rearranged.

Let $\sum a_n$ be an absolutely convergent series and $\sum b_n$ be its rearrangement. Since $\sum a_n$ converges absolutely, for $\epsilon > 0$, there exists a $n > N$ such that $|s_n - A| < \epsilon/2$ where s_n is the n th partial sum. Let t_n be the n th partial sum of $\sum b_n$. Then for some $n > N$

$$\begin{aligned} |t_n - A| &= |t_n - s_n + s_n - A| \\ &\leq |t_n - s_n| + |s_n - A| \\ &< |t_n - s_n| + \frac{\epsilon}{2} \end{aligned}$$

Since $\sum a_n$ is absolutely convergent, $\sum_{k=n+1}^{\infty} |a_k|$ converges to zero. Let the remainder be r_n . Then for some $N > n, n_1$, $|r_n - 0| < \epsilon/2$. Let $M = \max\{k_1, k_2, \dots, k_N\}$. Then for some $n > M$, we have

$$|t_n - s_n| = \left| \sum_n^N a_n \right| \leq \sum |a_n| \leq \sum_{k=n+1}^{\infty} |a_n| = r_n < \frac{\epsilon}{2}$$

Thus, $|t_n - s_n| < \epsilon$ and a rearrangement of an absolutely convergent series does not change its sum.

4. Discuss completely the convergence and uniform convergence of the sequence $\{nz^n\}_{n=1}^{\infty}$.

Consider when $|z| < 1$. Then $z^n = \frac{1}{w^n}$ where $|w| > 1$. By the ratio test, we have

$$\lim_{n \rightarrow \infty} \left| \frac{(n+1)w^n}{nw^{n+1}} \right| = \frac{1}{|w|} \lim_{n \rightarrow \infty} \frac{n+1}{n} = \frac{1}{|w|}$$

In order for convergence, the ratio test has to be less than one.

$$\frac{1}{|w|} = |z|$$

which is less than one by our assumption so $\{nz^n\}$ converges absolutely in the disc less than one. Now, let's consider $|z| \geq 1$. By the ratio test, we get $\lim_{n \rightarrow \infty} |a_{n+1}/a_n| = |z| \geq 1$ by our assumption. When the

limit is one, we can draw no conclusion about convergence, but when the limit is greater than one, the sequence diverges. For $|z| < 1$, $\epsilon > 0$, and $n > N$, $|nz^n - 0| < \epsilon$ for uniform convergence. Take $z = 9/10$, $n = 100$, and $\epsilon = 0.001$. Then

$$\begin{aligned} |nz^n| &= n|z|^n < \epsilon \\ |z|^n &< \frac{\epsilon}{n} \\ 0.000026 &\not\leq 0.00001 \end{aligned}$$

Thus, the sequence is not uniformly convergent in the disc with radius less than one. Let's consider the closed disc $|z| \leq R$ where $R \in (0, 1)$. Now $|nz^n|$ is bounded above by a convergent geometric series, say $\sum r^n$ where $|r| < 1$. Then $|nz^n| < ar^n$ for $|z| \leq R$ and a a real constant. Let $M_n = ar^n$ where M_n is the M in the Weierstrass M-test. Thus, $\{nz^n\}$ is uniformly convergent in a closed disc less than one.

5. Discuss the uniform convergence of the series

$$\sum_{n=1}^{\infty} \frac{x}{n(1+nx^2)}$$

for real values of x .

By the AM-GM inequality, $(x+y)/2 \geq \sqrt{xy}$, we have

$$1 + nx^2 \geq 2|x|\sqrt{n}$$

or $\frac{1}{2|x|\sqrt{n}} \geq \frac{1}{1+nx^2}$. Let $f_n(x) = \frac{x}{n(1+nx^2)}$. Then

$$|f_n(x)| \leq \left| \frac{x}{2xn^{3/2}} \right| = \left| \frac{1}{2n^{3/2}} \right| = M_n$$

For a fixed x , $\sum |f_n(x)| \leq M_n < \infty$ so $\sum |f_n(x)|$ is absolutely convergent. Thus, $\sum f_n(x)$ is pointwise convergent to $f(x)$. Let $\epsilon > 0$ be given and $s_n = \sum_{k=1}^n f_k(x)$ be n th partial sum. Let $n > N$ such that

$$|f(x) - s_n| = \left| \sum_{k=1}^{\infty} f_k(x) - \sum_{k=1}^n f_k(x) \right| = \left| \sum_{k=n+1}^{\infty} f_k(x) \right| \leq \sum_{k=n+1}^{\infty} |f_k(x)|$$

Since $\sum M_k$ converges to some limit, for n sufficiently large, $\sum_{k=n+1}^{\infty} M_k < \epsilon$. Select N such that this is true. Then

$$|f(x) - s_n| \leq \sum_{k=n+1}^{\infty} |f_k(x)| \leq \sum_{k=n+1}^{\infty} M_k < \epsilon$$

Therefore, $\sum f_n(x)$ where $f_n(x) = \frac{x}{n(1+nx^2)}$ is uniformly convergent by the Weierstrass M-test.

6. If $U = u_1 + u_2 + \dots$, $V = v_1 + v_2 + \dots$ are convergent series, prove that $UV = u_1v_1 + (u_1v_2 + u_2v_1) + (u_1v_3 + u_2v_2 + u_3v_1) + \dots$ provided that at least one of the series is absolutely convergent. (It is easy if both series are absolutely convergent. Try to rearrange the proof so economically that the absolute convergence of the second series is not needed.)

2.2.4 Power Series

1. Expand $(1-z)^{-m}$, m a positive integer, in powers of z .

The Binomial theorem states that $(1+x)^n = \sum_{k=0}^{\infty} \binom{n}{k} x^k$. In our case, we have

$$(1-z)^{-m} = \sum_{k=0}^{\infty} \binom{-m}{k} (-z)^k \quad (2.4)$$

where $\binom{-m}{k} = (-1)^k \binom{m+k-1}{k}$. Then equation (2.4) can be written as

$$(1-z)^{-m} = \sum_{k=0}^{\infty} \binom{m+k-1}{k} z^k = 1 + mz + \frac{m(m+1)}{2!} z^2 + \dots$$

2. Expand $\frac{2z+3}{z+1}$ in powers of $z-1$. What is the radius of convergence?

Let's just consider $\frac{1}{z+1}$ for the moment.

$$\frac{1}{z+1} = \frac{1}{z-1+2} = \frac{1/2}{1 + \frac{z-1}{2}} = \frac{1}{2} \sum_{n=0}^{\infty} (-1)^n \left(\frac{z-1}{2}\right)^n$$

From the full expressing, we obtain

$$\frac{2z+3}{z+1} = \frac{2z+3}{2} \sum_{n=0}^{\infty} (-1)^n \left(\frac{z-1}{2}\right)^n.$$

The radius of convergence can be found by $1/R = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$. Therefore, the radius of convergence is

$$R = 1/\limsup_{n \rightarrow \infty} \sqrt[n]{\left|(-1)^n \frac{1}{2^n}\right|} = |2| = 2$$

3. Find the radius of convergence of the following power series:

$$\sum n^p z^n, \quad \sum \frac{z^n}{n!}, \quad \sum n! z^n, \quad \sum q^{n^2} z^n, \quad \sum z^{n!}$$

where $|q| < 1$.

For $\sum n^p z^n$, we can use the inverse of argument of the ratio test to determine the radius of convergence; that is,

$$R = \lim_{n \rightarrow \infty} \left| \frac{n^p}{(n+1)^p} \right| = \lim_{n \rightarrow \infty} \frac{n^p}{(n+1)^p} = 1$$

For $\sum \frac{z^n}{n!}$, we can use the fact that the sum is e^z which is entire or the method used previously. Since e^z is entire, the radius of convergence is $R = \infty$.

$$R = \lim_{n \rightarrow \infty} \left| \frac{(n+1)!}{n!} \right| = \lim_{n \rightarrow \infty} \frac{n!(n+1)}{n!} = \infty$$

For $\sum n! z^n$, we use the modified ratio test again.

$$R = \lim_{n \rightarrow \infty} \left| \frac{n!}{(n+1)!} \right| = 0$$

For $\sum q^{n^2} z^n$, we will use the ratio test.

$$R = \lim_{n \rightarrow \infty} \left| \frac{q^{n^2}}{q^{(n+1)^2}} \right| = \frac{1}{|q|} \lim_{n \rightarrow \infty} \left| \frac{1}{q^{2n}} \right| = \infty$$

For $\sum z^{n!}$, we will use the root test.

$$R = 1/\limsup_{n \rightarrow \infty} \sqrt[n]{|z^{(n-1)!}|^n} = 1/\limsup_{n \rightarrow \infty} |z|^{(n-1)!}$$

When $|z| < 1$, $R = \infty$, and when $|z| > 1$, $R = 0$.

4. If $\sum a_n z^n$ has a radius of convergence R , what is the radius of convergence of $\sum a_n z^{2n}$? of $\sum a_n^2 z^n$?

Since $\sum a_n z^n$ has a radius of convergence R ,

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|.$$

For $\sum a_n z^{2n} = z^2 \sum a_n z^n$, we have

$$|z|^2 \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = |z|^2 R$$

so the radius of convergence is \sqrt{R} . For $\sum a_n^2 z^n$, we have

$$\lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|^2 = R^2.$$

5. If $f(z) = \sum a_n z^n$, what is $\sum n^3 a_n z^n$?

Let's write out the first few terms of

$$\sum n^3 a_n z^n = a_1 z + 8a_2 z^2 + 27a_3 z^3 + 64a_4 z^4 + \dots$$

Let's consider the first three derivatives of $f(z)$.

$$\begin{aligned} f'(z) &= \sum n a_n z^{n-1} \\ z f'(z) &= \sum n a_n z^n \\ &= a_1 z + 2a_2 z^2 + 3a_3 z^3 + \dots \end{aligned} \quad (2.5)$$

$$\begin{aligned} f''(z) &= \sum n(n-1) a_n z^{n-2} \\ z^2 f''(z) &= \sum n(n-1) a_n z^n \\ &= 2a_2 z^2 + 6a_3 z^3 + 12a_4 z^4 + \dots \end{aligned} \quad (2.6)$$

$$\begin{aligned} f'''(z) &= \sum n(n-1)(n-2) a_n z^{n-3} \\ z^3 f'''(z) &= \sum n(n-1)(n-2) a_n z^n \\ &= 6a_3 z^3 + 24a_4 z^4 + 60a_5 z^5 + \dots \end{aligned} \quad (2.7)$$

If we add equations (2.5) to (2.7), we have

$$z f'(z) + z^2 f''(z) + z^3 f'''(z) = a_1 z + 4a_2 z^2 + 15a_3 z^3 + \dots \neq \sum n^3 a_n z^n$$

However, consider $3z^2 f''(z) = 6a_2 z^2 + 18a_3 z^3 + 36a_4 z^4 + \dots$. Then

$$z f'(z) + 3z^2 f''(z) + z^3 f'''(z) = a_1 z + 8a_2 z^2 + 27a_3 z^3 + 64a_4 z^4 + \dots = \sum n^3 a_n z^n.$$

6. If $\sum a_n z^n$ and $\sum b_n z^n$ have radii of convergence R_1 and R_2 , show that the radii of convergence of $\sum a_n b_n z^n$ is at least $R_1 R_2$.

Let $\epsilon > 0$ be given. Then there exists $n > N$ such that

$$|a_n|^{1/n} < 1/R_1 + \epsilon, \quad |b_n|^{1/n} < 1/R_2 + \epsilon$$

since $\limsup_{n \rightarrow \infty} |a_n|^{1/n} = 1/R_1$ so $|a_n|^{1/n} < 1/R_1 + \epsilon$ and similarly for b_n . Multiplying we obtain

$$|a_n b_n|^{1/n} < \frac{1}{R_1 R_2} + \epsilon(1/R_1 + 1/R_2) + \epsilon^2$$

Then

$$\frac{1}{R} \leq \frac{1}{R_1 R_2} \Rightarrow R_1 R_2 \leq R$$

7. If $\lim_{n \rightarrow \infty} |a_n|/|a_{n+1}| = R$, prove that $\sum a_n z^n$ has a radius of convergence of R .

Let $\epsilon > 0$ be given. Suppose $|z| < R$. Pick ϵ such that $|z| < R - \epsilon$. Then for some $n > N$

$$\begin{aligned} \left| \frac{a_n}{a_{n+1}} - R \right| &\leq R - \left| \frac{a_n}{a_{n+1}} \right| < \epsilon \\ R - \epsilon &< \left| \frac{a_n}{a_{n+1}} \right| \end{aligned} \quad (2.8)$$

For $n > N$, we can write

$$\left| \frac{a_N}{a_n} \right| = \left| \frac{a_N a_{N+1} \cdots a_{n-1}}{a_{N+1} a_{N+2} \cdots a_n} \right| = \left| \frac{a_N}{a_{N+1}} \frac{a_{N+1}}{a_{N+2}} \cdots \frac{a_{n-1}}{a_n} \right| \quad (2.9)$$

For $n > N$, we have that from equation (2.8), $R - \epsilon < \frac{a_N}{a_{N+1}}$. Thus, we can equation (2.9) as

$$(R - \epsilon)^{n-N} < \left| \frac{a_N}{a_n} \right|$$

$$|a_n| < \frac{|a_N|}{(R - \epsilon)^{n-N}}$$

$$|a_n z^n| < |a_N z^N| \left(\frac{|z|}{R - \epsilon} \right)^{n-N}$$

Since ϵ was chosen such that $|z| < R - \epsilon$, we that $\frac{|z|}{R - \epsilon} < 1$ and

$$|a_n z^n| < |a_N z^N|$$

where $|a_N z^N| < \infty$ since it is a convergent geometric series. Therefore, $\sum a_n z^n$ converges absolutely with a radius of convergence of R .

8. For what values of z is

$$\sum_{n=0}^{\infty} \left(\frac{z}{1+z} \right)^n$$

convergent?

In order for series to converge $\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|} < 1$. Then

$$\limsup_{n \rightarrow \infty} \sqrt[n]{|z/(z+1)|^n} = \left| \frac{z}{z+1} \right| < 1$$

or $|z|^2 < (1+z)(1+\bar{z}) = 1 + 2\Re\{z\} + |z|^2$ so the series converges when

$$0 < 1 + 2\Re\{z\}.$$

9. Same question for

$$\sum_{n=0}^{\infty} \frac{z^n}{1+z^{2n}}.$$

Consider the following two equations:

$$\begin{aligned} |1| &= |1 + z^{2n} - z^{2n}| \\ &\leq |1 + z^{2n}| + |z^{2n}| \\ 1 - |z^{2n}| &\leq |1 + z^{2n}| \end{aligned} \tag{2.10}$$

$$\begin{aligned} |z^{2n}| &= |1 - 1 + z^{2n}| \\ &\leq |1 + z^{2n}| + 1 \\ |z^{2n}| - 1 &\leq |1 + z^{2n}| \end{aligned} \tag{2.11}$$

From equations (2.10) and (2.11), the triangle inequality, we have that

$$|1 - |z^{2n}|| \leq |1 + z^{2n}|.$$

There exists an $m > 1$ such that

$$\frac{|z^{2n}|}{m} \leq |1 - |z^{2n}||.$$

By the root test,

$$\limsup_{n \rightarrow \infty} \sqrt[n]{\frac{m|z|^n}{|z^2|^n}} = \limsup_{n \rightarrow \infty} \frac{\sqrt[n]{m}}{|z|} = \frac{1}{|z|} < 1$$

When $|z| > 1$, the convergence of the ratio test $\frac{1}{|z|} < 1$ leads to $|z| > 1$. If $|z| < 1$, then $1/|z| > 1$ where we can write $1/|z| = |z_1|$. Since the choice dummy variables is arbitrary, $|z| < 1$. In other words, the series will converge when $|z| > 1$ or $|z| < 1$. Suppose $|z| = 1$. Then by the limit test,

$$\lim_{n \rightarrow \infty} \frac{1}{1^n + 1^{-n}} = \frac{1}{2} \neq 0;$$

therefore, the series diverges.

2.2.5 Abel's Limit Theorem

2.3 The Exponential and Trigonometric Functions

2.3.1 The Exponential

2.3.2 The Trigonometric Functions

1. Find the values of $\sin(i)$, $\cos(i)$, and $\tan(1+i)$.

For $\sin(i)$, we can use the identity $\sin(z) = \frac{e^{iz} - e^{-iz}}{2i}$. Then

$$\sin(i) = \frac{e^{-1} - e^1}{2i} = i \frac{e^1 - e^{-1}}{2} = i \sinh(1).$$

Similarly, for $\cos(i)$, we have

$$\cos(i) = \frac{e^{-1} + e^1}{2} = \frac{e^1 + e^{-1}}{2} = \cosh(1).$$

For $\tan(1+i)$, we can use the identity $\tan(z) = -i \frac{e^{iz} - e^{-iz}}{e^{iz} + e^{-iz}}$. Then

$$\tan(1+i) = -i \frac{e^{i-1} - e^{1-i}}{e^{i-1} + e^{1-i}} = -i \tanh(i-1).$$

2. The hyperbolic cosine and sine are defined as $\cosh(z) = \frac{e^z + e^{-z}}{2}$ and $\sinh(z) = \frac{e^z - e^{-z}}{2}$. Express them through $\cos(iz)$ and $\sin(iz)$. Derive the addition formulas, and formulas for $\cosh(2z)$ and $\sinh(2z)$.

For the first part, we have

$$\begin{aligned}\cos(iz) &= \frac{e^{-z} + e^z}{2} \\ &= \cosh(z) \\ \sin(iz) &= \frac{e^{-z} - e^z}{2i} \\ &= i \frac{e^z - e^{-z}}{2} \\ &= i \sinh(z)\end{aligned}$$

For \cosh , we have that the addition formula is

$$\begin{aligned}\cosh(a+b) &= \cos[i(a+b)] \\ &= \frac{e^{a+b} + e^{-(a+b)}}{2} \\ &= \frac{2e^{a+b} + 2e^{-(a+b)}}{4} \\ &= (e^{a+b} + e^{a-b} + e^{b-a} + e^{-(a+b)} + e^{a+b} - e^{a-b} - e^{b-a} + e^{-(a+b)})/4 \\ &= \frac{e^a + e^{-a}}{2} \frac{e^b + e^{-b}}{2} + \frac{e^a - e^{-a}}{2} \frac{e^b - e^{-b}}{2} \\ &= \cosh(a) \cosh(b) + \sinh(a) \sinh(b)\end{aligned}$$

For \sinh , we have that the addition formula is

$$\begin{aligned}\sinh(a+b) &= -i \sin[i(a+b)] \\ &= \frac{e^{-(a+b)} - e^{a+b}}{-2} \\ &= \frac{2e^{a+b} - 2e^{-(a+b)}}{4} \\ &= (e^{a+b} + e^{a-b} - e^{b-a} - e^{-(a+b)} + e^{a+b} - e^{a-b} + e^{b-a} - e^{-(a+b)})/4 \\ &= \frac{e^a - e^{-a}}{2} \frac{e^b + e^{-b}}{2} + \frac{e^a + e^{-a}}{2} \frac{e^b - e^{-b}}{2}\end{aligned}$$

$$= \sinh(a) \cosh(b) + \cosh(a) \sinh(b)$$

For the double angle formulas, recall that $\cos(2z) = \cos^2(z) - \sin^2(z) = 2\cos^2(z) - 1 = 1 - 2\sin^2(z)$ and $\sin(2z) = 2\sin(z)\cos(z)$. Therefore, we have

$$\begin{aligned}\cosh(2z) &= \cos(2iz) \\ &= \cos^2(iz) - \sin^2(iz) \\ &= \left(\frac{e^z + e^{-z}}{2}\right)^2 - \left(\frac{e^z - e^{-z}}{2}\right)^2 \\ &= \cosh^2(z) + \sinh^2(z) \\ \cosh(2z) &= 2\cos^2(iz) - 1 \\ &= 2\cosh^2(z) - 1 \\ \cosh(2z) &= 1 - 2\sin^2(iz) \\ &= 1 - 2\sinh^2(z) \\ \sinh(2z) &= -i\sin(2iz) \\ &= -2i\sin(iz)\cos(iz) \\ &= 2\frac{e^z - e^{-z}}{2}\frac{e^z + e^{-z}}{2} \\ &= 2\sinh(z)\cosh(z)\end{aligned}$$

3. Use the addition formulas to separate $\cos(x + iy)$ and $\sin(x + iy)$ in real and imaginary parts.

$$\begin{aligned}\cos(x + iy) &= \cos(x)\cos(iy) - \sin(x)\sin(iy) \\ &= \cos(x)\cosh(y) - i\sin(x)\sinh(y) \\ \sin(x + iy) &= \sin(x)\cos(iy) + \sin(iy)\cos(x) \\ &= \sin(x)\cosh(y) + i\sinh(y)\cos(x)\end{aligned}$$

4. Show that

$$|\cos(z)|^2 = \sinh^2(y) + \cos^2(x) = \cosh^2(y) - \sin^2(x) = (\cosh(2y) + \cos(2x))/2$$

and

$$|\sin(z)|^2 = \sinh^2(y) + \sin^2(x) = \cosh^2(y) - \cos^2(x) = (\cosh(2y) - \cos(2x))/2.$$

For the identities, recall that $\cosh^2(z) - \sinh^2(z) = 1$ and $\cos^2(z) + \sin^2(z) = 1$. Then for the first identity, we have

$$\begin{aligned}|\cos(z)|^2 &= \cos(z)\cos(\bar{z}) \\ &= [\cos(x)\cosh(y) - i\sin(x)\sinh(y)][\cos(x)\cosh(y) + i\sin(x)\sinh(y)] \\ &= \cos^2(x)\cosh^2(y) + \sin^2(x)\sinh^2(y) \\ &= \cos^2(x)(1 + \sinh^2(y)) + \sin^2(x)\sinh^2(y) \\ &= \cos^2(x) + \sinh^2(y) \\ |\cos(z)|^2 &= \cos^2(x)\cosh^2(y) + \sin^2(x)(\cosh^2(y) - 1) \\ &= \cosh^2(y) - \sin^2(x) \\ |\cos(z)|^2 &= \\ &= (\cosh(2y) + \cos(2x))/2 \\ |\sin(z)|^2 &= \sin(z)\sin(\bar{z}) \\ &= [\sin(x)\cosh(y) + i\sinh(y)\cos(x)][\sin(x)\cosh(y) - i\sinh(y)\cos(x)] \\ &= \sin^2(x)\cosh^2(y) + \sinh^2(y)\cos^2(x) \\ &= \sin^2(x)(1 + \sinh^2(y)) + \sinh^2(y)\cos^2(x) \\ &= \sin^2(x) + \sinh^2(y)\end{aligned}$$

$$\begin{aligned}
|\sin(z)|^2 &= \sin^2(x) \cosh^2(y) + (\cosh^2(y) - 1) \cos^2(x) \\
&= \cosh^2(y) - \cos^2(x) \\
|\sin(z)|^2 &= \\
&= (\cosh(2y) - \cos(2x))/2
\end{aligned}$$

2.3.3 Periodicity

2.3.4 The Logarithm

1. For real y , show that every remainder in the series for $\cos(y)$ and $\sin(y)$ has the same sign as the leading term (this generalizes the inequalities used in the periodicity proof).

The series for both cosine and sine are

$$\begin{aligned}
\cos(y) &= \sum_{k=0}^{\infty} (-1)^k \frac{y^{2k}}{(2k)!} \\
&= 1 - \frac{y^2}{2!} + \frac{y^4}{4!} - \dots \\
\sin(y) &= \sum_{k=0}^{\infty} (-1)^k \frac{y^{2k+1}}{(2k+1)!} \\
&= y - \frac{y^3}{3!} + \frac{y^5}{5!} - \dots
\end{aligned}$$

We can write Taylor's formula as $f(y) = T_n(y) + R_n(y)$ where

$$f(y) = \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} y^k + \frac{1}{k!} \int_0^y (y-t)^k f^{(k+1)}(t) dt.$$

Now, we can write cosine and sine of y as

$$\begin{aligned}
\cos(y) &= \sum_{k=0}^n \frac{(-1)^k y^{2k}}{(2k)!} + \frac{1}{n!} \int_0^y (y-t)^n \cos^{n+1}(t) dt \\
\sin(y) &= \sum_{k=0}^{n-1} \frac{(-1)^k y^{2k+1}}{(2k+1)!} + \frac{1}{n!} \int_0^y (y-t)^n \sin^n(t) dt
\end{aligned}$$

For cosine and sine, let $n = 2m$ and $n = 2m - 1$, respectively. Then

$$\begin{aligned}
\cos(y) &= \sum_{k=0}^m \frac{(-1)^k y^{2k}}{(2k)!} + \frac{1}{(2m)!} \int_0^y (y-t)^{2m} \cos^{2m+1}(t) dt \\
\sin(y) &= \sum_{k=0}^{m-1} \frac{(-1)^k y^{2k+1}}{(2k+1)!} + \frac{1}{(2m-1)!} \int_0^y (y-t)^{2m-1} \sin^{2m-1}(t) dt
\end{aligned}$$

2. Prove, for instance, that $3 < \pi < 2\sqrt{3}$.
3. Find the value of e^z for $z = -i\pi/2, 3i\pi/4, 2i\pi/3$.

$$\begin{aligned}
e^{-i\pi/2} &= -i \\
e^{3i\pi/4} &= (-\sqrt{2} + i\sqrt{2})/2 \\
e^{2i\pi/3} &= (-1 + i\sqrt{3})/2
\end{aligned}$$

4. For that values of z is e^z equal to $2, -1, i, -i/2, -1 - i, 1 + 2i$?

For all problems, $k \in \mathbb{Z}$.

$$\begin{aligned}
e^z &= 2 & e^z &= -1 \\
z &= \log(2) + 2ki\pi & z &= \log(-1)
\end{aligned}$$

$$\begin{aligned}
e^z &= i & &= \log|i| + i(\arg(-1) + 2k\pi) \\
z &= \log|i| + i(\arg(i) + 2k\pi) & &= i\pi(1 + 2k\pi) \\
&= \frac{i\pi}{2}(1 + 4k) & &e^z = \frac{-i}{2} \\
e^z &= -1 - i & &= -\log(2) - \frac{i\pi}{2}(1 + 4k) \\
z &= \log|-1 - i| + i(\arg(-1 - i) + 2k\pi) & &e^z = 1 + 2i \\
&= \log(\sqrt{2}) - \frac{3i\pi}{4} + 2ki\pi & &z = \log(\sqrt{5}) + i(\arctan(2) + 2k\pi) \\
&= \frac{\log(2)}{2} - \frac{3i\pi}{4} + 2ki\pi & &= \frac{\log(5)}{2} + i(\arctan(2) + 2k\pi)
\end{aligned}$$

5. Find the real and imaginary parts of $\exp(e^z)$.

Let $z = x + iy$. Then

$$\begin{aligned}
\exp(e^z) &= \exp[e^x(\cos(y) + i\sin(y))] \\
&= \exp(e^x \cos(y)) \exp(ie^x \sin(y)) \\
&= \exp(e^x \cos(y)) [\cos(e^x \sin(y)) + i\sin(e^x \sin(y))] \\
u(x, y) &= \exp(e^x \cos(y)) \cos(e^x \sin(y)) \\
v(x, y) &= \exp(e^x \cos(y)) \sin(e^x \sin(y))
\end{aligned}$$

where $u(x, y)$ is the real and $v(x, y)$ is the imaginary part of $\exp(e^z)$.

6. Determine all values of $2^i, i^i, (-1)^{2i}$.

For all problems, $k \in \mathbb{Z}$.

$$\begin{aligned}
z &= 2^i \\
&= \exp[i \log(2)] \\
&= \cos(\log(2)) + i\sin(\log(2)) \\
z &= i^i \\
&= \exp[i \log(i)] \\
&= \exp[-\pi(1 + 4k)/2] \\
z &= (-1)^{2i} \\
&= i^{4i} \\
&= (i^i)^4 \\
&= \exp[-2\pi(1 + 4k)]
\end{aligned}$$

7. Determine the real and imaginary parts of z^z .

Let $z = x + iy$ and $k \in \mathbb{Z}$.

$$\begin{aligned}
z^z &= (x + iy)^{x+iy} \\
&= \exp[(x + iy) \log(x + iy)] \\
&= e^{x/2 \log(x^2+y^2) - y(\arctan(y/x) + 2k\pi)} \left[\cos(x(\arctan(y/x) + 2k\pi) + y/2 \log(x^2 + y^2)) + \right. \\
&\quad \left. i \sin(x(\arctan(y/x) + 2k\pi) + y/2 \log(x^2 + y^2)) \right]
\end{aligned}$$

Thus, the real part is

$$u(x, y) = e^{x/2 \log(x^2+y^2) - y(\arctan(y/x) + 2k\pi)} \cos(x(\arctan(y/x) + 2k\pi) + y/2 \log(x^2 + y^2))$$

and the imaginary part is

$$v(x, y) = e^{x/2 \log(x^2+y^2) - y(\arctan(y/x) + 2k\pi)} \sin(x(\arctan(y/x) + 2k\pi) + y/2 \log(x^2 + y^2))$$

8. Express $\arctan(w)$ in terms of the logarithm.

Let $\arctan(w) = z$. Then $w = \tan(z)$. Recall that $\tan(z) = -i \frac{e^{iz} - e^{-iz}}{e^{iz} + e^{-iz}}$. Now, let $e^{2iz} = x$. Then we have the following

$$w = -i \frac{x^2 - 1}{x^2 + 1}$$

which leads to

$$e^{2iz} = \frac{i - w}{i + w}.$$

By taking the log, we can recover z .

$$\begin{aligned} 2iz &= \log(i - w) - \log(i + w) \\ &= \log(i) + \log(1 - iw) - \log(i) - \log(1 + iw) \\ z &= \frac{i}{2} [\log(1 - iw) - \log(1 + iw)] \\ \arctan(w) &= z \\ &= \frac{i}{2} [\log(1 - iw) - \log(1 + iw)] \end{aligned}$$

9. Show how to define the "angles" in a triangle, bearing in mind that they should lie between 0 and π . With this definition, prove that the sum of the angles is π .

10. Show that the roots of the binomial equation $z^n = a$ are the vertices of a regular polygon (equal sides and angles).

3 Analytic Functions as Mappings

3.1 Elementary Point Set Topology

3.1.1 Sets and Elements

3.1.2 Metric Spaces

1. If S is a metric space with distance function $d(x, y)$, show that S with the distance function $\delta(x, y) = d(x, y)/[1 + d(x, y)]$ is also a metric space. The latter space is bounded in the sense that all distances lie under a fixed bound.

Since $d(x, y)$ is a metric on S , $d(x, y)$ satisfies

$$d(x, y) \geq 0, \text{ and zero only when } x = y \quad (3.1a)$$

$$d(x, y) = d(y, x) \quad (3.1b)$$

$$d(x, z) \leq d(x, y) + d(y, z) \quad (3.1c)$$

By equation (3.1a), for $x = y$, $d(x, y) = 0$ so $\delta(x, y) = 0/1 = 0$, and when $x \neq y$, $d(x, y) > 0$ so $\delta(x, y) > 0$ since a positive number divided by a positive number is positive. We have that $\delta(x, y) \geq 0$ and equal zero if and only if $x = y$. By equation (3.1b), we have

$$\begin{aligned} \delta(x, y) &= \frac{d(x, y)}{1 + d(x, y)} \\ &= \frac{d(y, x)}{1 + d(y, x)} \\ &= \delta(y, x) \end{aligned}$$

For the triangle inequality, we have

$$\frac{d(x, z)}{1 + d(x, z)} \leq \frac{d(x, y)}{1 + d(x, y)} + \frac{d(y, z)}{1 + d(y, z)}$$

Let's multiple through by the product of all three denominators. After simplifying, we obtain

$$d(x, z) \leq d(x, y) + d(y, z) + 2d(x, y)d(y, z) + d(x, y)d(y, z)d(x, z)$$

We have already shown that $d(x, y) \geq 0$ and zero if and only if $x = y$. If $x = y = z$, the triangle inequality is vacuously true. When $x \neq y \neq z$, the triangle inequality follows since each distance is positive and equation (3.1c); that is,

$$\delta(x, z) \leq \delta(x, y) + \delta(y, z).$$

2. Suppose that there are given two distance functions $d(x, y)$ and $d_1(x, y)$ on the same space S . They are said to be equivalent if they determine the same open sets. Show that d and d_1 are equivalent if to every $\epsilon > 0$ there exists a $\delta > 0$ such that $d(x, y) < \delta$ implies $d_1(x, y) < \epsilon$, and vice versa. Verify that this condition is fulfilled in item 1.
3. Show by strict application of the definition that the closure of $|z - z_0| < \delta$ is $|z - z_0| \leq \delta$.
4. If X is the set of complex numbers whose real and imaginary parts are rational, what is $\text{Int } X, \bar{X}, \partial X$?
5. It is sometimes typographically simpler to write X' for $\sim X$. With this notation, how is $X'^{-'}$ related to X ? Show that $X^{-' -' -' -'} = X^{-' -'}$.
6. A set is said to be discrete if all its points are isolated. Show that a discrete set in \mathbb{R} or \mathbb{C} is countable.