

# Spherical solutions for stars

Daniel Wysocki

Rochester Institute of Technology

General Relativity I Presentations

December 14th, 2015



# Introduction



# Spherically symmetric coordinates



## Two-sphere in flat spacetime

## General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)$$

## Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2\theta d\phi^2) \equiv r^2 d\Omega^2$$



2015-12-12

## Spherical stars

## └ Spherically symmetric coordinates

## └ Two-sphere in flat spacetime

## General metric:

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)$$

$$dl^2 = r^2(d\theta^2 + \sin^2\theta d\phi^2) \equiv r^2 d\Omega^2$$

## Two-sphere in flat spacetime

## General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

## Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

2015-12-12

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in flat spacetime

Two-sphere in flat spacetime

General metric:

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

Schutz (2009, p. 256)

## Two-sphere in curved spacetime

## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

2015-12-12

## Spherical stars

## └ Spherically symmetric coordinates

## └ Two-sphere in curved spacetime

## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

## Two-sphere in curved spacetime

## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

2015-12-12

Spherical stars

- └ Spherically symmetric coordinates
  - └ Two-sphere in curved spacetime

Two-sphere in curved spacetime

Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$


Relation to  $r$ 

$$f(r', t) \equiv r^2$$

---

 Schutz (2009, pp. 256–257)

# Meaning of $r$



Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

2015-12-12

## Spherical stars

- └ Spherically symmetric coordinates
- └ Meaning of  $r$

Meaning of  $r$ 

Mark Hannam


- “curvature” or “area” coordinate
- radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)



# Meaning of $r$



Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

2015-12-12

## Spherical stars

- └ Spherically symmetric coordinates
- └ Meaning of  $r$

Meaning of  $r$ 


Mark Hannam

- “curvature” or “area” coordinate
- radius of curvature and area
- *not* proper distance from center

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)

# Meaning of $r$



Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:

Surface with circular symmetry but no coordinate  $r = 0$ .

2015-12-12

## Spherical stars

- └ Spherically symmetric coordinates
- └ Meaning of  $r$



Mark Hannam

- “curvature” or “area” coordinate
- radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)

# Spherically symmetric spacetime

## General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

$g_{00}$ ,  $g_{0r}$ , and  $g_{rr}$  functions of  $t$  and  $r$



2015-12-12

Spherical stars

└ Spherically symmetric coordinates

└ Spherically symmetric spacetime

Spherically symmetric spacetime

General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

$g_{00}$ ,  $g_{0r}$ , and  $g_{rr}$  functions of  $t$  and  $r$

---

 Schutz (2009, p. 258)

# Static spacetimes



# Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Motivation

- George David Birkhoff

Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

# Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Motivation

- George David Birkhoff

Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations

• Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

# Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Motivation

- George David Birkhoff

Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

## Definition

A spacetime is static if we can find a time coordinate  $t$  for which

- (i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Definition

- a spacetime which only satisfies the first condition is *stationary*
  - e.g. rotating star

## Definition

A spacetime is static if we can find a time coordinate  $t$  for which

- (i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



# Definition

A spacetime is static if we can find a time coordinate  $t$  for which

(i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

(ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Definition

- a spacetime which only satisfies the first condition is *stationary*
  - e.g. rotating star

## Definition

A spacetime is static if we can find a time coordinate  $t$  for which

(i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

(ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

## Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

Spherical stars  
└ Static spacetimes

## └ Time reversal

- $g_{\bar{0}\bar{r}} = -g_{0r} = 0$

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{j}} = x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j$$

$$\Lambda^0_{\bar{i}} = x^0_{,\bar{i}} = x^0_{,i} = 0$$

## Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

# Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Time reversal

- $g_{\bar{0}\bar{r}} = -g_{0r} = 0$

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

### Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

### Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

Schutz (2009, p. 258)

## Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Time reversal

- $g_{\bar{0}\bar{r}} = -g_{0r} = 0$

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

Schutz (2009, p. 258)

# The metric

## Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

## Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

## Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ The metric

- constraint  $g_{00} < 0 < g_{rr}$  holds for stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Schutz (2009, pp. 258–259)

# The metric

## Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

## Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

## Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ The metric

- constraint  $g_{00} < 0 < g_{rr}$  holds for stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Schutz (2009, pp. 258–259)

# The metric

## Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

## Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

## Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ The metric

- constraint  $g_{00} < 0 < g_{rr}$  holds for stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Schutz (2009, pp. 258–259)

# Einstein Tensor

## General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

## Einstein tensor components

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Einstein Tensor

- $x' \equiv dx/dr$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Schutz (2009, pp. 165, 260)



# Einstein Tensor

## General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

## Einstein tensor components

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Einstein Tensor

- $x' \equiv dx/dr$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Schutz (2009, pp. 165, 260)

# Static perfect fluid



# Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

## Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

## Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-12

Spherical stars  
 └ Static perfect fluid  
   └ Four-velocity

- “conservation law” is the conservation of four-momentum

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

### Constraints

 $U^i = 0$  (static) $\vec{U} \cdot \vec{U} = -1$  (conservation law)

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Schutz (2009, p. 260)

# Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

## Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

## Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-12

Spherical stars  
 └ Static perfect fluid  
 └ Four-velocity

Four-velocity

Constraints	
$U^i = 0$ (static)	$\vec{U} \cdot \vec{U} = -1$ (conservation law)
Solving for $U^0$	
$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$	
$U_0 = g_{00}U^0 = -e^{\Phi}$	
Schutz (2009, p. 260)	

- “conservation law” is the conservation of four-momentum

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

# Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

## Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

## Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-12

Spherical stars  
 └ Static perfect fluid  
 └ Four-velocity

- “conservation law” is the conservation of four-momentum

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

### Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

### Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

### Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Schutz (2009, p. 260)

# Stress-energy tensor

## $T_{\alpha\beta}$ for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

## Components of $T_{\alpha\beta}$

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{\alpha\beta} = 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii}$$


$$T_{rr} = pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Stress-energy tensor

- $T_{\alpha\beta} = 0$  because
  - the cross terms make  $g_{\alpha\beta} = 0$
  - $T_{0i}$  makes one of the  $U$ 's  $U_i = 0$
- likewise,  $T_{ii} = pg_{ii}$  because  $U_i U_i = 0$



# Stress-energy tensor

## $T_{\alpha\beta}$ for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

## Components of $T_{\alpha\beta}$

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{\alpha\beta} = 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii}$$

$$T_{rr} = pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

## Spherical stars

- Static perfect fluid

## Stress-energy tensor

- $T_{\alpha\beta} = 0$  because
  - the cross terms make  $g_{\alpha\beta} = 0$
  - $T_{0i}$  makes one of the  $U$ 's  $U_i = 0$
- likewise,  $T_{ii} = pg_{ii}$  because  $U_i U_i = 0$

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{\alpha\beta} = 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii}$$

$$T_{rr} = pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

## Equation of state

## Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies



2015-12-12

Spherical stars  
└ Static perfect fluid

└ Equation of state

## Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies



# Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO:** prove

## Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

2015-12-12

 Spherical stars  
 └ Static perfect fluid

└ Equations of motion

Equations of motion

Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

 • symmetries make only non-trivial solution  $\alpha = r$   
 TODO: prove

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

Schutz (2009, pp. 175, 261)

# Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO: prove**

## Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

2015-12-12

## Spherical stars

- └ Static perfect fluid

- └ Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO: prove**

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

# Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$

**TODO: prove**

## Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Equations of motion

Equations of motion

Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$

**TODO: prove**

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

Schutz (2009, pp. 175, 261)

## Mass function

## Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

 $m(r)$ 

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

## Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-12

## Spherical stars


└ Static perfect fluid

└ Mass function

- inspect  $(0,0)$  component of Einstein equations
- in Newtonian limit,  $m(r)$  is mass within radius  $r$

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable



## Mass function

## Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

 $m(r)$ 

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

## Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Mass function

- inspect  $(0,0)$  component of Einstein equations
- in Newtonian limit,  $m(r)$  is mass within radius  $r$

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$m(r)$

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

$\frac{dm(r)}{dr} = 4\pi r^2 \rho$

Schutz (2009, pp. 260–262)

# Mass function

## Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

## $m(r)$

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

## Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Mass function

- inspect  $(0,0)$  component of Einstein equations
- in Newtonian limit,  $m(r)$  is mass within radius  $r$

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable

Mass function
Einstein field equations
$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$
$m(r)$
$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$
Relation to energy density
$\frac{dm(r)}{dr} = 4\pi r^2 \rho$
Schutz (2009, pp. 260–262)

What is  $\Phi$  called?!!!!!!?

## Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$\Phi(r)$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ What is  $\Phi$  called?!!!!!!?

- inspect  $(r, r)$  component of Einstein equations

What is  $\Phi$  called?!!!!!!?

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

Schutz (2009, pp. 260–262)

What is  $\Phi$  called?!

## Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

## $\Phi(r)$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ What is  $\Phi$  called?!

- inspect  $(r, r)$  component of Einstein equations

What is  $\Phi$  called?!

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

 $\Phi(r)$ 

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

Schutz (2009, pp. 260–262)



# Exterior Geometry



## Schwarzschild metric I

## Condition

$$\rho = p = 0$$

## Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$m(r) = M$$


$$\Phi(r) = \frac{1}{2} \log \left( 1 - \frac{2M}{r} \right)$$

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
- breaks down when matter surrounds star
- $M$  is a constant



## Schwarzschild metric I

## Condition

$$\rho = p = 0$$

## Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$m(r) \equiv M$$

$$\Phi(r) = \frac{1}{2} \log \left( 1 - \frac{2M}{r} \right)$$

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
- breaks down when matter surrounds star
- $M$  is a constant

Schwarzschild metric I

Condition	
	$\rho = p = 0$
Consequences	
$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$	$m(r) \equiv M$
$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$	$\Phi(r) = \frac{1}{2} \log \left( 1 - \frac{2M}{r} \right)$

Schutz (2009, pp. 262–263)

## Schwarzschild metric I

## Condition

$$\rho = p = 0$$

## Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$m(r) \equiv M$$

$$\Phi(r) = \frac{1}{2} \log \left( 1 - \frac{2M}{r} \right)$$

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
- breaks down when matter surrounds star
- $M$  is a constant

Condition	
	$\rho = p = 0$
Consequences	
$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$	$m(r) \equiv M$
$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$	$\Phi(r) = \frac{1}{2} \log \left( 1 - \frac{2M}{r} \right)$

## Schwarzschild metric II

## First two metric components

$$g_{tt} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right) \quad g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1}$$

## Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

$$g_{tt} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right) \quad g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1}$$

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

## Schwarzschild metric II

## First two metric components

$$g_{tt} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right) \quad g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1}$$

## Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$



2015-12-12

Spherical stars  
└ Exterior Geometry

└ Schwarzschild metric II

Schwarzschild metric II

First two metric components

$$g_{tt} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right) \quad g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1}$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Schutz (2009, pp. 258, 262–263)

## Far-field metric

## Condition

$$r \gg M$$

## Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

## Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-12


## Spherical stars

## └ Exterior Geometry

## └ Far-field metric

- far-field metric of a star

- mass  $M$
- distance  $R$



## Far-field metric

## Condition

$$r \gg M$$

## Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

## Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Far-field metric

- far-field metric of a star
  - mass  $M$
  - distance  $R$

Far-field metric

Condition	$r \gg M$
Far-field Schwarzschild metric	$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$
Far-field Schwarzschild metric (Cartesian)	$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$
	$R^2 \equiv x^2 + y^2 + z^2$
Schutz (2009, pp. 263)	



## Far-field metric

## Condition

$$r \gg M$$

## Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

## Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-12

## Spherical stars

## └ Exterior Geometry

## └ Far-field metric

- far-field metric of a star
  - mass  $M$
  - distance  $R$

Condition	$r \gg M$
Far-field Schwarzschild metric	$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$
Far-field Schwarzschild metric (Cartesian)	$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$
	$R^2 \equiv x^2 + y^2 + z^2$
Schutz (2009, pp. 263)	

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

- George David Birkhoff

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

- George David Birkhoff

Let the geometry of a given region of spacetime:

- be spherically symmetric

• be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

- George David Birkhoff

Let the geometry of a given region of spacetime:

- be spherically symmetric
- be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

---

 Birkhoff and Langer (1923)

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))



2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

- George David Birkhoff

Let the geometry of a given region of spacetime:

- be spherically symmetric
- be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

---

 Birkhoff and Langer (1923)

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))



2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

- George David Birkhoff

Let the geometry of a given region of spacetime:

- be spherically symmetric
- be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

---

 Birkhoff and Langer (1923)

# Interior structure



# Tolman–Oppenheimer–Volkov (T–O–V) equation

## Condition

$$\rho \neq 0; \quad p \neq 0$$

## Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

$$\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

## T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-12

## Spherical stars

### └ Interior structure

### └ Tolman–Oppenheimer–Volkov (T–O–V) equation

- T–O–V equation coupled with  $dm/dr$  and  $p(\rho)$ 
  - 3 equations
  - 3 unknowns ( $m$ ,  $\rho$ ,  $p$ )
  - $\Phi(r)$  only intermediate variable

Condition
$\rho \neq 0; \quad p \neq 0$
$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$
$\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
Schutz (2009, pp. 261–264)



# Tolman–Oppenheimer–Volkov (T–O–V) equation

## Condition

$$\rho \neq 0; \quad p \neq 0$$

## Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

## T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-12

## Spherical stars

### └ Interior structure

### └ Tolman–Oppenheimer–Volkov (T–O–V) equation

- T–O–V equation coupled with  $dm/dr$  and  $p(\rho)$ 
  - 3 equations
  - 3 unknowns ( $m$ ,  $\rho$ ,  $p$ )
  - $\Phi(r)$  only intermediate variable

Condition
$\rho \neq 0; \quad p \neq 0$
Recall
$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$ $\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
T–O–V equation
$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
Schutz (2009, pp. 261–264)

## Tolman–Oppenheimer–Volkov (T–O–V) equation

## Condition

$$\rho \neq 0; \quad p \neq 0$$

## Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

## T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-12

## Spherical stars

## └ Interior structure

## └ Tolman–Oppenheimer–Volkov (T–O–V) equation

- T–O–V equation coupled with  $dm/dr$  and  $p(\rho)$ 
  - 3 equations
  - 3 unknowns ( $m$ ,  $\rho$ ,  $p$ )
  - $\Phi(r)$  only intermediate variable

Tolman–Oppenheimer–Volkov (T–O–V) equation	
Condition	$\rho \neq 0; \quad p \neq 0$
Recall	$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$ $\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
T–O–V equation	$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
Schutz (2009, pp. 261–264)	

# Tolman–Oppenheimer–Volkov (T–O–V) equation

## Condition

$$\rho \neq 0; \quad p \neq 0$$

## Recall

$$(\rho + p) \frac{d\Phi}{dr} = - \frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

## T–O–V equation

$$\frac{dp}{dr} = - \frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-12

## Spherical stars

### └ Interior structure

### └ Tolman–Oppenheimer–Volkov (T–O–V) equation

- T–O–V equation coupled with  $dm/dr$  and  $p(\rho)$ 
  - 3 equations
  - 3 unknowns ( $m$ ,  $\rho$ ,  $p$ )
  - $\Phi(r)$  only intermediate variable

<b>Condition</b>	$\rho \neq 0; \quad p \neq 0$
<b>Recall</b>	$(\rho + p) \frac{d\Phi}{dr} = - \frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
<b>T–O–V equation</b>	$\frac{dp}{dr} = - \frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
<small>Schutz (2009, pp. 261–264)</small>	

# Newtonian hydrostatic equilibrium

## Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

## Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

2015-12-12

## Spherical stars

### └ Interior structure

### └ Newtonian hydrostatic equilibrium

- Newtonian limit reduces the T–O–V equation to HSE

#### Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

# Newtonian hydrostatic equilibrium

## Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

## Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

2015-12-12

## Spherical stars

### └ Interior structure

### └ Newtonian hydrostatic equilibrium

- Newtonian limit reduces the T–O–V equation to HSE

#### Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

#### Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

# References





G. D. Birkhoff and R. E. Langer. *Relativity and modern physics*. 1923.



S. M. Carroll. *Spacetime and geometry. An introduction to general relativity*. 2004.



C. J. Hansen and S. D. Kawaler. *Stellar Interiors. Physical Principles, Structure, and Evolution*. 1994.



C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. 1973.



B. Schutz. *A First Course in General Relativity*. May 2009.



N. Stergioulas. Rotating Stars in Relativity. *Living reviews in relativity*, 6:3, June 2003. [Online; accessed 2015-12-09]. DOI: 10.12942/lrr-2003-3. eprint: gr-qc/0302034.



-  G. D. Birkhoff and R. E. Langer. *Relativity and modern physics*. 1923.
-  S. M. Carroll. *Spacetime and geometry. An introduction to general relativity*. 2004.
-  C. J. Hansen and S. D. Kawaler. *Stellar Interiors. Physical Principles, Structure, and Evolution*. 1994.
-  C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. 1973.
-  B. Schutz. *A First Course in General Relativity*. May 2009.
-  N. Stergioulas. Rotating Stars in Relativity. *Living reviews in relativity*, 6:3, June 2003. [Online; accessed 2015-12-09]. DOI: 10.12942/lrr-2003-3. eprint: gr-qc/0302034.