

Spherical solutions for stars

Daniel Wysocki

Rochester Institute of Technology

General Relativity I Presentations

December 14th, 2015



Introduction

- model stars using spherical symmetry
- Schwarzschild metric
- T–O–V equation
- applications



2015-12-13

Spherical stars

└ Introduction

- I will model stars using GR assuming spherical symmetry
- I will derive the Schwarzschild metric and T–O–V equation
- finally I will relate these equations to modeling specific types of stars

Introduction

- model stars using spherical symmetry
- Schwarzschild metric
- T–O–V equation
- applications

Spherically symmetric coordinates

- First we need to derive our coordinate system



Two-sphere in flat spacetime

General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$



2015-12-13

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in flat spacetime

- we start with the simplest spherically symmetric coordinates
- 2-sphere in Minkowski space

General metric:

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

Two-sphere in flat spacetime

General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

2015-12-13

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in flat spacetime

- we start with the simplest spherically symmetric coordinates
- 2-sphere in Minkowski space

Two-sphere in flat spacetime

General metric:

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

 Schutz (2009, p. 256)

Two-sphere in curved spacetime

Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to r

$$f(r', t) \equiv r^2$$



2015-12-13

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in curved spacetime

- generalize to 2-sphere in arbitrary curved spherically symmetric spacetime
- inclusion of curvature makes r^2 some function of r' and t

$$dl^2 = f(r', t) d\Omega^2$$

$$f(r', t) \equiv r^2$$

Two-sphere in curved spacetime

Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to r

$$f(r', t) \equiv r^2$$



2015-12-13

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in curved spacetime

- generalize to 2-sphere in arbitrary curved spherically symmetric spacetime
- inclusion of curvature makes r^2 some function of r' and t


Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to r

$$f(r', t) \equiv r^2$$

Meaning of r



Mark Hannam

- *not* proper distance from center
- “curvature” or “area” coordinate
 - radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$


Figure:
Surface with circular
symmetry but no
coordinate $r = 0$.

2015-12-13

Spherical stars


└ Spherically symmetric coordinates

└ Meaning of r



- r is not necessary the “distance from the center”
- it is merely a coordinate
- for instance, we may have a spacetime where the center is missing
 - example: wormhole spacetime
- surface of constant (r, t) is a two-sphere of area A and circumference C

Meaning of r



Mark Hannam

- *not* proper distance from center
- “curvature” or “area” coordinate
 - radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$


Figure:
Surface with circular
symmetry but no
coordinate $r = 0$.

2015-12-13

Spherical stars


└ Spherically symmetric coordinates

└ Meaning of r



- r is not necessary the “distance from the center”
- it is merely a coordinate
- for instance, we may have a spacetime where the center is missing
 - example: wormhole spacetime
- surface of constant (r, t) is a two-sphere of area A and circumference C

Meaning of r



Mark Hannam

- *not* proper distance from center
- “curvature” or “area” coordinate
 - radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$

Figure:

Surface with circular symmetry but no coordinate $r = 0$.

2015-12-13

Spherical stars

└ Spherically symmetric coordinates

└ Meaning of r



Figure:
Surface with circular
symmetry but no
coordinate $r = 0$.

Schutz (2009, p. 257)

- not proper distance from center
- “curvature” or “area” coordinate
- radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$

- r is not necessary the “distance from the center”
- it is merely a coordinate
- for instance, we may have a spacetime where the center is missing
 - example: wormhole spacetime
- surface of constant (r, t) is a two-sphere of area A and circumference C

Spherically symmetric spacetime

General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

g_{00} , g_{0r} , and g_{rr} functions of t and r



2015-12-13

Spherical stars

└ Spherically symmetric coordinates

└ Spherically symmetric spacetime

- now consider not only surface of 2-sphere, but whole spacetime
- now we have some unknown g_{00} , g_{rr} , and cross term g_{0r}
- cross terms g_{0i} for $i \in \{\theta, \phi\}$ are zero from symmetry
- need more constraints to say anything particular about them

General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

g_{00} , g_{0r} , and g_{rr} functions of t and r

Static spacetimes

2015-12-13

Spherical stars
└ Static spacetimes

Static spacetimes

- now I will impose the static constraint



Motivation

- leads to simple derivation of Schwarzschild metric
- unique solution to spherically symmetric, asymptotically flat Einstein vacuum field equations (Birkhoff's theorem)

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-13

Spherical stars
 └ Static spacetimes
 └ Motivation

- we choose the constraint of a static spacetime because
 - it allows us to easily derive the Schwarzschild metric
 - according to Birkhoff's theorem, this metric is the unique solution to the Einstein vacuum field equations for spherically symmetric, asymptotically flat spacetimes
- George David Birkhoff

Motivation

- leads to simple derivation of Schwarzschild metric

• unique solution to spherically symmetric, asymptotically flat Einstein vacuum field equations (Birkhoff's theorem)

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

Motivation

- leads to simple derivation of Schwarzschild metric
- unique solution to spherically symmetric, asymptotically flat Einstein vacuum field equations (Birkhoff's theorem)

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-13

Spherical stars
 └ Static spacetimes
 └ Motivation

- we choose the constraint of a static spacetime because
 - it allows us to easily derive the Schwarzschild metric
 - according to Birkhoff's theorem, this metric is the unique solution to the Einstein vacuum field equations for spherically symmetric, asymptotically flat spacetimes
- George David Birkhoff

Motivation

- leads to simple derivation of Schwarzschild metric
- unique solution to spherically symmetric, asymptotically flat Einstein vacuum field equations (Birkhoff's theorem)

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

Definition

A spacetime is static if we can find a time coordinate t for which

- (i) the metric independent of t

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



2015-12-13

Spherical stars
 └ Static spacetimes
 └ Definition

- now I define “static”
- first condition is that the metric is independent of time
 - by itself, this condition is called “stationary”
- second condition is that metric unaffected by time reversal
- e.g. rotating stars are stationary but not static

Definition

A spacetime is static if we can find a time coordinate t for which

- (i) the metric independent of t

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

 Schutz (2009, p. 258)

Definition

A spacetime is static if we can find a time coordinate t for which

- (i) the metric independent of t

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



2015-12-13

Spherical stars
 └ Static spacetimes
 └ Definition

- now I define “static”
- first condition is that the metric is independent of time
 - by itself, this condition is called “stationary”
- second condition is that metric unaffected by time reversal
- e.g. rotating stars are stationary but not static

Definition

A spacetime is static if we can find a time coordinate t for which

- (i) the metric independent of t

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

 Schutz (2009, p. 258)

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$



2015-12-13

Spherical stars
└ Static spacetimes

└ Time reversal

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

Schutz (2009, p. 258)

- now I use the static constraint to simplify the metric
- transformation
 - (0,0) term is $dt/d(-t)$
 - spatial terms are 1 if transformed to themselves
 - cross-terms are all zero, as coordinates independent of each other
- transformed metric
 - (0,0) term is unchanged, as -1 is squared
 - (r,r) term is unchanged, as transformation is 1
 - (0,r) term is negated, but must still be equal, so it's zero
 - no cross terms

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

Spherical stars

└ Static spacetimes

└ Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

Schutz (2009, p. 258)

- now I use the static constraint to simplify the metric
- transformation
 - (0,0) term is $dt/d(-t)$
 - spatial terms are 1 if transformed to themselves
 - cross-terms are all zero, as coordinates independent of each other
- transformed metric
 - (0,0) term is unchanged, as -1 is squared
 - (r,r) term is unchanged, as transformation is 1
 - (0,r) term is negated, but must still be equal, so it's zero
 - no cross terms

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

Spherical stars
└ Static spacetimes

└ Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

- now I use the static constraint to simplify the metric
- transformation
 - $(0,0)$ term is $dt/d(-t)$
 - spatial terms are 1 if transformed to themselves
 - cross-terms are all zero, as coordinates independent of each other
- transformed metric
 - $(0,0)$ term is unchanged, as -1 is squared
 - (r,r) term is unchanged, as transformation is 1
 - $(0,r)$ term is negated, but must still be equal, so it's zero
 - no cross terms

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Spherical stars

- Static spacetimes
 - The metric

- now we simplify the metric, since the cross term is zero
- we assume g_{00} to be negative, and g_{rr} to be positive
 - signature is $(-, +, +, +)$
 - holds inside stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*
 - $\Phi = \Lambda = 0 \implies e^{2\Phi} = e^{2\Lambda} = 1$ and $\mathbf{g} = \eta$

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Spherical stars

- Static spacetimes
 - The metric

- now we simplify the metric, since the cross term is zero
- we assume g_{00} to be negative, and g_{rr} to be positive
 - signature is $(-, +, +, +)$
 - holds inside stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*
 - $\Phi = \Lambda = 0 \implies e^{2\Phi} = e^{2\Lambda} = 1$ and $\mathbf{g} = \eta$

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Spherical stars

- Static spacetimes
- The metric

- now we simplify the metric, since the cross term is zero
- we assume g_{00} to be negative, and g_{rr} to be positive
 - signature is $(-, +, +, +)$
 - holds inside stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*
 - $\Phi = \Lambda = 0 \implies e^{2\Phi} = e^{2\Lambda} = 1$ and $\mathbf{g} = \eta$

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{00} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Spherical stars

└ Static spacetimes

└ Einstein Tensor

- now we can use the metric to derive the Riemann tensor
- from that the Einstein tensor
- the derivation is involved, so we will just take them as is
- we're going to use some of these components later on
- $x' \equiv dx/dr$

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

$$G_{00} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{00} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

2015-12-13

Spherical stars
└ Static spacetimes

└ Einstein Tensor

- now we can use the metric to derive the Riemann tensor
- from that the Einstein tensor
- the derivation is involved, so we will just take them as is
- we're going to use some of these components later on
- $x' \equiv dx/dr$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{00} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Schutz (2009, pp. 165, 260)

Static perfect fluid

2015-12-13

Spherical stars
└ Static perfect fluid

Static perfect fluid

- stars are fluids – for simplicity we assume perfect
- thus we will impose additional constraints accordingly



Four-velocity

Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for U^0

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for U_0

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-13

Spherical stars
 └ Static perfect fluid
 └ Four-velocity

- static fluid, so in MCRF three-velocity components all zero
- we find the only non-zero term, U^0 , by relating to the dot product
- lower it with the metric, to use in next part

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

Constraints

 $U^i = 0$ (static) $\vec{U} \cdot \vec{U} = -1$ (conservation law)

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Schutz (2009, p. 260)

Four-velocity

Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for U^0

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for U_0

$$U_0 = g_{00}U^0 = -e^{\Phi}$$



2015-12-13

Spherical stars
 └ Static perfect fluid
 └ Four-velocity

- static fluid, so in MCRF three-velocity components all zero
- we find the only non-zero term, U^0 , by relating to the dot product
- lower it with the metric, to use in next part

$$\begin{aligned} g_{00}U^0U^0 = -1 &\implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

Constraints	
$U^i = 0$ (static)	$\vec{U} \cdot \vec{U} = -1$ (conservation law)
Solving for U^0	
$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$	
$U_0 = g_{00}U^0 = -e^{\Phi}$	
Schutz (2009, p. 260)	

Four-velocity

Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for U^0

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for U_0

$$U_0 = g_{00}U^0 = -e^{\Phi}$$



2015-12-13

Spherical stars
 └ Static perfect fluid
 └ Four-velocity

- static fluid, so in MCRF three-velocity components all zero
- we find the only non-zero term, U^0 , by relating to the dot product
- lower it with the metric, to use in next part

$$\begin{aligned} g_{00}U^0U^0 = -1 &\implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

Constraints	
$U^i = 0$ (static)	$\vec{U} \cdot \vec{U} = -1$ (conservation law)
Solving for U^0	
$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$	
Solving for U_0	
$U_0 = g_{00}U^0 = -e^{\Phi}$	
Schutz (2009, p. 260)	

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{00} = \rho c^2$$

$T_{\alpha\beta}$ is diagonal

$$T_{0i} = (\rho + p)U_0U_i + pg_{0i} = 0$$

$$T_{ij} = (\rho + p)U_iU_j + pg_{ij} = pg_{ij}$$

2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{00} = \rho c^2$$

$T_{\alpha\beta}$ is diagonal

$$T_{0i} = (\rho + p)U_0U_i + pg_{0i} = 0$$

$$T_{ij} = (\rho + p)U_iU_j + pg_{ij} = pg_{ij}$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

 $T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Schutz (2009, p. 260)

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

 $T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Schutz (2009, p. 260)

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$



2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

 $T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Schutz (2009, p. 260)

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = 2\pi r^2 \sin^2 \theta$$



2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

 $T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Schutz (2009, p. 260)

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor
Stress-energy tensor for perfect fluid
$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$
Components of $T_{\alpha\beta}$
$T_{i\alpha} = pg_{i\alpha}$ $T_{\alpha\beta}$ is diagonal $T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$ $T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$
Schutz (2009, p. 260)

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

 $T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{ii} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Schutz (2009, p. 260)

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$



2015-12-13

Spherical stars
└ Static perfect fluid

└ Stress-energy tensor

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

 $T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

Schutz (2009, p. 260)

- $T_{i\alpha} = pg_{i\alpha}$ because spatial components of U are zero
- $T_{\alpha\beta}$ is diagonal because of previous condition and $g_{\alpha\beta}$ is diagonal
- T_{00} requires a little algebra
- T_{ii} just need to multiply metric by p
- $T_{\phi\phi}$ can be written in terms of $T_{\theta\theta}$

Equation of state

Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies



2015-12-13

Spherical stars
└ Static perfect fluid

└ Equation of state

- in a static fluid we have local thermodynamic equilibrium
- pressure a function of density and specific entropy
- specific entropy assumed negligibly small

Equation of state

Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies

Schutz (2009, p. 261)

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

2015-12-13

Spherical stars
└ Static perfect fluid

└ Equations of motion

- first equation follows from conservation of 4-momentum
- due to symmetry, the only non-trivial solution is for $\alpha = r$
- equation of motion for perfect fluid

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

• symmetries make only non-trivial solution $\alpha = r$
TODO: prove

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

Schutz (2009, pp. 175, 261)

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$



2015-12-13

Spherical stars

- └ Static perfect fluid

- └ Equations of motion

- first equation follows from conservation of 4-momentum
- due to symmetry, the only non-trivial solution is for $\alpha = r$
- equation of motion for perfect fluid

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

Schutz (2009, pp. 175, 261)

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$



2015-12-13

Spherical stars
└ Static perfect fluid

└ Equations of motion

- first equation follows from conservation of 4-momentum
- due to symmetry, the only non-trivial solution is for $\alpha = r$
- equation of motion for perfect fluid

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

• symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

Schutz (2009, pp. 175, 261)

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$m(r)$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-13

Spherical stars

└ Static perfect fluid

└ Mass function

- inspect $(0, 0)$ component of Einstein equations
- define the mass function, $m(r)$
- in Newtonian limit, $m(r)$ is mass within radius r

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} \left[r(1 - e^{-2\Lambda}) \right] = 8\pi \rho e^{2\Phi}$$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Schutz (2009, pp. 260–262)

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

 $m(r)$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-13

Spherical stars


└ Static perfect fluid

└ Mass function

- inspect $(0, 0)$ component of Einstein equations
- define the mass function, $m(r)$
- in Newtonian limit, $m(r)$ is mass within radius r

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable



Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

 $m(r)$

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-13

Spherical stars

└ Static perfect fluid

└ Mass function

- inspect $(0,0)$ component of Einstein equations
- define the mass function, $m(r)$
- in Newtonian limit, $m(r)$ is mass within radius r

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$m(r)$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

2015-12-13

Spherical stars

└ Static perfect fluid

└ Mass function

- inspect $(0,0)$ component of Einstein equations
- define the mass function, $m(r)$
- in Newtonian limit, $m(r)$ is mass within radius r

$$m(r) = 4\pi \int_0^r (r')^2 \rho(r') dr'$$

- doesn't work in GR, because total energy is not localizable

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

 $m(r)$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Schutz (2009, pp. 260–262)

$$\Phi(r)$$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\Phi(r)$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

2015-12-13

Spherical stars
 └ Static perfect fluid

$$\lrcorner \Phi(r)$$

- inspect (r, r) component of Einstein equations
- gives us an expression for $\Phi(r)$

$\Phi(r)$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

Schutz (2009, pp. 260–262)

$$\Phi(r)$$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\Phi(r)$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

2015-12-13

Spherical stars
└ Static perfect fluid

$$\lrcorner \Phi(r)$$

- inspect (r, r) component of Einstein equations
- gives us an expression for $\Phi(r)$

 $\Phi(r)$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

Schutz (2009, pp. 260–262)

$$\Phi(r)$$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\Phi(r)$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

2015-12-13

Spherical stars
└ Static perfect fluid

$$\lrcorner \Phi(r)$$

- inspect (r, r) component of Einstein equations
- gives us an expression for $\Phi(r)$

$$\Phi(r)$$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\Phi(r)$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

Schutz (2009, pp. 260–262)

Exterior Geometry

2015-12-13

Spherical stars
└ Exterior Geometry

Exterior Geometry

- until now, we've not considered whether we were inside or outside star
- properties inside different than outside (obviously)
- we're going to inspect both cases, starting with outside



Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$2\pi r^2 \frac{d\Phi(r)}{dr} = 4\pi r^2 p = 0$$

$$\frac{d\Phi(r)}{dr} = 0 \quad \Phi(r) = \frac{1}{2} \ln \left(\frac{r-2M}{r} \right)$$



2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
 - breaks down when matter surrounds star
- $m(r)$ is constant, we call it M
- $d\Phi/dr$ simplifies, and we can now integrate it to find $\Phi(r)$

$$\rho = p = 0$$

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
 - breaks down when matter surrounds star
- $m(r)$ is constant, we call it M
- $d\Phi/dr$ simplifies, and we can now integrate it to find $\Phi(r)$

Condition	
	$\rho = p = 0$
Consequences	
$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$	$m(r) \equiv M$
$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$	$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)} \quad \Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

Schutz (2009, pp. 262–263)

2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
 - breaks down when matter surrounds star
- $m(r)$ is constant, we call it M
- $d\Phi/dr$ simplifies, and we can now integrate it to find $\Phi(r)$

Schwarzschild metric I

Condition	
$\rho = p = 0$	
Consequences	
$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$	$m(r) \equiv M$
$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$	$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$

Schutz (2009, pp. 262–263)

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
 - breaks down when matter surrounds star
- $m(r)$ is constant, we call it M
- $d\Phi/dr$ simplifies, and we can now integrate it to find $\Phi(r)$

Schwarzschild metric I

Condition	
$\rho = p = 0$	
Consequences	
$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$	$m(r) \equiv M$
$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$	$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$

Schutz (2009, pp. 262–263)

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)} \quad \Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric I

- the external conditions just state we are in a vacuum
 - breaks down when matter surrounds star
- $m(r)$ is constant, we call it M
- $d\Phi/dr$ simplifies, and we can now integrate it to find $\Phi(r)$

Schwarzschild metric I

Condition	
$\rho = p = 0$	
Consequences	
$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$	$m(r) \equiv M$
$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$	$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$
Schutz (2009, pp. 262–263)	

Schwarzschild metric II

First two metric components

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric II

- recall g_{rr} from earlier
- substituting our expression from $\Phi(r)$ into $-e^{2\Phi}$ gives g_{00}
- we have found the Schwarzschild metric!

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Schwarzschild metric II

First two metric components

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$



2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric II

- recall g_{rr} from earlier
- substituting our expression from $\Phi(r)$ into $-e^{2\Phi}$ gives g_{00}
- we have found the Schwarzschild metric!

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Schwarzschild metric II

First two metric components

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$



2015-12-13

Spherical stars
└ Exterior Geometry

└ Schwarzschild metric II

- recall g_{rr} from earlier
- substituting our expression from $\Phi(r)$ into $-e^{2\Phi}$ gives g_{00}
- we have found the Schwarzschild metric!

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Far-field metric

Condition

$$r \gg M$$

Schwarzschild metric

$$ds^2 = \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$


$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-13

Spherical stars
└ Exterior Geometry

└ Far-field metric

- far-field metric of a star (far away)
 - mass M
 - distance R



Far-field metric

Condition

$$r \gg M$$

Far-field Schwarzschild metric

$$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx - \left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-13

Spherical stars

└ Exterior Geometry

└ Far-field metric

- far-field metric of a star (far away)
 - mass M
 - distance R

Condition	$r \gg M$
Schwarzschild metric	$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$
Far-field Schwarzschild metric (Cartesian)	$ds^2 \approx - \left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$
	$R^2 \equiv x^2 + y^2 + z^2$
Schutz (2009, pp. 263)	

Far-field metric

Condition

$$r \gg M$$

Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-13

Spherical stars
└ Exterior Geometry

└ Far-field metric

- far-field metric of a star (far away)
 - mass M
 - distance R

Far-field metric

Condition	$r \gg M$
Far-field Schwarzschild metric	$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$
Far-field Schwarzschild metric (Cartesian)	$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$
	$R^2 \equiv x^2 + y^2 + z^2$
Schutz (2009, pp. 263)	

Far-field metric

Condition

$$r \gg M$$

Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

2015-12-13

Spherical stars

└ Exterior Geometry

└ Far-field metric

- far-field metric of a star (far away)
 - mass M
 - distance R

Far-field metric

Condition	$r \gg M$
Far-field Schwarzschild metric	$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$
Far-field Schwarzschild metric (Cartesian)	$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$
	$R^2 \equiv x^2 + y^2 + z^2$
Schutz (2009, pp. 263)	

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)



2015-12-13

Spherical stars
└ Exterior Geometry

└ Birkhoff's Theorem

- the Schwarzschild metric generalizes to all spherically symmetric spacetimes in a vacuum
- George David Birkhoff

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)



2015-12-13

Spherical stars
└ Exterior Geometry

└ Birkhoff's Theorem

- the Schwarzschild metric generalizes to all spherically symmetric spacetimes in a vacuum
- George David Birkhoff

If the geometry of a given region of spacetime is:

- spherically symmetric

• a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)



2015-12-13

Spherical stars
└ Exterior Geometry

└ Birkhoff's Theorem

- the Schwarzschild metric generalizes to all spherically symmetric spacetimes in a vacuum
- George David Birkhoff

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- spherically symmetric
- a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)



2015-12-13

Spherical stars
└ Exterior Geometry

└ Birkhoff's Theorem

- the Schwarzschild metric generalizes to all spherically symmetric spacetimes in a vacuum
- George David Birkhoff

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- spherically symmetric
- a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)



2015-12-13

Spherical stars
└ Exterior Geometry

└ Birkhoff's Theorem

- the Schwarzschild metric generalizes to all spherically symmetric spacetimes in a vacuum
- George David Birkhoff

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- spherically symmetric
- a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff and Langer (1923)

Interior structure



Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$


2015-12-13

Spherical stars

└ Interior structure

└ Tolman–Oppenheimer–Volkov (T–O–V) equation

- inside a star, we cannot assume density and pressure are zero
- revisit two earlier equations
- substitute one into the other
- arrive at the T–O–V equation



Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-13

Spherical stars

└ Interior structure

└ Tolman–Oppenheimer–Volkov (T–O–V) equation

- inside a star, we cannot assume density and pressure are zero
- revisit two earlier equations
- substitute one into the other
- arrive at the T–O–V equation

Condition
$\rho \neq 0 \quad p \neq 0$
Recall
$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$ $\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
Schutz (2009, pp. 261–264)

Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-13

Spherical stars

└ Interior structure

└ Tolman–Oppenheimer–Volkov (T–O–V) equation

- inside a star, we cannot assume density and pressure are zero
- revisit two earlier equations
- substitute one into the other
- arrive at the T–O–V equation

Condition
$\rho \neq 0 \quad p \neq 0$
Recall
$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
Schutz (2009, pp. 261–264)

Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

2015-12-13

Spherical stars

└ Interior structure

└ Tolman–Oppenheimer–Volkov (T–O–V) equation

- inside a star, we cannot assume density and pressure are zero
- revisit two earlier equations
- substitute one into the other
- arrive at the T–O–V equation

Condition	$\rho \neq 0 \quad p \neq 0$
Recall	$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$ $\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$
T–O–V equation	$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$
Schutz (2009, pp. 261–264)	

System of equations

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

Mass function

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Equation of state

$$p = p(\rho)$$



2015-12-13

Spherical stars
└ Interior structure

└ System of equations

- T–O–V equation coupled with dm/dr and $p(\rho)$
 - 3 equations
 - 3 unknowns (m , ρ , p)
 - $\Phi(r)$ only intermediate variable

System of equations

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

Mass function

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Equation of state

$$p = p(\rho)$$

Schutz (2009, pp. 261–262, 264)

Newtonian hydrostatic equilibrium

Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

2015-12-13

Spherical stars
└ Interior structure

└ Newtonian hydrostatic equilibrium

- Newtonian limit reduces the T–O–V equation to HSE

Newtonian hydrostatic equilibrium

Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

Newtonian hydrostatic equilibrium

Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

2015-12-13

Spherical stars

└ Interior structure

└ Newtonian hydrostatic equilibrium

- Newtonian limit reduces the T–O–V equation to HSE

Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

References





G. D. Birkhoff and R. E. Langer. *Relativity and modern physics*. 1923.



S. M. Carroll. *Spacetime and geometry. An introduction to general relativity*. 2004.



C. J. Hansen and S. D. Kawaler. *Stellar Interiors. Physical Principles, Structure, and Evolution*. 1994.



C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. 1973.



B. Schutz. *A First Course in General Relativity*. May 2009.



N. Stergioulas. Rotating Stars in Relativity. *Living reviews in relativity*, 6:3, June 2003. [Online; accessed 2015-12-09]. DOI: 10.12942/lrr-2003-3. eprint: gr-qc/0302034.

