

Spherical solutions for stars

Daniel Wysocki

Rochester Institute of Technology

General Relativity I Presentations

December 14th, 2015



Introduction

- model stars using spherical symmetry
- Schwarzschild metric
- T–O–V equation
- applications

Spherically symmetric coordinates



Two-sphere in flat spacetime

General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

Two-sphere in flat spacetime

General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

Two-sphere in curved spacetime

Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to r

$$f(r', t) \equiv r^2$$

Two-sphere in curved spacetime


Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to r

$$f(r', t) \equiv r^2$$

Meaning of r




Mark Hannam

Figure:

Surface with circular symmetry but no coordinate $r = 0$.

- *not* proper distance from center
- “curvature” or “area” coordinate
 - radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$

Meaning of r



Mark Hannam


- *not* proper distance from center
- “curvature” or “area” coordinate
 - radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$

Figure:

Surface with circular symmetry but no coordinate $r = 0$.

Schutz (2009, p. 257)

Meaning of r



Mark Hannam

- *not* proper distance from center
- “curvature” or “area” coordinate
 - radius of curvature and area
- $r = \text{const}, t = \text{const}$
 - $A = 4\pi r^2$
 - $C = 2\pi r$

Figure:

Surface with circular symmetry but no coordinate $r = 0$.

Schutz (2009, p. 257)

Spherically symmetric spacetime

General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

g_{00} , g_{0r} , and g_{rr} functions of t and r

Static spacetimes



Motivation

- leads to simple derivation of Schwarzschild metric
- unique solution to spherically symmetric, asymptotically flat Einstein vacuum field equations (Birkhoff's theorem)

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

Motivation

- leads to simple derivation of Schwarzschild metric
- unique solution to spherically symmetric, asymptotically flat Einstein vacuum field equations (Birkhoff's theorem)

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

Definition

A spacetime is static if we can find a time coordinate t for which

(i) the metric independent of t

$$g_{\alpha\beta,t} = 0$$

(ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

Definition

A spacetime is static if we can find a time coordinate t for which

(i) the metric independent of t

$$g_{\alpha\beta,t} = 0$$

(ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{i}} = x^i_{,\bar{i}} = x^i_{,i} = 1$$

Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R^{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{00} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{00} = \frac{1}{r^2}e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Static perfect fluid



Four-velocity

Constraints

$$U^i = 0 \text{ (static)} \qquad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for U^0

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for U_0

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Four-velocity

Constraints

$$U^i = 0 \text{ (static)} \qquad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for U^0

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for U_0

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Four-velocity

Constraints

$$U^i = 0 \text{ (static)} \qquad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for U^0

$$g_{00}U^0U^0 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for U_0

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{00} = \rho g_{00}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)U^0U_0 + p(-1) = -\rho$$

$$T_{11} = (\rho + p)U^1U_1 + p(1) = p$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Stress–energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Stress–energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

Stress-energy tensor

Stress-energy tensor for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_{\alpha}U_{\beta} + pg_{\alpha\beta}$$

Components of $T_{\alpha\beta}$

$$T_{i\alpha} = pg_{i\alpha}$$

$T_{\alpha\beta}$ is diagonal

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{rr} = pe^{2\Lambda}, \quad T_{\theta\theta} = pr^2, \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

Equation of state

Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

Equations of motion

Conservation of 4-momentum

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution $\alpha = r$
TODO: prove

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$$m(r)$$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$$m(r)$$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$m(r)$

$$m(r) \equiv \frac{1}{2} r(1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Mass function

Einstein field equations

$$G_{00} = 8\pi T_{00} \implies \frac{1}{r^2} e^{2\Phi} \frac{d}{dr} [r(1 - e^{-2\Lambda})] = 8\pi \rho e^{2\Phi}$$

$m(r)$

$$m(r) \equiv \frac{1}{2} r (1 - e^{-2\Lambda}) \quad \text{or} \quad g_{rr} = e^{2\Lambda} \equiv \left(1 - \frac{2m(r)}{r} \right)^{-1}$$

Relation to energy density

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

$$\Phi(r)$$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\Phi(r)$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

$$\Phi(r)$$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

$$\Phi(r)$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

$\Phi(r)$

Einstein field equations

$$G_{rr} = 8\pi T_{rr} \implies -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi' = 8\pi p e^{2\Lambda}$$

 $\Phi(r)$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

Exterior Geometry



Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{u'(r)}{u(r)} = \frac{2\alpha(r) + 4\pi r^3 \rho}{r(r - 2\alpha(r))} = \frac{2\alpha(r)}{r(r - 2\alpha(r))} \quad \alpha(r) = \frac{r}{2} \ln \left(\frac{r - 2\alpha(r)}{r} \right)$$

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

Schwarzschild metric I

Condition

$$\rho = p = 0$$

Consequences

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho = 0$$

$$m(r) \equiv M$$

$$\frac{d\Phi(r)}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]} = \frac{M}{r(r - 2M)}$$

$$\Phi(r) = \frac{1}{2} \log \left(1 - \frac{2M}{r} \right)$$

Schwarzschild metric II

First two metric components

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \qquad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Schwarzschild metric II

First two metric components

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \qquad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Schwarzschild metric II

First two metric components

$$g_{rr} = e^{2\Lambda} = \left(1 - \frac{2M}{r}\right)^{-1} \quad g_{00} = -e^{2\Phi} = -\left(1 - \frac{2M}{r}\right)$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Far-field metric

Condition

$$r \gg M$$

Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

Far-field metric

Condition

$$r \gg M$$

Far-field Schwarzschild metric

$$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx - \left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

Far-field metric

Condition

$$r \gg M$$

Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

Far-field metric

Condition

$$r \gg M$$

Far-field Schwarzschild metric

$$ds^2 \approx -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 d\Omega^2$$

Far-field Schwarzschild metric (Cartesian)

$$ds^2 \approx -\left(1 - \frac{2M}{R}\right) dt^2 + \left(1 + \frac{2M}{R}\right) (dx^2 + dy^2 + dz^2)$$

$$R^2 \equiv x^2 + y^2 + z^2$$

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Birkhoff's Theorem

If the geometry of a given region of spacetime is:

- ① spherically symmetric
- ② a solution to the Einstein field equations in vacuum

then that geometry is necessarily a subset of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Interior structure



Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = - \frac{dp}{dr}$$

$$\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = - \frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

$$\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

$$\frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

Tolman–Oppenheimer–Volkov (T–O–V) equation

Condition

$$\rho \neq 0 \quad p \neq 0$$

Recall

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr} \qquad \frac{d\Phi}{dr} = \frac{m(r) + 4\pi r^3 p}{r[r - 2m(r)]}$$

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

System of equations

T–O–V equation

$$\frac{dp}{dr} = -\frac{(\rho + p)[m(r) + 4\pi r^3 p]}{r[r - 2m(r)]}$$

Mass function

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$

Equation of state

$$p = p(\rho)$$

Schutz (2009, pp. 261–262, 264)

Newtonian hydrostatic equilibrium

Newtonian limit

$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

Newtonian hydrostatic equilibrium

Newtonian limit







$$p \ll \rho; \quad 4\pi r^3 p \ll m; \quad m \ll r$$

Equation of hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{\rho m(r)}{r^2}$$

Schutz (2009, pp. 265–266) and Hansen and Kawaler (1994, p. 3)

References

-  G. D. Birkhoff and R. E. Langer. *Relativity and modern physics*. 1923.
-  S. M. Carroll. *Spacetime and geometry. An introduction to general relativity*. 2004.
-  C. J. Hansen and S. D. Kawaler. *Stellar Interiors. Physical Principles, Structure, and Evolution*. 1994.
-  C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. 1973.
-  B. Schutz. *A First Course in General Relativity*. May 2009.
-  N. Stergioulas. Rotating Stars in Relativity. *Living reviews in relativity*, 6:3, June 2003. [Online; accessed 2015-12-09]. DOI: 10.12942/lrr-2003-3. eprint: gr-qc/0302034.