

# Spherical solutions for stars

Daniel Wysocki

Rochester Institute of Technology

General Relativity I Presentations

December 14th, 2015



# Introduction



# Spherically symmetric coordinates



## Two-sphere in flat spacetime

## General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)$$

## Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2\theta d\phi^2) \equiv r^2 d\Omega^2$$



2015-12-12

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in flat spacetime

Two-sphere in flat spacetime

General metric:

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)$$

$$dl^2 = r^2(d\theta^2 + \sin^2\theta d\phi^2) \equiv r^2 d\Omega^2$$

Schutz (2009, p. 256)

## Two-sphere in flat spacetime

## General metric

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

## Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

2015-12-12

Spherical stars

└ Spherically symmetric coordinates

└ Two-sphere in flat spacetime

Two-sphere in flat spacetime

General metric:

$$ds^2 = -dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

Metric on 2-sphere

$$dl^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega^2$$

Schutz (2009, p. 256)

## Two-sphere in curved spacetime

## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

2015-12-12

## Spherical stars

## └ Spherically symmetric coordinates

## └ Two-sphere in curved spacetime

## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

## Two-sphere in curved spacetime

## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

2015-12-12

## Spherical stars

## └ Spherically symmetric coordinates

## └ Two-sphere in curved spacetime


## Metric on 2-sphere

$$dl^2 = f(r', t) d\Omega^2$$

Relation to  $r$ 

$$f(r', t) \equiv r^2$$

# Meaning of $r$



Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

2015-12-12

## Spherical stars

- └ Spherically symmetric coordinates
- └ Meaning of  $r$



Mark Hannam


- “curvature” or “area” coordinate
- radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)



# Meaning of $r$



Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:

Surface with circular symmetry but no coordinate  $r = 0$ .

2015-12-12

## Spherical stars

- └ Spherically symmetric coordinates
- └ Meaning of  $r$

Meaning of  $r$ 


Mark Hannam

- “curvature” or “area” coordinate
- radius of curvature and area
- *not* proper distance from center

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)

# Meaning of $r$



Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)

2015-12-12

## Spherical stars

- └ Spherically symmetric coordinates
  - └ Meaning of  $r$

Meaning of  $r$ 

Mark Hannam

- “curvature” or “area” coordinate
  - radius of curvature and area
- *not* proper distance from center
- $r = \text{const}, t = \text{const}$ 
  - $A = 4\pi r^2$
  - $C = 2\pi r$

Figure:  
Surface with circular  
symmetry but no  
coordinate  $r = 0$ .

Schutz (2009, p. 257)

# Spherically symmetric spacetime

## General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

$g_{00}$ ,  $g_{0r}$ , and  $g_{rr}$  functions of  $t$  and  $r$

2015-12-12

Spherical stars

└ Spherically symmetric coordinates

└ Spherically symmetric spacetime

Spherically symmetric spacetime

General metric

$$ds^2 = g_{00} dt^2 + 2g_{0r} dr dt + g_{rr} dr^2 + r^2 d\Omega^2$$

$g_{00}$ ,  $g_{0r}$ , and  $g_{rr}$  functions of  $t$  and  $r$

---

 Schutz (2009, p. 258)

# Static spacetimes



# Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Motivation

- Birkhoff's theorem says that the Schwarzschild metric applies to point 2

Motivation

- leads to simple derivation of Schwarzschild metric

• generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations

• Birkhoff's theorem

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

# Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Motivation

- Birkhoff's theorem says that the Schwarzschild metric applies to point 2

## Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

# Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Motivation

- Birkhoff's theorem says that the Schwarzschild metric applies to point 2

Motivation

- leads to simple derivation of Schwarzschild metric
- generalizes to spherically symmetric, asymptotically flat Einstein vacuum field equations
- Birkhoff's theorem

---

Schutz (2009, p. 263) and Misner, Thorne, and Wheeler (1973, p. 843)

# Definition

A spacetime is static if we can find a time coordinate  $t$  for which

- (i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ Definition

- a spacetime which only satisfies the first condition is *stationary*
  - e.g. rotating star

## Definition

A spacetime is static if we can find a time coordinate  $t$  for which

- (i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

- (ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$



# Definition

A spacetime is static if we can find a time coordinate  $t$  for which

(i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

(ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

## Spherical stars

- Static spacetimes
  - Definition

A spacetime is static if we can find a time coordinate  $t$  for which

(i) the metric independent of  $t$

$$g_{\alpha\beta,t} = 0$$

(ii) the geometry unchanged by time reversal

$$t \rightarrow -t$$

- a spacetime which only satisfies the first condition is *stationary*
  - e.g. rotating star

## Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^\alpha_{\bar{\alpha}} \Lambda^\beta_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

Spherical stars  
└ Static spacetimes

## └ Time reversal

- $g_{\bar{0}\bar{r}} = -g_{0r} = 0$

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^\alpha_{\bar{\alpha}} \Lambda^\beta_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{j}} = x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j$$

$$\Lambda^0_{\bar{i}} = x^0_{,\bar{i}} = x^0_{,i} = 0$$

## Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

# Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Time reversal

- $g_{\bar{0}\bar{r}} = -g_{0r} = 0$

Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

### Transformation

$$\Lambda^0_{\bar{0}} = x^0_{,\bar{0}} = -x^0_{,0} = -1$$

$$\Lambda^i_{\bar{j}} = x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j$$

$$\Lambda^0_{\bar{i}} = x^0_{,\bar{i}} = x^0_{,i} = 0$$

### Metric

$$g_{\bar{0}\bar{0}} = (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00}$$

$$g_{\bar{0}\bar{r}} = \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r}$$

$$g_{\bar{r}\bar{r}} = (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}$$

Schutz (2009, p. 258)

## Time reversal

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

Spherical stars  
└ Static spacetimes

## └ Time reversal

- $g_{\bar{0}\bar{r}} = -g_{0r} = 0$

$$\Lambda : (t, x, y, z) \rightarrow (-t, x, y, z)$$

$$g_{\bar{\alpha}\bar{\beta}} = \Lambda^{\alpha}_{\bar{\alpha}} \Lambda^{\beta}_{\bar{\beta}} g_{\alpha\beta} = g_{\alpha\beta}$$

## Transformation

$$\begin{aligned}\Lambda^0_{\bar{0}} &= x^0_{,\bar{0}} = -x^0_{,0} = -1 \\ \Lambda^i_{\bar{j}} &= x^i_{,\bar{j}} = x^i_{,j} = \delta^i_j \\ \Lambda^0_{\bar{i}} &= x^0_{,\bar{i}} = x^0_{,i} = 0\end{aligned}$$

## Metric

$$\begin{aligned}g_{\bar{0}\bar{0}} &= (\Lambda^0_{\bar{0}})^2 g_{00} = g_{00} \\ g_{\bar{0}\bar{r}} &= \Lambda^0_{\bar{0}} \Lambda^r_{\bar{r}} g_{0r} = -g_{0r} \\ g_{\bar{r}\bar{r}} &= (\Lambda^r_{\bar{r}})^2 g_{rr} = g_{rr}\end{aligned}$$

# The metric

## Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

## Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

## Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ The metric

- constraint  $g_{00} < 0 < g_{rr}$  holds for stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Schutz (2009, pp. 258–259)

# The metric

## Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

## Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

## Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ The metric

- constraint  $g_{00} < 0 < g_{rr}$  holds for stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Schutz (2009, pp. 258–259)

# The metric

## Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

## Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

## Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

2015-12-12

Spherical stars  
 └ Static spacetimes  
   └ The metric

- constraint  $g_{00} < 0 < g_{rr}$  holds for stars but not black holes
- limits at infinity tell us that spacetime is *asymptotically flat*

The metric

Simplified metric

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega^2$$

Replacement

$$g_{00} \rightarrow -e^{2\Phi}, \quad g_{rr} \rightarrow e^{2\Lambda}, \quad \text{provided } g_{00} < 0 < g_{rr}$$

Static spherically symmetric metric

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2 d\Omega^2$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \lim_{r \rightarrow \infty} \Lambda(r) = 0$$

Schutz (2009, pp. 258–259)

# Einstein Tensor

## General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

## Einstein tensor components

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Einstein Tensor

- $x' \equiv dx/dr$

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})]$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r]$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$



# Einstein Tensor

## General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

## Einstein tensor components

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

2015-12-12

Spherical stars  
└ Static spacetimes

└ Einstein Tensor

- $x' \equiv dx/dr$

Einstein Tensor

General Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$$

Einstein tensor components

$$G_{tt} = \frac{1}{r^2}e^{2\Phi}\frac{d}{dr}[r(1 - e^{-2\Lambda})],$$

$$G_{rr} = -\frac{1}{r^2}e^{2\Lambda}(1 - e^{-2\Lambda}) + \frac{2}{r}\Phi'$$

$$G_{\theta\theta} = r^2e^{-2\Lambda}[\Phi'' + (\Phi')^2 + \Phi'/r - \Phi'\Lambda' - \Lambda'/r],$$

$$G_{\phi\phi} = \sin^2\theta G_{\theta\theta}$$

Schutz (2009, pp. 165, 260)

# Static perfect fluid



# Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

## Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

## Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-12

Spherical stars  
 └ Static perfect fluid  
   └ Four-velocity

- “conservation law” is the conservation of four-momentum

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

### Constraints

$$U^i = 0 \text{ (static)}$$

$$\vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Schutz (2009, p. 260)

# Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

## Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

## Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-12

Spherical stars  
 └ Static perfect fluid  
   └ Four-velocity

- “conservation law” is the conservation of four-momentum

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

Four-velocity

Constraints	
$U^i = 0$ (static)	$\vec{U} \cdot \vec{U} = -1$ (conservation law)
Solving for $U^0$	
$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$	
$U_0 = g_{00}U^0 = -e^{\Phi}$	
Schutz (2009, p. 260)	

# Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

## Solving for $U^0$

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

## Solving for $U_0$

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

2015-12-12

Spherical stars  
 └ Static perfect fluid  
 └ Four-velocity

Four-velocity

## Constraints

$$U^i = 0 \text{ (static)} \quad \vec{U} \cdot \vec{U} = -1 \text{ (conservation law)}$$

Solving for  $U^0$ 

$$g_{00}(U^0)^2 = -1 \implies U^0 = (-g_{00})^{-1/2} = e^{-\Phi}$$

Solving for  $U_0$ 

$$U_0 = g_{00}U^0 = -e^{\Phi}$$

Schutz (2009, p. 260)

- “conservation law” is the conservation of four-momentum

$$\begin{aligned} g_{00}U^0U^0 &= -1 \implies (U^0)^2 = (-g_{00})^{-1} \\ &\implies U^0 = (-g_{00})^{-1/2} \\ &\implies U^0 = (e^{2\Phi})^{-1/2} = e^{-\Phi} \end{aligned}$$

# Stress-energy tensor

## $T_{\alpha\beta}$ for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

## Components of $T_{\alpha\beta}$

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{\alpha\beta} = 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii}$$

$$T_{rr} = pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Stress-energy tensor

- $T_{\alpha\beta} = 0$  because
  - the cross terms make  $g_{\alpha\beta} = 0$
  - $T_{0i}$  makes one of the  $U$ 's  $U_i = 0$
- likewise,  $T_{ii} = pg_{ii}$  because  $U_i U_i = 0$

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

$$\begin{aligned} T_{00} &= (\rho + p)U^0 U_0 + p(-e^{2\Phi}) = \rho e^{2\Phi} \\ T_{0i} &= 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii} \\ T_{rr} &= pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta \end{aligned}$$

# Stress-energy tensor

## $T_{\alpha\beta}$ for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

## Components of $T_{\alpha\beta}$

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{\alpha\beta} = 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii}$$

$$T_{rr} = pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Stress-energy tensor

- $T_{\alpha\beta} = 0$  because
  - the cross terms make  $g_{\alpha\beta} = 0$
  - $T_{0i}$  makes one of the  $U$ 's  $U_i = 0$
- likewise,  $T_{ii} = pg_{ii}$  because  $U_i U_i = 0$

Stress-energy tensor

 $T_{\alpha\beta}$  for perfect fluid

$$T_{\alpha\beta} = (\rho + p)U_\alpha U_\beta + pg_{\alpha\beta}$$

Components of  $T_{\alpha\beta}$ 

$$T_{00} = (\rho + p)e^{2\Phi} + p(-e^{2\Phi}) = \rho e^{2\Phi}$$

$$T_{\alpha\beta} = 0 \text{ for } \alpha \neq \beta; \quad T_{ii} = pg_{ii}$$

$$T_{rr} = pe^{2\Lambda}; \quad T_{\theta\theta} = pr^2; \quad T_{\phi\phi} = pr^2 \sin^2 \theta = T_{\theta\theta} \sin^2 \theta$$

Schutz (2009, p. 260)

## Equation of state

## Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies



2015-12-12

Spherical stars  
└ Static perfect fluid

└ Equation of state

Equation of state

Local thermodynamic equilibrium

$$p = p(\rho, S) \approx p(\rho)$$

- pressure related to energy density and specific entropy
- we often deal with negligibly small entropies

Schutz (2009, p. 261)



# Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO:** prove

## Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

2015-12-12

 Spherical stars  
 └ Static perfect fluid

└ Equations of motion

Equations of motion

Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

 • symmetries make only non-trivial solution  $\alpha = r$   
 TODO: prove

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

---

 Schutz (2009, pp. 175, 261)

# Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO: prove**

## Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO: prove**

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{d\rho}{dr}$$

# Equations of motion

## Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO: prove**

## Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Equations of motion

Equations of motion

Conservation laws

$$T^{\alpha\beta}_{;\beta} = 0$$

- symmetries make only non-trivial solution  $\alpha = r$   
**TODO: prove**

Equation of motion

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}$$

---

 Schutz (2009, pp. 175, 261)

## Equation of motion (continued)

**TODO**  
Show 10.31**TODO**

Show 10.31

2015-12-12

Spherical stars  
└ Static perfect fluid

└ Equation of motion (continued)



# Exterior Geometry



# Schwarzschild metric

TODO



2015-12-12

Spherical stars

└ Exterior Geometry

└ Schwarzschild metric

Schwarzschild metric

TODO

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

Let the geometry of a given region of spacetime:

- be spherically symmetric

• be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))



# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))



2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- be spherically symmetric
- be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

---

 Birkhoff and Langer (1923)

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))



2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- be spherically symmetric
- be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

# Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- ① be spherically symmetric
- ② be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))



2015-12-12

Spherical stars  
└ Exterior Geometry

└ Birkhoff's Theorem

Let the geometry of a given region of spacetime:

- be spherically symmetric
- be a solution to the Einstein field equations in vacuum.

Then that geometry is necessarily a piece of the Schwarzschild geometry.

(Proof given in Misner, Thorne, and Wheeler (1973, pp. 843–844))

# References





G. D. Birkhoff and R. E. Langer. *Relativity and modern physics*. 1923.



S. M. Carroll. *Spacetime and geometry. An introduction to general relativity*. 2004.



C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. 1973.



B. Schutz. *A First Course in General Relativity*. May 2009.



N. Stergioulas. Rotating Stars in Relativity. *Living reviews in relativity*, 6:3, June 2003. [Online; accessed 2015-12-09]. DOI: 10.12942/lrr-2003-3. eprint: gr-qc/0302034.



-  G. D. Birkhoff and R. E. Langer. *Relativity and modern physics*. 1923.
-  S. M. Carroll. *Spacetime and geometry. An introduction to general relativity*. 2004.
-  C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. 1973.
-  B. Schutz. *A First Course in General Relativity*. May 2009.
-  N. Stergioulas. Rotating Stars in Relativity. *Living reviews in relativity*, 6:3, June 2003. [Online; accessed 2015-12-09]. DOI: 10.12942/lrr-2003-3. eprint: gr-qc/0302034.