

My Study Notes 2024 / QFT ver.

Itsuki Miyane

Last modified: February 28, 2024

Contents

1	Introduction	2
2	Functional Method	3
3	Geometry of the Spacetime	4
A	Some notes	5
A.1	Normalization of Maxwell Lagrangian	5

1 Introduction

2 Functional Method

3 Geometry of the Spacetime

A Some notes

A.1 Normalization of Maxwell Lagrangian

The Maxwell lagrangian is the form

$$\mathcal{L} = N F^{\mu\nu} F_{\mu\nu} \quad (\text{A.1})$$

where $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$ is a field strength. We will determine the constant N to lagrangian contains the term

$$\mathcal{L} = \frac{1}{2} \dot{A}_1^2 + \frac{1}{2} \dot{A}_2^2 + \frac{1}{2} \dot{A}_3^2 + \dots \quad (\text{A.2})$$

Now expanding the term $F^{\mu\nu} F_{\mu\nu}$ carefully, we obtain

$$\begin{aligned} F^{\mu\nu} F_{\mu\nu} &= (\partial^\mu A^\nu - \partial^\nu A^\mu)(\partial_\mu A_\nu - \partial_\nu A_\mu) \\ &= 2((\partial^\mu A^\nu)(\partial_\mu A_\nu) - (\partial_\mu A^\nu)(\partial_\nu A_\mu)) \\ &= -2(\dot{A}_1^2 + \dot{A}_2^2 + \dot{A}_3^2) + \dots \end{aligned} \quad (\text{A.3})$$

and thus, the constant N satisfies the condition $-2N = 1/2$ and $N = -1/4$. That's why we get Maxwell Lagrangian as^{*1}

$$\mathcal{L} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} \quad (\text{A.4})$$

^{*1}aaa

References

- [AKOS12] H. Abe, T. Kobayashi, H. Ohki, and K. Sumita, *Superfield description of 10D SYM theory with magnetized extra dimensions*, *Nuclear Physics B* **863** (2012) 1–18, [arxiv:1204.5327 \[hep-ph, physics:hep-th\]](#).
- [PS95] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley Pub. Co, Reading, Mass, 1995.
- [WB92] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, Princeton University Press, Princeton, N.J, 1992.