

# My Study Notes 2024 / QFT

Itsuki Miyane

Last modified: March 11, 2024

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Functional Method</b>	<b>3</b>
<b>3</b>	<b>Geometry of the Spacetime</b>	<b>4</b>
<b>4</b>	<b>Perturbation Theory Anomalies</b>	<b>5</b>
4.1	Intuitive Example: The Axial Current in Two Dimensions . . . . .	5
<b>A</b>	<b>Some notes</b>	<b>6</b>
A.1	Normalization of Maxwell Lagrangian . . . . .	6

# **1 Introduction**

## **2 Functional Method**

### **3 Geometry of the Spacetime**

## **4 Perturbation Theory Anomalies**

In specific circumstances, quantum corrections can destroy symmetries of the classical equations of motion.

### **4.1 Intuitive Example: The Axial Current in Two Dimensions**

## A Some notes

### A.1 Normalization of Maxwell Lagrangian

The Maxwell lagrangian is the form

$$\mathcal{L} = N F^{\mu\nu} F_{\mu\nu} \quad (\text{A.1})$$

where  $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$  is a field strength. We will determine the constant  $N$  to lagrangian contains the term

$$\mathcal{L} = \frac{1}{2} \dot{A}_1^2 + \frac{1}{2} \dot{A}_2^2 + \frac{1}{2} \dot{A}_3^2 + \cdots . \quad (\text{A.2})$$

Now expanding the term  $F^{\mu\nu} F_{\mu\nu}$  carefully, we obtain

$$\begin{aligned} F^{\mu\nu} F_{\mu\nu} &= (\partial^\mu A^\nu - \partial^\nu A^\mu)(\partial_\mu A_\nu - \partial_\nu A_\mu) \\ &= 2((\partial^\mu A^\nu)(\partial_\mu A_\nu) - (\partial_\mu A^\nu)(\partial_\nu A_\mu)) \\ &= -2(\dot{A}_1^2 + \dot{A}_2^2 + \dot{A}_3^2) + \cdots \end{aligned} \quad (\text{A.3})$$

and thus, the constant  $N$  satisfies the condition  $-2N = 1/2$  and  $N = -1/4$ . That's why we get Maxwell Lagrangian as<sup>\*1</sup>

$$\mathcal{L} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} . \quad (\text{A.4})$$

---

<sup>\*1</sup>aaa