

Spring School 2024 @Izukawana

Moduli stabilization

on (for/at/in ?) supersymmetric magnetized D9-brane
model

Abe Lab. M1

Itsuki Miyane

Saturday, April 6th, 2024

*I will speak in Japanese though this slide is written in English.

Topics

- Review the senior thesis
- Report progress and difficulties I met

Introduction

Motivation

To solve the problems in the Standard Model, **higher dimensional models** were proposed.

Motivation

To solve the problems in the Standard Model, **higher dimensional models** were proposed.

In general, these theories contain extra fields related to

- the size and shape of the compactified extra dimension,
- hence the metric and the gravity

in its 4d effective field theory.

Motivation

To solve the problems in the Standard Model, **higher dimensional models** were proposed.

In general, these theories contain extra fields related to

- the size and shape of the compactified extra dimension,
- hence the metric and the gravity

in its 4d effective field theory.

Such a field is called **moduli fields**.

Moduli stabilization

The metrics in 10-dimensional spacetime are dynamical fields:

$$\begin{aligned} ds^{10} &= G_{MN} dX^M dX^N \\ &= g_{\mu\nu}(x, y) dx^\mu dx^\nu + g_{mn}(x, y) dy^m dy^n \end{aligned}$$

Thus, its vacuum expectation value (VEV) should be determined **by its dynamics**.

Moduli stabilization

The metrics in 10-dimensional spacetime are dynamical fields:


$$\begin{aligned} ds^{10} &= G_{MN} dX^M dX^N \\ &= g_{\mu\nu}(x, y) dx^\mu dx^\nu + g_{mn}(x, y) dy^m dy^n \end{aligned}$$

Thus, its vacuum expectation value (VEV) should be determined **by its dynamics**.

How?

What we have to do is just

- Write down the potential for moduli (e.g. g_{mn})
- Compute the minimum and identify the value $\langle g_{mn} \rangle$



These procedures are called **moduli stabilization**.

Purpose of my study

We will discuss the **moduli stabilization** on **magnetized torus model**.

.....

Magnetized torus model

Torus compactification and Magnetic flux

Torus compactification

- Compactifying 6d extra dimensions for three tori $(T^2)^3$.


 $\mathcal{A}^{(1)}$ $\mathcal{A}^{(2)}$ $\mathcal{A}^{(3)}$: area of the i -th torus

- $\mathcal{A}^{(i)}(x)$ are also **moduli** since it is related to the metric.

Torus compactification and Magnetic flux

Torus compactification

- Compactifying 6d extra dimensions for three tori $(T^2)^3$.




- $\mathcal{A}^{(i)}(x)$ are also **moduli** since it is related to the metric.

Magnetic flux

- Assigning the magnetic flux

$$M_a^{(i)} \quad (a = 1, 2)$$

for two gauge fields on each tori.



Finding the VEVs by fluxes

By magnetic fluxes in an extra dimension, moduli $\mathcal{A}^{(i)}(x)$ obtain its potential $V^{(D)}$.



Finding minima and determining the VEVs of $\langle \mathcal{A}^i \rangle$.

Finding the VEVs by fluxes

By **magnetic fluxes** in an extra dimension, **moduli** $\mathcal{A}^{(i)}(x)$ obtain its potential $V^{(D)}$.



Finding minima and determining the VEVs of $\langle \mathcal{A}^i \rangle$.

Potential by **magnetic fluxes**

$$F^{MN}F_{MN} = F^{\mu\nu}F_{\mu\nu} + \underbrace{F^{mn}F_{mn}} + \dots$$



$$V^{(D)} = \pi^2 \prod_i \mathcal{A}^i \times \left\{ \underbrace{\left(\sum_i \frac{M_1^{(i)}}{\mathcal{A}^{(i)}} \right)^2} + \underbrace{\left(\sum_i \frac{M_2^{(i)}}{\mathcal{A}^{(i)}} \right)^2} \right\}$$

Finding the VEVs by fluxes

By **magnetic fluxes** in an extra dimension, **moduli** $\mathcal{A}^{(i)}(x)$ obtain its potential $V^{(D)}$.



Finding minima and determining the VEVs of $\langle \mathcal{A}^i \rangle$.

Potential by **magnetic fluxes**

$$F^{MN}F_{MN} = F^{\mu\nu}F_{\mu\nu} + \underbrace{F^{mn}F_{mn}} + \dots$$



$$V^{(D)} = \pi^2 \prod_i \mathcal{A}^i \times \left\{ \underbrace{\left(\sum_i \frac{M_1^{(i)}}{\mathcal{A}^{(i)}} \right)^2} + \underbrace{\left(\sum_i \frac{M_2^{(i)}}{\mathcal{A}^{(i)}} \right)^2} \right\}$$

$$\longrightarrow \frac{M_a^{(1)}}{\langle \mathcal{A}^{(1)} \rangle} + \frac{M_a^{(2)}}{\langle \mathcal{A}^{(2)} \rangle} + \frac{M_a^{(3)}}{\langle \mathcal{A}^{(3)} \rangle} = 0 \quad \text{for } a = 1, 2$$

Finding the VEVs by fluxes

Relation of the VEVs $\langle \mathcal{A}^{(1)} \rangle$, $\langle \mathcal{A}^{(2)} \rangle$, $\langle \mathcal{A}^{(3)} \rangle$

$$\frac{M_a^{(1)}}{\langle \mathcal{A}^{(1)} \rangle} + \frac{M_a^{(2)}}{\langle \mathcal{A}^{(2)} \rangle} + \frac{M_a^{(3)}}{\langle \mathcal{A}^{(3)} \rangle} = 0 \quad \text{for } a = 1, 2$$

Finding the VEVs by fluxes

Relation of the VEVs $\langle \mathcal{A}^{(1)} \rangle$, $\langle \mathcal{A}^{(2)} \rangle$, $\langle \mathcal{A}^{(3)} \rangle$

$$\frac{M_a^{(1)}}{\langle \mathcal{A}^{(1)} \rangle} + \frac{M_a^{(2)}}{\langle \mathcal{A}^{(2)} \rangle} + \frac{M_a^{(3)}}{\langle \mathcal{A}^{(3)} \rangle} = 0 \quad \text{for } a = 1, 2$$

$$\longrightarrow M_a^{(1)} + M_a^{(2)} \frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(2)} \rangle} + M_a^{(3)} \frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(3)} \rangle} = 0$$

\downarrow

$$\frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(2)} \rangle} = \frac{M_1^{(3)} M_2^{(1)} - M_1^{(1)} M_2^{(3)}}{M_1^{(2)} M_2^{(3)} - M_1^{(3)} M_2^{(2)}}, \quad \frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(3)} \rangle} = -\frac{M_1^{(2)} M_2^{(1)} - M_1^{(1)} M_2^{(2)}}{M_1^{(2)} M_2^{(3)} - M_1^{(3)} M_2^{(2)}}$$

Finding the VEVs by fluxes

Relation of the VEVs $\langle \mathcal{A}^{(1)} \rangle$, $\langle \mathcal{A}^{(2)} \rangle$, $\langle \mathcal{A}^{(3)} \rangle$

$$\frac{M_a^{(1)}}{\langle \mathcal{A}^{(1)} \rangle} + \frac{M_a^{(2)}}{\langle \mathcal{A}^{(2)} \rangle} + \frac{M_a^{(3)}}{\langle \mathcal{A}^{(3)} \rangle} = 0 \quad \text{for } a = 1, 2$$

$$\longrightarrow M_a^{(1)} + M_a^{(2)} \frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(2)} \rangle} + M_a^{(3)} \frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(3)} \rangle} = 0$$

\downarrow

$$\frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(2)} \rangle} = \frac{M_1^{(3)} M_2^{(1)} - M_1^{(1)} M_2^{(3)}}{M_1^{(2)} M_2^{(3)} - M_1^{(3)} M_2^{(2)}}, \quad \frac{\langle \mathcal{A}^{(1)} \rangle}{\langle \mathcal{A}^{(3)} \rangle} = -\frac{M_1^{(2)} M_2^{(1)} - M_1^{(1)} M_2^{(2)}}{M_1^{(2)} M_2^{(3)} - M_1^{(3)} M_2^{(2)}}$$

The ratios of the moduli's VEVs are determined by the fluxes potential.

Summary so far

- We could stabilize the moduli
and obtain the ratio of its VEVs $\langle \mathcal{A}^{(1)} \rangle / \langle \mathcal{A}^{(2)} \rangle$ & $\langle \mathcal{A}^{(1)} \rangle / \langle \mathcal{A}^{(3)} \rangle$.

Summary so far

- We could stabilize the moduli
and obtain the ratio of its VEVs $\langle \mathcal{A}^{(1)} \rangle / \langle \mathcal{A}^{(2)} \rangle$ & $\langle \mathcal{A}^{(1)} \rangle / \langle \mathcal{A}^{(3)} \rangle$.
- But the **overall factor** T is undetermined.

$$T \propto \langle \mathcal{A}^{(1)} \rangle, \langle \mathcal{A}^{(2)} \rangle, \langle \mathcal{A}^{(3)} \rangle$$

Summary so far

- We could stabilize the moduli
and obtain the ratio of its VEVs $\langle \mathcal{A}^{(1)} \rangle / \langle \mathcal{A}^{(2)} \rangle$ & $\langle \mathcal{A}^{(1)} \rangle / \langle \mathcal{A}^{(3)} \rangle$.
- But the **overall factor T** is undetermined.

$$T \propto \langle \mathcal{A}^{(1)} \rangle, \langle \mathcal{A}^{(2)} \rangle, \langle \mathcal{A}^{(3)} \rangle$$

Introducing a potential that **has a different origin** than the magnetic fluxes
to stabilize **overall factor T** .

Determination of the overall factor

F -term potential

Effective potential for moduli T

- Its effective theory remains supersymmetric.
- Supersymmetric action is determined
by **super potential W** and **Kähler potential K** .
- We will study the following potential now[1]:

$$\begin{cases} W = w_0 - Ae^{-aT} + BX \\ K = -3 \ln(T + \bar{T}) + |X|^2 \end{cases}$$

Introducing a new scalar field X and w_0, A, B, a are parameters.


F -term potential

*We set the Planck constant as $M_{\text{Pl}}(\sim 2.4 \times 10^{18} \text{ GeV}) = 1$.

Scalar potential

$$V^{(F)} = e^K (K^{I\bar{J}} (D_I W)(D_{\bar{J}} \bar{W}) - 3|W|^2)$$

$$\begin{cases} D_I W \equiv \partial_I W + K_I W \\ K^{I\bar{J}}: \text{inverse matrix of } K_{I\bar{J}} \end{cases} \quad (I = X, T)$$



Parameters

$$w_0 \sim 2.17 \times 10^{-18}, \quad a = 4\pi^2, \quad A = 1, \quad B = e^{-4\pi^2}$$

$$\text{and } \langle X \rangle = \sqrt{3} - 1$$

$$\rightarrow \langle T \rangle \sim 1.085$$

Report progresses

In my senior thesis, we

- fix the value of the fluxes $M_a^{(i)}$
- and compute the area of the tori $\langle \mathcal{A}^{(i)} \rangle$ ($i = 1, 2, 3$).

Report progresses

In my senior thesis, we

- fix the value of the fluxes $M_a^{(i)}$
- and compute the area of the tori $\langle \mathcal{A}^{(i)} \rangle$ ($i = 1, 2, 3$).



During this vacation, I mainly studied the F -term potential more general form:

$$\begin{cases} W = w_0 - Ae^{-aT} + B \underbrace{e^{-bT}} \\ K = -3 \ln(T + \bar{T}) + |X|^2 - \frac{1}{\underbrace{\Lambda^2}} |X|^4 \end{cases}$$

Wavy factors are the difference from the previous one.

Report progresses

We can stabilize the moduli T , X but



Parameters

$$A = B = 1, a = b = 4\pi^2, w_0 = 10^{-17}, \Lambda = 10^{-4}$$

$$\longrightarrow \langle T \rangle \sim 1.07, \langle X \rangle \sim 10^{-8}, V_{\min} \sim 10^{-35}$$

Difficulties & Future Works

- We should tune the parameters A, B, a, b, w_0, Λ to **uplift the potential minima V_{\min} to the order $+10^{-120}$** .
- But it is difficult (at least for me) since

Difficulties & Future Works

- We should tune the parameters A, B, a, b, w_0, Λ to **uplift the potential minima V_{\min} to the order $+10^{-120}$** .
- But it is difficult (at least for me) since
I tried and/or will try following the method

Summary

Reference

- [1] H. Abe, T. Higaki, T. Kobayashi, and Y. Omura, *Moduli stabilization, F -term uplifting and soft supersymmetry breaking terms*, **Physical Review D** **75** (2007) 025019, [arxiv:hep-th/0611024](#).
- [2] J. Wess and J. Bagger, *Supersymmetry and Supergravity*. Princeton University Press, Princeton, N.J, 1992.