

Genetic and Functional Drivers of Diffuse Large B Cell Lymphoma¹

Anupama Reddy, Jenny Zhang, Nicholas S. Davis, ..., Jyotishka Datta, David B. Dunson, Sandeep S. Dave

Presented by Dominik Yang and Diana Lin

Diffuse Large B Cell Lymphoma (DLBCL)


- Most common type of non-Hodgkin lymphoma (NHL)²
- Under a microscope, lymphoma cells appear larger than normal lymphocytes²
- Most of the time, starts in the lymph nodes but it can also start in organs and tissues outside the lymph nodes²
- 30-40% of cases are diagnosed in stage 1 or 2 (localized)²
- Rest of the cases have spread to lymph nodes above and below the diaphragm or to liver, spleen, or bone marrow²
- Treated with R-CHOP chemotherapy³
 - **R**ituximab
 - **C**yclophosphamide
 - **H**ydroxydaunomycin
 - **O**ncovin
 - **P**rednisolone




Background

- Most common hematologic malignancy (annual incidence of over 100,000 cases worldwide)
- **Striking molecular and clinical heterogeneity makes it difficult to study this disease**
- Requires a large sample size to discover rare but important mutations

Goals


- Define the link between the functional impact of these genetic mutations to the growth of lymphoma cells
 - Ultimately be able to create therapeutics that target specific genetic drivers of DLBCL
- 

Discovery of Genetic Drivers



- Whole-exome sequencing of 1001 DLBCLs
- 150 possible driver genes identified
- Mean of 7.75 mutations per DLBCL patient
- Driver genes express common mutation patterns:
- Missense mutation and copy number gains - oncogene (BCL2, CARD11, IRF4)
- Truncating mutations and copy number losses - tumor suppressor (SPEN, CDKN2A, TNFAIP3)

Cell-of-Origin Effects



- RNA sequencing of gene expression in 775 patients to determine tumor cell of origin (A)
 - 313 activated B cell-like (ABC)
 - 331 germinal Center B cell-like (GCB)
 - 131 unclassified
- Validated by 2 complementary methods (B)
 - Nanostring assays
 - Immunohistochemistry-based Hans algorithm
- ABC DLBCLs had significantly worse overall survival than GCB DLBCLs



Cell-of-Origin Effects

Question:

Why do ABC DLBCLs have worse survival than GCB DLBCLs in response to standard therapy?



Connectivity of Driver Genes



- ABC and GCB DLBCLs share most driver genes
- 20 genes had significantly different mutations (C)
- Of the 150 driver genes, 61 were significantly related to other driver genes (D)
 - MLL2 mutations mostly exclusive with MYC
 - TP53 mutations mutually exclusive with KLHL6

Functional Genomics via CRISPR Screening

- 3 ABC, 2 GCB DLBCL, 1 BJAB (similar to GCB DLBCL)
- Selectively knock-out each gene in the DLBCLs to observe the effect on cell growth

QUESTION:


What did the authors expect to see from the effect of oncogenes vs tumor-suppressor genes?



Functional Genomics via CRISPR Screening

- Expected outcome: cells with oncogene-targeting sgRNAs would be depleted, cells expressing tumor-suppressor targeting sgRNA would be enriched

A



CRISPR Screen (sgRNAs)


| Gene A | Gene B | + ~20,000 Genes |
|-----------|------------|---------------------|
| [Red bar] | [Blue bar] | [Multi-colored bar] |

$$\text{sgRNA score} = \text{z-score} \left[\log_2 \left(\frac{\text{Day 5 sgRNA count}}{\text{Day 21 sgRNA count}} \right) \right]$$

CRISPR gene score = Avg(sgRNA scores for gene)


Functional Genomics via CRISPR Screening

- 1956 essential genes with significant role on cell fitness
- CRISPR score: degree of alteration in each gene as a function of altered abundance of sgRNAs targeting the gene




Functional Genomics via CRISPR Screening

c




- 35 driver genes identified as oncogenes
 - 9 sub-type specific ABC or GCB
- Four functional groups:
 - Signalling
 - Cell growth
 - B cell development
 - Transcription and translation
- Lethal effects of gene knockdown also affects non-mutated cells


Gene Expression Signature Associations with Genetic Alterations and Outcome



- Gene expression signatures with significant correlation fall into two classes:
 - Immune cell types (regulatory T cells, myeloid cells, NK cells)
 - Stromal connective tissue (epithelial and endothelial cells)
- *RHOA* and *MYC* mutations associated with proliferation-related signatures




Gene Expression Signature Associations with Genetic Alterations and Outcome



- No reliable association between signalling pathways and mutations in those pathways
 - Non-malignant cells may have affected effective measurements
- No association between stromal signatures and mutations (insignificant)

Gene Expression Signature Associations with Genetic Alterations and Outcome




- Gene sets related to cancer-related processes contained significantly more CRISPR-identified driver genes
 - DNA damage, oxidative phosphorylation, DNA replication, cell-cycle, RNA processing
- GCB DLBCL associated with better survival rate
- Stromal and immune signatures also associated with better survival rate
 - Regulatory T cells
- Proliferation signatures associated with worse survival rates
 - MYC/Nucleotide biosynthesis



Clinical Characteristics of DLBCL Driver Genes

A



- GCB DLBCL patients have a better survival probability than ABC DLBCL
- High expression of *MYC* & *BCL2* are associated with a worse prognosis and lower survival probability


Question:

What are the proteins that *MYC* & *BCL2* encode involved in?

B

Clinical Characteristics of DLBCL Driver Genes


A



B


- GCB DLBCL patients have a better survival probability than ABC DLBCL
- High expression of *MYC* & *BCL2* are associated with a worse prognosis and lower survival probability
 - *MYC*: codes for a transcription factor involved in processes of the cell cycle, **apoptosis**, etc.⁴
 - *BCL2*: codes Bcl-2 family of regulator proteins involved in the regulation of **apoptosis**⁴

Clinical Characteristics of DLBCL Driver Genes



- Alterations in *NF1* and *SGK1* were associated with more favourable survival
- Alterations in *MYC*, *CD79B* and *ZFAT* associated with poor survival

Clinical Characteristics of DLBCL Driver Genes



- Within ABC DLBCL...
 - Alterations in *KLHL14*, *BTG1*, *PAX5*, *CDKN2A* associated with significantly poorer survival
 - Alterations in *CREBBP* associated with favourable survival
- Within GCB DLBCL...
 - Alterations in *NCOR1* and *NFKBIA* are associated with poorer prognosis
 - Alterations in *EZH2*, *MYD88*, and *ARID5B* are associated with a significantly better prognosis

Clinical Characteristics of DLBCL Driver Genes

Question:

What do a lot of these genes have in common?



Clinical Characteristics of DLBCL Driver Genes

- *NF1*: codes for protein neurofibromin which acts as a **tumour suppressor** protein⁵
- *SGK1*: codes for Serine/Threonine Kinase, involved in regulating **cell proliferation** and **apoptosis**⁶
- *CD79B*: codes for Ig β protein of **B cell** antigen⁷
- *ZFAT*: codes a zinc finger protein⁸
- *KLHL14*: codes for adapter protein for certain **ubiquitin** ligases⁹
- *BTG1*: codes for **anti-proliferative** protein involved in regulating cell growth and **proliferation**¹⁰
- *PAX5*: codes for **B-cell** lineage specific activator protein¹¹
- *CDKN2A*: codes for a protein that stabilizes **tumour suppressor** protein p53¹²
- *CREBBP*: encodes CREB binding protein that is involved in growth control and **chromatin remodeling**
- *NCOR1*: codes for protein that promotes **chromatin condensation**¹³
- *NFKBIA*: codes for **NF κ B** inhibitor¹⁴
- *EZH2*: codes for histone methyltransferase involved in **transcriptional repression**¹⁵
- *MYD88*: codes for adaptor protein involved in **immune** cell signaling
- *ARID5B*: encodes a DNA binding protein that is involved in **transcription regulation**¹⁶

Comparison of Clinical vs Genomic Risk Model


- Clinical Risk Model

- Based on the International Prognostic Index (IPI) score¹⁸

- Age > 60 (1 pt)
 - Stage III or IV disease (1pt)
 - Elevated serum LDH (1 pt)
 - High levels of LDH (lactate dehydrogenase) indicate tissue damage and cellular destruction
 - WHO score of 2, 3 or 4 (1 pt)¹⁸
 - 2: symptomatic; <50% in bed during the day
 - 3: symptomatic; >50% in bed during the day but not bedbound
 - 4: bedbound
 - > 1 extranodal site (1 pt)




Comparison of Clinical vs Genomic Risk Model




- Genomic Risk Model
 - Based on genetic and molecular features
 - Known expression subgroups
 - ABC, GCB subgroups
 - *MYC*, *BCL2* high expression
 - Mutations and copy-number events in the 150 DLBCL associated genes identified in this study

Comparison of Clinical vs Genomic Risk Model




- Clinical Risk Model is highly prognostic in early stages (early mortality)
- Genomic Risk Model **better** at predicting long-term mortality (early and late mortality)

Comparison of Clinical vs Genomic Risk Model



- Early mortality of those that completely respond to treatment can be predicted by the Genomic Risk Model
- Indicates that genomic profile of a patient strongly influences their prognostic risk

Summary

- DLBCL is the most prevalent blood cancer in the world
 - Diverse number of mutations causing DLBCL make it difficult to understand the impact of the mutations for therapeutic reasons
 - 150 driver genes were identified, including 27 newly identified genes
 - Functional screening showed many possible therapeutically targetable genes
 - Can now distinguish which DLBCL patients would likely not benefit well from standard therapy, and can pursue new targeted therapies
 - Genomic risk model can be used clinically: measure cell of origin, BCL2 and MYC expression, targeted sequencing of driver genes
- 

Future Outlook

- Better understanding of the combined effects of the gene combinations and potential to aid in developing new therapeutic approaches
- Apply genomic methodologies to large cohorts and uncover patterns in a variety of other cancers with genetic heterogeneity that would not be discovered in small sample sizes
- Clinical and functional drivers of DLBCL identified → first step in improving outcomes for DLBCL




Strengths and limitations

- Pros:

- Large sample size
- Lots of statistical verification for their results
- Provides novel and valuable information that can be translated into a clinical setting

- Cons:

- A lot of their starting data is reliant on other researchers' data
 - There were many cases where the authors did not explain why they chose to do certain things
 - Requires a lot of statistical educational background to fully understand some tests they conducted
 - Requires a deep understanding of biological methodology and techniques to fully understand the paper on another level
- 

Questions?



References

1. Reddy A, Zhang J, Davis NS, et al. Genetic and Functional Drivers of Diffuse Large B+Cell Lymphoma. *CELL Cell*. 2017;171(2):481-494.e415.
2. Dearden C, Matutes E. Non-Hodgkin's lymphoma. *MPMED Medicine*. 2004;32(6):78-84.
3. Schmitz N, Zeynalova S, Nickelsen M, et al. CNS International Prognostic Index: A Risk Model for CNS Relapse in Patients With Diffuse Large B-Cell Lymphoma Treated With R-CHOP. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2016;34(26):3150-3156.
4. Miao Y, Hu S, Lu X, et al. Double-hit follicular lymphoma with MYC and BCL2 translocations: a study of 7 cases with a review of literature. *YHUPA Human Pathology*. 2016;58:72-77.
5. Susanne AMT, Lauren F, Margaret RW. Review Article : NF1 Mutations and Molecular Testing. *Journal of Child Neurology*. 2002;17(8):555-561.
6. Di Cristofano A. SGK1 : The Dark Side of PI3K Signaling. *BS:CTDB Current Topics in Developmental Biology*. 2017;123:49-71.
7. Chu PG, Arber DA. CD79: A Review. *Applied Immunohistochemistry & Molecular Morphology*. 2001;9(2):97-106.
8. Tsunoda T, Shirasawa S. Roles of ZFAT in haematopoiesis, angiogenesis and cancer development. *Anticancer research*. 2013;33(7):2833-2837.
9. Braybrook C, Warry G, Howell G, et al. Identification and characterization of KLHL4, a novel human homologue of the Drosophila Kelch gene that maps within the X-linked cleft palate and Ankyloglossia (CPX) critical region. *Genomics*. 2001;72(2):128-136.
10. Matsuda S, Rouault J, Magaud J, Berthet C. In search of a function for the TIS21/PC3/BTG1/TOB family. *FEBS letters*. 2001;497(2-3):2-3.
11. Medvedovic J, Ebert A, Tagoh H, Busslinger M. Pax5 : A Master Regulator of B Cell Development and Leukemogenesis. *BS:AI*. 2011;111:179-206.
12. Kannengiesser C, Avril MF, Spatz A, Laud K, Lenoir GM, Bressac-de-Paillerets B. CDKN2A as a uveal and cutaneous melanoma susceptibility gene. *Genes, chromosomes & cancer*. 2003;38(3):265-268.
13. Wang W, Song X-W, Bu X-M, Zhang N, Zhao C-H. PDCD2 and NCoR1 as putative tumor suppressors in gastric gastrointestinal stromal tumors. *Cell Oncol Cellular Oncology : The official journal of the International Society for Cellular Oncology*. 2016;39(2):129-137.
14. Geng P, Ou J, Li J, et al. Genetic Association Between NFKBIA -881A>G Polymorphism and Cancer Susceptibility. *Medicine*. 2015;94(31).
15. Crea F, Fornaro L, Bocci G, et al. EZH2 inhibition: targeting the crossroad of tumor invasion and angiogenesis. *Cancer Metastasis Rev Cancer and Metastasis Reviews*. 2012;31(3-4):753-761.
16. Gutiérrez-Camino, LÚpez-LÚpez E, Martín-Guerrero I, et al. Intron 3 of the ARID5B gene: a hot spot for acute lymphoblastic leukemia susceptibility. *Journal of cancer research and clinical oncology*. 2013;139(11):1879-1886.
17. Wilder RB, Rodriguez MA, Medeiros LJ, et al. International prognostic index-based outcomes for diffuse large B-cell lymphomas. *CNCR Cancer*. 2002;94(12):3083-3088.
18. Oken MM, Creech RH, Tormey DC, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *AMERICAN JOURNAL OF CLINICAL ONCOLOGY*. 1982;5(6):649-656.