

David Yang

Section 0.3 (Floating Point Representation of Real Numbers), Problem 3

Solution. First, note that the number 5 is represented in double precision floating point as

$$5 = 1.01 \times 2^2.$$

[illegible]
$$\begin{aligned} 2^k &\leq 2^{-52} \times 2^2 \\ &= 2^{-50}. \end{aligned}$$

Thus, the number $5 + 2^{-k}$ will be represented exactly in double precision floating arithmetic for any positive integer k from $\boxed{1 \text{ to } 50}$. ■

Section 0.3 (Floating Point Representation of Real Numbers), Problem 5(a)

Do the following sums by hand in IEEE double precision computer arithmetic, using the Rounding to Nearest Rule.

a) $(1 + (2^{-51} + 2^{-53}) - 1)$

Solution. Note that

[illegible]

[illegible]

and so

[illegible]

By the Rounding to the Nearest Rule, since the 53rd bit is 1, the 52nd bit is 0, so $2^{-51} + 2^{-53}$ rounds down to

[illegible]

[illegible]

[illegible]



Section 0.3 (Floating Point Representation of Real Numbers), Problem 6(a)

Do the following sums by hand in IEEE double precision computer arithmetic, using the Rounding to Nearest Rule.

a) $(1 + (2^{-51} + 2^{-52} + 2^{-54}) - 1)$

Solution. Note that

[illegible]

[illegible]

$$2^{-54} = 0.[\text{00}]01.$$

This means that

[illegible]

By the Rounding to the Nearest Rule, since the 53rd bit is 0, so $2^{-51} + 2^{-52} + 2^{-54}$ rounds down to

[illegible]

Does the associative law hold for IEEE computer addition?

$$\left(1 + \frac{\epsilon_{\text{mach}}}{2}\right) + \frac{\epsilon_{\text{mach}}}{2} \neq 1 + \left(\frac{\epsilon_{\text{mach}}}{2} + \frac{\epsilon_{\text{mach}}}{2}\right).$$

Identify for which values of x there is subtraction of nearly equal numbers, and find an alternate form that avoids the problem.

a) $\frac{1 - \sec x}{\tan^2(x)}$

Solution. Subtraction of nearly equal numbers occurs when

$$1 \approx \sec(x) = \frac{1}{\cos(x)}$$

which occurs when $\cos(x)$ is very close to 1. Values of x at which this occur include x which are close to $2\pi n$ for integer n .

We want to find an alternate form of the expression

$$\frac{1 - \sec x}{\tan^2 x}.$$

We can multiply both the numerator and denominator of the fraction by $1 + \sec x$, which gives

$$\begin{aligned} \frac{1 - \sec x}{\tan^2 x} &= \frac{1 - \sec x}{\tan^2 x} \cdot \frac{1 + \sec x}{1 + \sec x} \\ &= \frac{1 - \sec^2(x)}{\tan^2(x)(1 + \sec x)} \end{aligned}$$

Using the identity $\tan^2(x) = \sec^2(x) - 1$, we can rewrite the numerator as $-\tan^2(x)$. Thus, we get that

$$\begin{aligned} \frac{1 - \sec x}{\tan^2 x} &= \frac{1 - \sec^2(x)}{\tan^2(x)(1 + \sec x)} \\ &= \frac{-\tan^2(x)}{\tan^2(x)(1 + \sec x)} \\ &= -\frac{1}{1 + \sec x}. \end{aligned}$$

Thus, an alternate form of the expression $\frac{1 - \sec x}{\tan^2(x)}$ that avoids the potential problem of subtraction of nearly equal numbers is

$$\boxed{-\frac{1}{1 + \sec x}}.$$

■

Section 0.4 (Loss of Significance), Problem 3

Explain how to most accurately compute the two roots of the equation $x^2 + bx - 10^{-12} = 0$, where b is a number greater than 100.

Solution. The quadratic formula tells us that the roots of equation

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{b^2 + 4 \cdot 10^{-12}}}{2}.$$

Since $\sqrt{b^2 + 4 \cdot 10^{-12}} \approx b$ as $b \gg 10^{-12}$, the root

$$x = \frac{-b + \sqrt{b^2 + 4 \cdot 10^{-12}}}{2}$$

may lead to rounding error caused by the subtraction of nearly equal numbers.

As derived in Example 0.6, since b is positive, we can use the alternate form of the quadratic formula, which gives two roots

$$x_1 = -\frac{b + \sqrt{b^2 - 4ac}}{2a} \text{ and } x_2 = -\frac{2c}{(b + \sqrt{b^2 - 4ac})}.$$

Note that these forms avoid the rounding errors discussed above. Thus, in this instance, we can most accurately compute the two roots of the given equation by using the formulas

$$x_1 = -\frac{b + \sqrt{b^2 + 4 \cdot 10^{-12}}}{2} \text{ and } x_2 = \frac{2 \cdot 10^{-12}}{(b + \sqrt{b^2 + 4 \cdot 10^{-12}})}$$

■