
Homework 7

David Yang

Chapter 9 (The Fundamental Group) Problems.

Section 60 (Fundamental Groups of Some Surfaces), 60.2

Let X be the quotient space obtained from B^2 by identifying each point x of S^1 with its antipode $-x$. Show that X is homeomorphic to the projective plane P^2 .

Solution. ¹ Let π represent the quotient map from B^2 to X defined by identifying each point x of S^1 with its antipode. Consider the projection of the upper hemisphere of S^2 onto B^2 , a homeomorphism. Let h represent the inverse of this projection; h is a homeomorphism from B^2 to S^2 mapping the unit disk to the upper hemisphere. Finally, let p represent the canonical quotient map from S^2 to P^2 , defined by identifying each point of S^2 with its antipode.

■

¹constructed with Hillary's help.

Let C_1 and C_2 be disjoint simple closed curves in S^2 .

- a) **Show that $S^2 - C_1 - C_2$ has precisely three components. [Hint: If W_1 is the component of $S^2 - C_1$ disjoint from C_2 , and if W_2 is the component of $S^2 - C_2$ disjoint from C_1 , show that $\overline{W_1} \cup \overline{W_2}$ does not separate S^2 .]**

Solution. ² By the Jordan Curve Theorem, C_1 separates S^2 into two components, which we will denote as W_1 and V_1 . Similarly, by the Jordan Curve Theorem, C_2 separates S^2 into two components, which we will denote as W_2 and V_2 . Furthermore, define W_1 and W_2 to be the components matching the hint: W_1 is the component of $S^2 - C_1$ disjoint from C_2 and W_2 is the component of $S^2 - C_2$ disjoint from C_1 .

Note that since C_1 and C_2 are disjoint, it follows that $W_1 \subset V_2$ and $W_2 \subset V_1$, with $W_1 \cap W_2 = \emptyset$. Consequently, W_1 and W_2 are two components of $S^2 - C_1 - C_2$. Note that by construction, $V_1 = S^2 - \overline{W_1}$, and $V_2 = S^2 - \overline{W_2}$. Furthermore, C_1 and C_2 are disjoint, so $\overline{W_1} \cap \overline{W_2} = \emptyset$, and $S^2 - (\overline{W_1} \cap \overline{W_2}) = S^2$ is simply connected. Finally, neither $\overline{W_1}$ nor $\overline{W_2}$ separate S^2 , so by Theorem 63.3 (the general nonseparation theorem), $\overline{W_1} \cup \overline{W_2}$ does not separate S^2 . Thus, the third component of $S^2 - C_1 - C_2$ is precisely $\overline{W_1} \cup \overline{W_2} = V_1 \cap V_2$.

We conclude that $S^2 - C_1 - C_2$ has three components: W_1 , W_2 , and $V_1 \cap V_2$. ■

- b) **Show that these three components have boundaries C_1 and C_2 and $C_1 \cup C_2$, respectively.**

Solution. ³ By construction, the boundaries of W_1 and W_2 , two of the components of $S^2 - C_1 - C_2$, are C_1 and C_2 , respectively. It remains to show that the boundary of $V_1 \cap V_2$ is $C_1 \cup C_2$. Since $V_1 = S^2 - \overline{W_1}$, and $V_2 = S^2 - \overline{W_2}$, we have that

$$\begin{aligned} \overline{V_1} \cap \overline{V_2} - V_1 \cap V_2 &= ((S^2 - W_1) \cap (S^2 - W_2)) - ((S^2 - \overline{W_1}) \cap (S^2 - \overline{W_2})) \\ &= (\overline{W_1} \cup \overline{W_2}) - (W_1 \cup W_2) \\ &= (\overline{W_1} - W_1) \cup (\overline{W_2} - W_2) \\ &= C_1 \cup C_2. \end{aligned}$$

Thus, the boundary of $V_1 \cap V_2$ is by definition $C_1 \cup C_2$, as desired. ■

²constructed with Hillary's help.

³constructed with Hillary's help.