
Homework 1

Partial Differential Equations, Spring 2023

David Yang

Logan Chapter 1.1, Problem 6

Verify that

$$u(x, t) = \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds$$

is a solution to the wave equation $u_{tt} = c^2 u_{xx}$, where c is a constant and g is a given continuously differentiable function.

Solution. We will calculate the partial derivatives u_{tt} and u_{xx} separately. We will first calculate u_t and then u_{tt} . Note that

$$u_t = \frac{d}{dt} u(x, t) = \frac{d}{dt} \left(\frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds \right).$$

By Leibniz's rule, we know that

$$\frac{d}{dt} \left(\frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds \right) = \frac{1}{2c} \left[g(x+ct) \frac{d}{dt}(x+ct) - g(x-ct) \frac{d}{dt}(x-ct) \right].$$

Simplifying, we find that

$$u_t = \frac{1}{2c} [g(x+ct) \cdot c + g(x-ct) \cdot c] = \frac{g(x+ct) + g(x-ct)}{2}.$$

We can now take the partial of u_t with respect to t to find u_{tt} . We get that

$$u_{tt} = \frac{g'(x+ct) \frac{d}{dt}(x+ct) + g'(x-ct) \frac{d}{dt}(x-ct)}{2} = c \frac{g'(x+ct) - g'(x-ct)}{2}.$$

We will follow a similar procedure to calculate u_x and u_{xx} . We find that

$$u_x = \frac{d}{dx} u(x, t) = \frac{d}{dx} \left(\frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds \right).$$

Applying Leibniz's Rule, we get that

$$\frac{d}{dx} \left(\frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds \right) = \frac{1}{2c} \left[g(x+ct) \frac{d}{dx}(x+ct) - g(x-ct) \frac{d}{dx}(x-ct) \right].$$

Simplifying, we get that

$$u_x = \frac{g(x+ct) - g(x-ct)}{2c}.$$

Taking the partial of u_x with respect to x to determine u_{xx} , we find that

$$u_{xx} = \frac{g'(x+ct) \frac{d}{dx}(x+ct) - g'(x-ct) \frac{d}{dx}(x-ct)}{2c} = \frac{g'(x+ct) - g'(x-ct)}{2c}.$$

We see that

$$u_{tt} = c \frac{g'(x+ct) - g'(x-ct)}{2} = c^2 \left(\frac{g'(x+ct) - g'(x-ct)}{2c} \right) = c^2 u_{xx}$$

and so $u_{tt} = c^2 u_{xx}$, as desired. ■