

## Homework 4

David Yang

- 
1. Suppose  $V_1$  and  $V_2$  are independent  $\text{Gamma}(1, \lambda)$  random variables that represent waiting times in a Poisson process with rate  $\lambda$  events per unit time. Let  $X = V_1$  be the time of the first event and let  $Y = V_1 + V_2$  be the time of the second event.
- a) Find the joint CDF for  $X$  and  $Y$ :  $F_{xy}(x, y) = P(X \leq x, Y \leq y)$ . Hint: Graph the positive quadrant of the plane with axes  $V_1$  and  $V_2$ , and mark the region where  $V_1 \leq x$  and  $V_1 + V_2 \leq y$ . Integrate the joint pdf for  $V_1$  and  $V_2$  over this region to obtain the function  $F_{xy}(x, y)$ . Note that if there are restrictions on the arguments  $x$  and  $y$  that you do not specify, then you have failed to define the function.
- b) Show how to get the marginal CDF  $F_x$  by taking the upper limit for  $y$ , and  $F_y$  by taking the upper limit for  $x$ . Differentiate each marginal CDF to get the marginal pdfs.
- c) Show that taking partial derivatives of  $F_{xy}$  with respect to  $x$  and  $y$  yields the joint pdf:

$$\frac{\partial^2}{\partial x \partial y} F_{xy}(x, y) = \lambda^2 e^{-\lambda y} I(0 < x < y).$$

2. Suppose  $Z_1$  and  $Z_2$  have joint pdf

$$f_{12}(z_1, z_2) = \exp \left[ -\log(\pi) - 2(z_1^2 + z_2^2 + \sqrt{3}z_1z_2) \right].$$

- a) **Identify this as a bivariate Normal density by specifying the means  $\mu_1$  and  $\mu_2$ , standard deviations  $\sigma_1$  and  $\sigma_2$ , and the correlation  $\rho$ .**

*Solution.* Note that

$$\begin{aligned} f_{12}(z_1, z_2) &= \exp \left[ -\log(\pi) - 2(z_1^2 + z_2^2 + \sqrt{3}z_1z_2) \right] \\ &= \frac{1}{\pi} \exp \left[ -2(z_1^2 + z_2^2 + \sqrt{3}z_1z_2) \right]. \end{aligned}$$

Rewriting this expression, we have that

$$\begin{aligned} f_{12}(z_1, z_2) &= \frac{1}{\pi} \exp \left[ -2(z_1^2 + z_2^2 + \sqrt{3}z_1z_2) \right] \\ &= \frac{1}{2\pi \left(\frac{1}{2}\right)} \exp \left[ -\frac{1}{2 \left(\frac{1}{4}\right)} (z_1^2 + z_2^2 - 2 \left(-\frac{\sqrt{3}}{2}\right) z_1z_2) \right]. \end{aligned}$$

We recognize this as a bivariate Normal density, with

$$\boxed{\mu_1 = \mu_2 = 0, \sigma_1 = \sigma_2 = 1, \text{ and } \rho = -\frac{\sqrt{3}}{2}}. \quad \blacksquare$$

- b) **Any joint pdf may be expressed as a marginal pdf multiplied by a conditional pdf. Show that  $f_{12}(z_1, z_2)$  may be written as the product of a standard Normal density for  $Z_1$  and a Normal density for  $Z_2 \mid z_1$  that depends on  $z_1$ . Give the conditional mean and variance for  $Z_2 \mid z_1$  and show that they agree with the formulas  $\mathbb{E}[Z_2 \mid z_1] = \beta_0 + \beta_1 z_1$  with  $\beta_1 = \rho \frac{\sigma_2}{\sigma_1}$ ,  $\beta_0 = \mu_2 - \beta_1 \mu_1$ , and  $\text{Var}[Z_2 \mid z_1] = (1 - \rho^2) \sigma_2^2$ .**

*Solution.* Let  $Z_1 \sim N(0, 1)$ . We know that  $f_{Z_1}(z_1) = \frac{1}{\sqrt{2\pi}} \exp \left[ -\frac{z_1^2}{2} \right]$ .

$$\begin{aligned} f_{12}(z_1, z_2) &= \frac{1}{\pi} \exp \left[ -2(z_1^2 + z_2^2 + \sqrt{3}z_1z_2) \right] \\ &= \frac{1}{\sqrt{2\pi}} \exp \left[ -\frac{z_1^2}{2} \right] \frac{\sqrt{2}}{\sqrt{\pi}} \exp \left[ -\left( \frac{3}{2}z_1^2 + 2z_2^2 + 2\sqrt{3}z_1z_2 \right) \right] \end{aligned}$$

Recognizing the first term as  $f_{Z_1}(z_1)$ , the pdf for  $Z_1$ , we have that

$$f_{12}(z_1, z_2) = f_{Z_1}(z_1) \frac{\sqrt{2}}{\sqrt{\pi}} \exp \left[ -\left( \frac{3}{2}z_1^2 + 2z_2^2 + 2\sqrt{3}z_1z_2 \right) \right].$$

Rearranging the second term and simplifying, we get that

$$\begin{aligned} f_{12}(z_1, z_2) &= f_{Z_1}(z_1) \frac{\sqrt{2}}{\sqrt{\pi}} \exp \left[ -\left( \frac{3}{2}z_1^2 + 2z_2^2 + 2\sqrt{3}z_1z_2 \right) \right] \\ &= f_{Z_1}(z_1) \frac{\sqrt{2}}{\pi} \exp \left[ -2 \left( z_2^2 + \sqrt{3}z_1z_2 + \frac{3}{4}z_1^2 \right) \right] \\ &= f_{Z_1}(z_1) \frac{1}{\sqrt{2\pi \cdot \frac{1}{4}}} \exp \left[ -\frac{\left( z_2 + \frac{\sqrt{3}}{2}z_1 \right)^2}{2 \left( \frac{1}{4} \right)} \right]. \end{aligned}$$

We recognize the second term now as the pdf for  $Z_2 \mid z_1 \sim N\left(-\frac{\sqrt{3}}{2}z_1, \frac{1}{4}\right)$ , so we conclude

$$f_{12}(z_1, z_2) = f_{Z_1}(z_1)f_{Z_2|z_1}(z_2 \mid z_1).$$

Note that  $\mathbb{E}[Z_2 \mid z_1] = -\frac{\sqrt{3}}{2}z_1$ , and

$$\begin{aligned}\beta_0 + \beta_1 z_1 &= \left(\mu_2 - \left(\rho \frac{\sigma_2}{\sigma_1}\right) \mu_1\right) + \rho \frac{\sigma_2}{\sigma_1} z_1 \\ &= (0 - 0) - \frac{\sqrt{3}}{2} z_1 \\ &= -\frac{\sqrt{3}}{2} z_1.\end{aligned}$$

Similarly,  $\text{Var}[Z_2 \mid z_1] = \frac{1}{4}$  and

$$\begin{aligned}(1 - \rho^2) \sigma_2^2 &= \left(1 - \left(-\frac{\sqrt{3}}{2}\right)^2\right) 1^2 \\ &= \frac{1}{4},\end{aligned}$$

matching the results from the given formulas. ■

- c) **You can also show conditional results using representation. For  $Z_o \sim N(0, 1)$  independent of  $Z_2$ , define  $Z_1 = \rho Z_2 + \sqrt{1 - \rho^2} Z_o$  to have correlation  $\rho$  with  $Z_2$ . Show that conditioning on  $Z_2 = z_2$  and treating this as constant in the representation of  $Z_1$  results in a conditional distribution  $Z_1 \mid z_2$  that mirrors that of  $Z_2 \mid z_1$  from part (b).**

*Solution.* For  $Z_o \sim N(0, 1)$  independent of  $Z_2$ , define  $Z_1 = \rho Z_2 + \sqrt{1 - \rho^2} Z_o$  as stated. Note that  $\sqrt{1 - \rho^2} Z_o \sim N(0, 1 - \rho^2)$ . Conditioning on  $Z_2 \mid z_2$ , we find that

$$Z_1 \mid z_2 \sim N(\rho Z_2, 1 - \rho^2)$$

For  $\rho = -\frac{\sqrt{3}}{2}$  (giving  $1 - \rho^2 = \frac{1}{4}$ ), we have

$$Z_1 \mid z_2 \sim N\left(-\frac{\sqrt{3}}{2} z_2, \frac{1}{4}\right)$$

which matches the distribution of  $Z_2 \mid z_1$  in part (b). ■

3. Suppose  $X$  and  $Y$  have joint pdf  $f_{xy}(x, y) = I(0 < x < 1, -x < y < x)$ .

- a) **Explain how you can tell, without finding the marginal densities, that the conditional densities are Uniform. Write out the conditional densities  $f_{x|y}(x | y)$  and  $f_{y|x}(y | x)$ .**

*Solution.* The conditional densities are uniform as they are proportional to the joint density, which can be thought of as Uniform (as it is an indicator variable).

Since by construction,  $|y| < x$  and  $x < 1$  it follows that the conditional density is

$$f_{x|y}(x | y) = \frac{1}{1 - |y|} I(|y| < x < 1).$$

Similarly, we must have that  $-x < y < x$ , so

$$f_{y|x}(y | x) = \frac{1}{2x} I(-x < y < x). \quad \blacksquare$$

- b) **Explain how you can tell, without finding the marginal densities, that  $X$  and  $Y$  are not independent. Find the marginal pdf's  $f_x(x)$  and  $f_y(y)$  and verify that  $f_{xy}(x, y) \neq f_x(x)f_y(y)$ .**

*Solution.* Note that very clearly,  $f_{x|y}(x, y)$  are not equal for different values of  $y$ , so  $X$  and  $Y$  cannot be independent. We can verify this with the marginal pdfs:

$$f_x(x) = \frac{f_{xy}(x, y)}{f_{y|x}(y | x)} = \frac{I(0 < x < 1, -x < y, x)}{\frac{1}{2x} I(-x < y < x)} = 2x I(0 < x < 1).$$

Similarly,

$$f_y(y) = \frac{f_{xy}(x, y)}{f_{x|y}(x | y)} = \frac{I(0 < x < 1, -x < y, x)}{\frac{1}{1 - |y|} I(|y| < x < 1)} = (1 - |y|) I(-1 < y < 1).$$

We check that

$$\begin{aligned} f_X(x)f_Y(y) &= 2x(1 - |y|)I(0 < x < 1)I(-1 < y < 1) \\ &\neq I(0 < x < 1, -x < y, x) = f_{XY}(x, y) \end{aligned}$$

confirming that  $X$  and  $Y$  are not independent.  $\blacksquare$

- c) **Show that  $X$  and  $Y$  are uncorrelated.**

*Solution.* Note that

$$\mathbb{E}[XY] = \int_0^1 \int_{-x}^x xy I(0 < x < 1, -x < y < x) dy dx = 0$$

by symmetry. Similarly,

$$\mathbb{E}[X] = \int_0^1 x(2x) dx = 1$$

and

$$\mathbb{E}[Y] = \int_{-1}^1 y(1 - |y|) dy = 0$$

by symmetry. Thus, we have that

$$\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] = 0$$

so  $X$  and  $Y$  are uncorrelated, as desired.  $\blacksquare$

4. a) Suppose  $X_1$  and  $X_2$  are Bernoulli random variables with expectations  $p_1$  and  $p_2$ . Show that  $X_1$  and  $X_2$  are independent if and only if they are uncorrelated. This shows the Bernoulli distribution is special like the multivariate Normal distribution in that uncorrelated implies independence.
- b) Suppose  $Y = X_1 + X_2$  with  $X_1$  and  $X_2$  independent. If you learn that  $Y$  and  $X_1$  are Normal variables, prove that  $X_2$  is also a Normal random variable.