





A Molecular Dynamics Approach to Study Seed-Region Modifications in siRNA


Dylan J. Anderson, Cameron M. MacKenzie, Glenn A. Burley, David S. Palmer

Pure and Applied Chemistry, University of Strathclyde, 295 Cathedral Street, Glasgow, G1 1XL, UK




Background

RNA therapeutics offer an alternative mechanism of action to traditional small molecule drugs. Their target is dependent on **base sequence**, and their properties can be independently optimised by introducing **chemical modifications** to the backbone, sugar, and base.¹




Small interfering RNAs (siRNAs) are loaded into the **RNA-induced silencing complex** to facilitate the binding of an **mRNA target** and inhibit gene expression. This is a catalytic mechanism (RNA interference) where target binding initiates at the guide strand's seed-region: residues 2-8 from the 5'-end.^{2,3}




Six of seven marketed siRNA therapeutics employ chemical modifications (cm) to improve binding affinities, physiological stability, and reduce off-targets.

Molecular dynamics study of cm-siRNAs is hindered by the lack of unnatural residue force-field parameters. In response, toolkits like modXNA have been developed to calculate and assemble those missing values.⁵

This study assesses the impact of position-specific cm's on the structural dynamics, and binding free-energies, of an 8-residue siRNA sequence targeting HTT mRNA.




Methods




Modifications

Added to 2, 5, or 7 position (25 variants)




Global Analysis

Across the 25 sequence variants, RMSD values were consistent and indicative of the maintenance of duplex structure. **Per-residue RMSF's** and **hydrogen-bond distances** between Watson-Crick base-pairs convey weaker terminal base-pairing (especially at the 5'-region) due to A/U repeats and/or reduced base-stacking.⁶





Strong agreement shown between rankings of free-energies and enthalpies from MMGBSA binding energies, however, sequences modified with **DNA** at the **2nd** (s14) and **5th** (s15) residues had a significant difference in **ranking placement**.




Modification Analysis

Binding affinities followed the expected literature trend for the key stabilising and destabilising residues. The unforeseen outlier was the weak binding of **MOE** variants. As the implicit solvent of MMGBSA won't capture the known crystal water interactions, lower solvation energies reduced affinity.⁷



LNA's demonstrated strong binding affinities, particularly at positions **2** and **7**, due to favourable enthalpies from hydrogen-bonding and base-stacking interactions. The simulations also replicated the tight inter-base **C3'-endo pseudoangle** (favourable for A-RNA, circ.) and **large pucker amplitude** (rad.).⁸

Cytosine-mismatches were among the weakest binders: unsurprising, as they employ non-canonical Watson-Crick base-pairs. Position **2 cmis** had the most penalising binding enthalpy, along with: global instability (largest avg. RMSD), occupation of the A-RNA competing **C2'-endo pseudoangle** (**144° to 180°**), and movement of the **residue 2 χ -angle** towards the syn-conformation (**90° to -90°**).⁹



Conclusions

- modXNA parameters were easy to generate and output stable sims. of cm-siRNAs with the expected trend in per-residue deviations and H-bond distances.
- The strongest and weakest binders at choice seed positions were predicted using this workflow, and the properties of global/local deviations, base-pair and sugar-angles, and thermodynamics, were consistent with literature observations.
- Due to the misalignment between free-energy and enthalpic rankings for DNA-modified residues, NMA-entropies were considered necessary for comparisons.
- A hybrid implicit/explicit solvent method may be required to reproduce specific water-interactions seen in the crystal-structures of modified oligonucleotides.

Future work


- An experimental component: to determine binding thermodynamics (melting points) and kinetics (on/off rates) for seed-region modifications, and compare to the aforementioned computational output.
- The incorporation of the hAgo2 protein (from the RISC) in molecular dynamics simulations, in addition to a range of cm's, to probe their effects on structure and energetics in a biologically-relevant context.
- Investigating machine-learning algorithms, and fingerprinting methods appropriate for the representation of cm-nucleotides, to predict binding affinities and thus accelerate the experimental pipeline.

Acknowledgements

We would like to thank: the ARCHIE-WeSt team for computational resources; Drs. Otto Linden and Andrea Taladriz-Sender, and Theo Minetto, for ongoing feedback and support; Dr. Ross Urquhart and Jack Rodden for help in coding and implementation; Profs. DSP and GAB for the project opportunity and supervision. LNA graphic generated using UCSF ChimeraX software. Funding provided by the University of Strathclyde.

References

- Nat. Biotechnol. **2017**, 35, 238; 2. Front. Pharmacol. **2019**, 10, 444; 3. Health Sci. Rev. **2024**, 11, 100157; 4. Signal Transduct. Target. Ther. **2020**, 5, 101; 5. J. Chem. Theory Comput. **2024**, 20, 9354; 6. J. Am. Chem. Soc., **2006**, 128, 6885; 7. Nucleic Acids Res., **1998**, 26, 3694–3699; 8. Datta et al, Chem. Commun., **2024**, 60, 13024; 9. Nucleic Acids Res., **2020**, 48, e74



Links and Contact Info.
dylan.anderson@strath.ac.uk