

DATA 308: Lab 5b

Hypothesis Testing with SLR

Dylan Wikel

Our research question that we will investigate in this lab is:

Among first-year Elmhurst College students represented by these data, is family income linearly associated with mean gift aid received?

```
library(openintro)
library(broom)
library(tidyverse)
data("elmhurst")
```

Question 1

Take a look at the `elmhurst` dataset and familiarize yourself with it. You can run the code chunk below to view help documentation for this dataset and the structure. I have the `eval` argument chunk set to false, meaning when you render this document the code chunk will not be run.


```
?elmhurst
glimpse(elmhurst)
```

Question 2

Produce a plot to visualize the relationship between gift aid and family income. Make sure to plot the response variable on the y-axis and include a regression line.

Response

```
ggplot(elmhurst, aes(x = family_income, y = gift_aid)) +
  geom_point() +
  geom_smooth(method = "lm", color = "green")
```



Question 3

We will learn about the conditions for linear regression next week, but for now just visually assess, does a line appear to be a good model?

Response

It doesn't appear to be a particularly good model, but it is reasonable to say that there's a negative relationship that is somewhat linear. Loosely.

Question 4

Below is the population simple linear regression model, in a general form written with LaTeX. Replace the Y and X in the equation with the response and explanatory variables for this context.

Response

$$\text{Gift Aid Received} = \beta_0 + \beta_1(\text{Family Income}) + \epsilon$$

Question 5

State the null and alternative hypotheses, in symbols and in words, to test our research question stated at the top of this document.

Response

There is no linear association between family income and the amount of gift aid a student receives.

$$H_0 = \beta_1 = 0$$

There is a linear association between family income and the amount of gift aid a student receives.

$$H_a = \beta_1 \neq 0$$

Question 6

Fit a simple linear regression model.

Response

```
aid_model <- lm(gift_aid ~ family_income, data = elmhurst)
tidy_model <- tidy(aid_model)

summary(aid_model)
```

Call:

```
lm(formula = gift_aid ~ family_income, data = elmhurst)
```

Residuals:

Min	1Q	Median	3Q	Max
-10.1128	-3.6234	-0.2161	3.1587	11.5707

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	24.31933	1.29145	18.831	< 2e-16 ***
family_income	-0.04307	0.01081	-3.985	0.000229 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.783 on 48 degrees of freedom

Multiple R-squared: 0.2486, Adjusted R-squared: 0.2329

F-statistic: 15.88 on 1 and 48 DF, p-value: 0.0002289

Question 7

What is the p-value for this hypothesis test? With a significance level of 0.05, what is your statistical decision?

Response

```
p_value <- tidy_model |>
  filter(term == "family_income") |>
  select(p.value)
```

```
p_value
```

```
# A tibble: 1 x 1
  p.value
  <dbl>
1 0.000229
```

$$p = 0.0002289$$

$$p < 0.05$$

With a 0.00023 we can infer that `family_income` is a significant predictor of `gift_aid`.

Question 8

We should not just report a p-value on its own, we should also provide context. Compute a confidence interval for β_1 .

Response

Question 9

Provide your conclusion in context for this hypothesis test. One sentence should explain the results of your hypothesis test, with p-value reported in parentheses. The next sentence should provide your estimate and confidence interval interpreted in context (you can choose to report it as an interval or as an margin of error).

Response

Final note

A condition on our conclusion:

Our hypothesis test is valid only if certain conditions about the data and regression model are satisfied. Next week, we will learn how to identify these conditions and use diagnostic plots to assess them. For now, our conclusion assumes that the required conditions hold.