

SEVEN OPEN UNIT EQUILATERAL TRIANGLES DO NOT COVER A REGULAR UNIT HEXAGON

JINEON BAEK, JEEWON KIM

ABSTRACT. We prove that the regular hexagon of side length one cannot be covered by seven open equilateral triangles of side length one. Equivalently, no regular hexagon of side length greater than one is covered by seven closed unit equilateral triangles. Seven distinguished points force one center role and six vertex roles. An exhaustive classification of those roles reduces the proof to four mechanisms: direct boundary, radial, diagonal, and conditional-skeleton length sums; exact propagation of boundary and radial demands; normalized area loss; and a center-independent direct nine-point obstruction. The last mechanism forces six radial and three asymmetric witnesses into the center role. A reader-facing guide first gives the entire proof architecture; simplified body sections then present the geometric mechanisms and terminal implications, while the complete formula catalogues and exact certificates appear in technical appendices. The terminal mixed-overlap signs are certified by exact rational reductions and computer-verified global Bernstein identities, with no interval subdivision or branch-and-bound.

CONTENTS

1. A Guide to the Proof	3
1.1. The theorem	3
1.2. A geometric dictionary	3
1.3. The exhaustive routing table	5
1.4. The four strategies	6
1.5. Strategy 1: direct length-sum obstructions	7
1.6. Strategy 2: exact boundary–radial demand propagation	8
1.7. Strategy 3: area loss	9
1.8. Strategy 4: the direct nine-point obstruction	10
2. Structural Reduction to Finite Cases	12
2.1. Open, closed, and scaled formulations	13
2.2. The center classification	14
2.3. Vertex types and the T3-like normalization	14
2.4. Reach coordinates, supercritical rows, and gaps	16
2.5. Midpoint forcing	19
3. Trace Budgets and Length Contradictions	21
3.1. Exact neighboring-arm capacity	21
3.2. Conditional skeleton trace caps	23
3.3. Diagonal trace caps	27
3.4. Vd1/Vd2 corner estimates	28

2020 *Mathematics Subject Classification.* 52C15, 52A40, 51M04.

Key words and phrases. equilateral triangles, geometric covering, regular hexagon, boundary propagation, area loss, caliper certificate.

3.5. Exact center intervals and boundary trace caps	29
3.6. Terminal length contradictions	33
4. Strategy 2: Exact Demand Propagation	34
4.1. The local propagation interface	34
4.2. A universal normal form for every selected T_+ arc	35
4.3. Threshold routing	36
4.4. The CE1 one-gap chain with the shortened $X > 1/2$ proof	37
4.5. CE2 one-gap routing	39
4.6. The 407X high-sheet package	40
4.7. Strategy 2 branch assembly	40
5. Area-Loss Obstructions	41
5.1. The local square-loss inequalities	41
5.2. The direct T3-like loss	43
5.3. The cyclic area certificates	45
6. Strategy 4: Direct Nine-Point Forcing	46
6.1. Direct forcing of the disk and three frontier points	46
6.2. Newton inner points	47
6.3. The cap-chain lemma	48
6.4. The four verified overlaps	48
6.5. Enclosure and the center-independent contradiction	49
7. Exhaustive Assembly	49
Appendix A. Exact Local Demand Calculus	50
A.1. The exact local feasible set	50
A.2. The exact capped map	52
A.3. Exact center traces and complementary demands	54
Appendix B. Cyclic Propagation in the All-Vd0 Cases	55
B.1. The exactly-one-supercritical all-Vd0 branch	55
Appendix C. Nonsupercritical, T3-like, and Vd1/Vd2 Demand Branches	66
C.1. The exactly-one-T3 branch	66
C.2. The nonsupercritical CE1/CE2 boundary packages	67
C.3. Two reusable capped-loss inequalities	74
C.4. Half-edge envelopes for adjacent CE2 placements	81
C.5. The exactly-one-Vd1/Vd2 placements	87
C.6. Exact-demand terminal routing	90
Appendix D. The Direct Nine-Point Obstruction	91
D.1. The finite caliper criterion	91
D.2. Corner-clipped AB -unions	92
D.3. Exact-one handoffs and common demands	94
D.4. Six directly forced radial points	94
D.5. Tangent reduction and exact mixed overlaps	99
D.6. The exact global certificate for the mixed overlaps	100
D.7. The enclosure inequality	103
D.8. The center-independent obstruction	105
Appendix E. Table of Symbols	106
Appendix F. Exact Certificate Record	107

1. A GUIDE TO THE PROOF

This section gives the reader the entire proof architecture before any long calculation. It introduces the geometric objects and case labels, states the exhaustive routing table, and previews the four mechanisms that close its rows. Section 2 proves that the labels and reductions are exhaustive. The complete proofs then appear in dependency order in Sections 3–D, before the final audit in Section 7.

1.1. The theorem. Let H be the regular hexagon of side length one.

Theorem 1.1 (Main theorem). *The hexagon H cannot be covered by seven open equilateral triangles of side length one.*

The compactness and scaling argument in Proposition 2.1 gives the equivalent closed form.

Corollary 1.2 (Expanded closed formulation). *For every $L > 1$, the regular hexagon H_L of side length L cannot be covered by seven closed unit equilateral triangles.*

1.2. A geometric dictionary. Indices are taken in $\mathbb{Z}/6\mathbb{Z}$. Put

$$O = (0, 0), \quad V_i = \left(\cos \frac{i\pi}{3}, \sin \frac{i\pi}{3} \right), \quad H = \text{conv}\{V_0, \dots, V_5\}.$$

Write $e_{i,i+1} = [V_i, V_{i+1}]$ for a boundary edge, $r_i = [O, V_i]$ for a radial arm, and $M_i = V_i/2$. The radial target and the full skeleton are

$$D = \bigcup_{i=0}^5 r_i, \quad S = \partial H \cup D,$$

with lengths $\mathcal{H}^1(\partial H) = \mathcal{H}^1(D) = 6$ and $\mathcal{H}^1(S) = 12$.

Seven distinguished points force seven distinct roles: U_C contains O and U_i contains V_i . Their closures are denoted by T_C and T_i . The distinction is essential because closing an open trace may add an endpoint or a tangency.




FIGURE 1. Exact target geometry in the normalization above. The labels show the seven forced role anchors; they do not prescribe the orientations of the covering triangles.

The center type records the number of boundary edges on which T_C has a positive-length trace. These numbers are 0, 1, 2, giving respectively CE0, CE1, CE2.




FIGURE 2. Illustrative examples of the three center-role types. The screenshots are not to scale, and no proof depends on visual inspection.

For a vertex role, let o be the number of its triangle vertices outside H and n the number of adjacent arms met in positive length. The exhaustive names are

type	(o, n)
Vd0	$(1, 0), (2, 0), (3, 0)$
Vd1	$(1, 1)$
Vd2	$(1, 2)$
T3-like	$(2, 1)$.

These classifications are mathematical rather than pictorial; the proofs of exhaustiveness appear in Propositions 2.3 and 2.5.

For an original open vertex role, let A_i, B_i, C_i be its actual maximal incoming boundary, outgoing boundary, and inward radial reaches. Thus row i is *supercritical* when $A_i + B_i > 1$, and

$$N_+ = |\{i : A_i + B_i > 1\}|.$$

Lowercase (a_i, b_i, c_i) will always denote smaller selected demands, never the actual maxima used to define N_+ .




FIGURE 3. Illustrative vertex-role examples. The top row shows three Vd0 configurations; the bottom row shows the remaining three role types. The screenshots are not to scale, and no proof depends on visual inspection.

On $e_{i,i+1}$, parametrized from V_i to V_{i+1} , the two incident open traces are $[0, B_i)$ and $(1 - A_{i+1}, 1]$. Their nonempty complement

$$[B_i, 1 - A_{i+1}]$$

is a boundary gap. Equality gives a singleton gap, which remains uncovered because both incident traces are open at that point.

1.3. The exhaustive routing table. Table 1 is the logical spine of the proof. Each row is a mutually exclusive refinement after the center class and N_+ are fixed, except for the one row explicitly assigned jointly to Strategies 1 and 2. Its exhaustiveness is proved in Section 2, and the row-by-row audit is carried out in Section 7.

center and row count	exhaustive refinement	route
CE0, $N_+ = 0$	all vertex types	1
CE0, $N_+ = 1$	all six roles Vd0	4
CE0, $N_+ = 1$	a Vd1 or Vd2 role	1
CE0, $N_+ = 1$	no Vd1/Vd2 role and a T3-like role	3
CE0, $N_+ \geq 2$	all vertex types	3
CE1/CE2, $N_+ = 0$	all six roles Vd0	2
CE1/CE2, $N_+ = 0$	a Vd1 or Vd2 role	1
CE1/CE2, $N_+ = 0$	no Vd1/Vd2 role and a T3-like role	2
CE1/CE2, $N_+ = 1$	all Vd0, zero gaps	4
CE1/CE2, $N_+ = 1$	all Vd0, exactly one gap	2
CE2, $N_+ = 1$	all Vd0, exactly two gaps	2
CE1, $N_+ = 1$	a Vd1 or Vd2 role	1
CE2, $N_+ = 1$	at least two Vd1/Vd2 roles	1
CE2, $N_+ = 1$	exactly one Vd1/Vd2 role	1+2
CE1/CE2, $N_+ = 1$	no Vd1/Vd2 role and exactly one T3-like role	2
CE1/CE2, $N_+ = 1$	no Vd1/Vd2 role and at least two T3-like roles	1
CE1/CE2, $N_+ \geq 2$	all vertex types	1

TABLE 1. The complete terminal routing. Route j means Strategy j ; $1 + 2$ is the single hybrid package.

1.4. The four strategies.

Strategy 1: one-dimensional trace budgets. Choose one of ∂H , D , or the conditional skeleton S . Bound the trace of each role on that target according to its center class, vertex type, and supercritical status. In every routed branch the seven upper bounds add to strictly less than the target length. The target, rather than the case label, is the organizing idea. Subsection 1.5 gives the reader-level argument, and Section 3 gives the complete trace bounds and terminal contradictions.

Strategy 2: exact demand propagation. A center trace supplies boundary inputs and complementary radial demands. For a nonsupercritical vertex row, a monotone capped map converts an incoming boundary demand into a lower bound on the next row. Iterating around the hexagon returns to row 0 with a demand exceeding its exact remaining capacity. Subsection 1.6 displays the actual-row propagation mechanism. Sections A–C give the piecewise map, the individual five-link inequalities, the rational estimates, and the terminal branches.

Strategy 3: area loss. Normalize a unit triangle to area one, so H has area six. Each selected boundary handoff forces a square amount of a vertex triangle outside the local hexagonal wedge. Two ascents, or one ascent together with a T3-like row, force total loss greater than one; after adding the center triangle, the available inside area is strictly less than six. Subsection 1.7 previews the two cyclic sums, and Section 5 proves their local hypotheses.

Strategy 4: a direct nine-point obstruction. In the zero-gap, all-Vd0, exact-one state, the five nonsupercritical handoffs form one strict descent chain. It produces common lower demands that force six radial points and three asymmetric points into U_C . Their convex hull cannot fit in an open unit equilateral triangle: an exact caliper argument gives enclosing side at least one, whereas compact containment in U_C would give side strictly below one. The reader-level construction is in Subsection 1.8, and the complete argument is in Section D.

The proof uses no global noncoverage claim for S , no interval subdivision, and no branch-and-bound. The last mixed-overlap step is an exact computer-verified

Bernstein certificate; its precise mathematical input, output, and rational reduction are proved in Subsection D.6.

1.5. Strategy 1: direct length-sum obstructions. For a closed triangle T and a rectifiable target E , put $L_E(T) = \mathcal{H}^1(T \cap E)$. If the original open roles cover E , then their closures satisfy

$$(1) \quad \mathcal{H}^1(E) \leq L_E(T_C) + \sum_{i=0}^5 L_E(T_i).$$

Strategy 1 contradicts this inequality on one of three targets. Figure 4 is a visual index; the skeleton estimate is always conditional on the branch hypotheses and is never asserted globally.




FIGURE 4. Exact targets and their one-dimensional measures. The coloring is explanatory. The full-skeleton panel is conditional, not a claim that seven unit triangles can never cover S .

1.5.1. *The human-level mechanism.* There are three ways for the sum in (1) to fail.

- (1) On ∂H , center and vertex classifications give a short trace table. A Vd1/Vd2 role pays a half-unit boundary cap, while a supercritical row forces compensating restrictions on neighboring traces. This closes the perimeter branches.
- (2) On D , a nonsupercritical Vd0 role contributes at most one, a supercritical Vd0 role strictly less than one half, and a T3-like role strictly less than one half. If $k \geq 2$ rows are T3-like and exactly one row is supercritical, the total is

$$L_D(T_C) + \sum_i L_D(T_i) < \frac{3}{2} + \frac{1}{2} + \frac{k}{2} + (5 - k) = 7 - \frac{k}{2} \leq 6,$$

contradicting $\mathcal{H}^1(D) = 6$.

- (3) On S , the center, supercritical, positive-support, and no-support caps combine only under the hypotheses of the conditional skeleton theorem. They close $N_+ \geq 2$ in the CE1/CE2 classes and the indicated hybrid CE2 subcases.

The exact neighboring-arm formula, every trace table, and all endpoint strictness checks are proved in Section 3.

Branches closed by the length argument. Section 3 derives the boundary, radial, diagonal, and conditional-skeleton contradictions and ends with the exact list in Proposition 3.18. It closes every row of Table 1 assigned to route 1. Within the hybrid 1 + 2 row, it closes the Vd2 neighboring-midpoint and additional-positive-support subcases; the complementary exact placements are treated by Strategy 2.

1.6. Strategy 2: exact boundary–radial demand propagation. Length alone does not record where a trace must continue. Strategy 2 keeps that positional information. In local unit vectors

$$u = \left(\frac{1}{2}, \frac{\sqrt{3}}{2} \right), \quad v = \left(\frac{1}{2}, -\frac{\sqrt{3}}{2} \right),$$

define

$$K(a, b, c) = \text{conv}\{0, au, bv, c(u+v)\}.$$

Let $B_c(a)$ be the largest outgoing demand b for which $K(a, b, c)$ fits in a closed unit equilateral triangle, and set

$$F_c(a) = \min\{B_c(a), 1 - a\}, \quad G_c(a) = 1 - F_c(a).$$

The admissible set is coordinatewise down-closed. Hence F_c is nonincreasing, G_c is nondecreasing and extensive, and reflection gives an exact duality. The piecewise formula has a genuine downward jump; no argument below uses continuity at that jump.

1.6.1. How an actual row propagates a demand. Suppose an actual nonsupercritical row has $A_i \geq z$, radial reach $C_i \geq c_i$, and $A_i + B_i \leq 1$. Then its realized demand triple gives

$$B_i \leq F_{c_i}(z).$$

If the next incident open traces cover their boundary point, the next actual incoming reach satisfies

$$(2) \quad A_{i+1} \geq 1 - B_i \geq G_{c_i}(z).$$

Thus each successive use of (2) is an inequality for an actual row, not merely a formal composition of scalar functions.

The nonlinear selected- T_+ part of every capped map is one universal curve. After normalizing the output by the radial deficit, its increment is

$$\psi(x) = 1 - \sqrt{1 - x + x^2}.$$

This gives strict concavity and both chord bounds without a separate implicit differentiation in each branch. Once a chain input crosses a low-root threshold $e(d)$, the map G_{1-d} gives an output at least $1 - e(d)$ and all later extensive maps preserve it. Section 4 proves this normal form and the one-hit and two-threshold routing lemmas.

For a CE1 center trace, exact center geometry supplies one interval $[s, t] \subset e_{0,1}$ and six exits d_i from O ; the complementary radial demand is $c_i = 1 - d_i$. For CE2 there are coupled intervals $[x, u] \subset e_{5,0}$ and $[y, v] \subset e_{0,1}$. Figure 7 in the Strategy 2 technical appendix gives a common schematic for the CE1/CE2, $N_+ = 0$, all-Vd0 package. It shows only three representative Vd0 roles; the argument treats all six.

1.6.2. The branch-level contradictions. For a one-gap branch, repeated use of (2) gives the five actual-row chain $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 5 \rightarrow 0$. The exact terminal estimates have the forms

$$A_0 \geq Z > B_{c_0}(s) \geq A_0$$

for CE1 and

$$A_0 > Z_R > B_{c_0}(y) \geq A_0$$

for the CE2 right gap. Reflection $i \mapsto -i$ gives the left chain $5 \rightarrow 4 \rightarrow 3 \rightarrow 2 \rightarrow 1 \rightarrow 0$ and

$$B_0 > Z_L > B_{c_0}(x) \geq B_0.$$

The shortened CE1 proof uses the universal T_+ chords, a concavity argument forcing $X > 1/2$, and one terminal threshold trigger. The CE2 proof uses the two-threshold routing lemma directly. The two-gap CE2 state uses the exact two-endpoint capped loss; the no-active-gap $N_+ = 0$ state uses the strict overlap of a finite relatively open cover of the connected boundary circle. The remaining branches combine the same interface with the T3-like side tradeoff or the exact Vd1/Vd2 placement classification.

All full radical comparisons, contact-label tables, capped-loss inequalities, and placement polynomials are retained in the Strategy 2 technical appendix, Sections A–C. In particular, the half-edge rational envelope is used only on its valid domain

$$p \geq \frac{1}{2}, \quad 0 \leq h \leq p, \quad p + h < 1;$$

Subsection C.4 records a counterexample outside that domain.

Branches closed by exact demand. Proposition 4.5 closes every row of Table 1 assigned to route 2, including the complementary exact placements in the hybrid 1 + 2 row. The list consists of the all-Vd0 and T3-like $N_+ = 0$ packages, every positive-gap all-Vd0 state with $N_+ = 1$, the exactly-one-T3-like state, and the complementary CE2 placements. Singleton gaps are included. Equation (2) is the handoff from each scalar inequality to the actual role geometry.


1.7. Strategy 3: area loss. Normalize the area of a unit equilateral triangle to one. Then $\text{area}(H) = 6$. At a vertex, affine wedge coordinates identify $T_i \cap H$ with the portion of T_i inside the first quadrant. For lower boundary reaches a, b , the local square-loss theorem gives

$$\text{loss}(T_i) \geq \min(a, b)^2, \quad a + b > 1 \implies \text{loss}(T_i) \geq \max(a, b)^2.$$

A T3-like row has the separate bound

$$a, b \geq m \implies \text{loss}(T_i) \geq 2m - 4m^2.$$

Their coordinate proofs, with every orientation retained, are in Section 5.



(a) local retained area (b) global retained areas
 FIGURE 5. Author-supplied schematic views of Strategy 3, not to scale. The local panel shows one vertex-role triangle with boundary reaches a, b , and the global panel shows the six cyclic vertex-role triangles. Only the darker colored portions represent area retained inside H . The exact loss inequalities and cyclic sums are proved independently in Section 5.

1.7.1. *The two global sums.* Strict handoffs have $(a_i, b_i) = (1 - x_{i-1}, x_i)$, and selected supercritical rows are precisely the cyclic ascents $x_i > x_{i-1}$. If there are at least two ascents, let $m = \min_i x_i$ and reflect so that $m \leq 1 - \max_i x_i$. Four ordinary rows lose at least m^2 , a row leaving the minimum loses at least $(1 - m)^2$, and a second ascent loses strictly more than $1/4$. Hence the total loss is greater than

$$4m^2 + (1 - m)^2 + \frac{1}{4} = 5 \left(m - \frac{1}{5} \right)^2 + \frac{21}{20} > 1.$$

If there is one ascent and a T3-like row, the supercritical row loses at least $(1 - m)^2$, the T3-like row at least $2m - 4m^2$, and the other four rows at least m^2 each. Their total is

$$(1 - m)^2 + (2m - 4m^2) + 4m^2 = 1 + m^2 > 1.$$

The six vertex roles therefore contribute total inside area less than five; adding the center role still gives less than the required six.

Branches closed by area loss. Strategy 3 closes precisely the two rows of Table 1 assigned to route 3: CE0 with $N_+ \geq 2$, and CE0 with $N_+ = 1$, no Vd1/Vd2 role, and at least one T3-like role. The strict-handoff theorem transfers the actual N_+ pattern to the selected ascents, and the two sums above give the global certificate. Section 5 verifies the local hypotheses and the strict normalization, culminating in Proposition 5.6.

1.8. **Strategy 4: the direct nine-point obstruction.** Assume all six vertex roles are Vd0, they cover ∂H , and exactly one actual boundary row is supercritical.

Rotate the supercritical index to 4. The strict handoffs from Proposition 2.9 obey

$$(3) \quad x_3 < x_2 < x_1 < x_0 < x_5 < x_4.$$

Put

$$a = 1 - x_3, \quad b = x_4, \quad p = 1 - b, \quad q = 1 - a.$$

Then $a + b > 1$, $a^2 + ab + b^2 < 1$, and every selected row has incoming and outgoing demands at least p and q . This single descent chain is why the strategy is specific to $N_+ = 1$.

1.8.1. *Forcing nine points into the center role.* Let $c_* = c_{\max}(p, q)$ be the exact maximal radial reach for the common boundary demands. The six points

$$D_i = (1 - c_*)V_i \quad (0 \leq i \leq 5)$$

are excluded from their matching Vd0 role by maximality and openness, from the adjacent roles by Vd0 locality, and from the remaining roles by the diameter-one condition. Thus all six lie in U_C , and their convex hull forces a centered disk $\mathcal{D}_\eta \subset U_C$.

Three parameter-dependent points Q_- , Q_0 , Q_+ are then placed on the two actual neighboring handoff lines. The unique supercritical row is excluded by its exact strict frontier; rows 0, 1, 2 are excluded by direct vertex distances; and rows 3, 5 are excluded by the actual handoffs. Hence these three points also lie in U_C .




FIGURE 6. Author-supplied illustrative Strategy 4 core-case configuration, not to scale. The displayed placement is an example only; no exclusion, forcing, or enclosure inequality is inferred from the picture.

1.8.2. *Why the witness cannot fit.* For a compact set K , let $\Lambda(K)$ be the least side length of a closed equilateral triangle containing K . The exact outer frontier points contain two independent quadratic radicals, so the proof replaces Q_- and Q_+ by one-step Newton points lying strictly inside the segments from Q_0 to those outer points. This produces an inner witness set with rational coordinates in the single radical $D = \sqrt{4(a^2 + ab + b^2) - 3}$.

The geometric core is then a four-cap chain in the support-direction circle. The concise proof in Section 6 consists of direct forcing, the Newton reduction, a cyclic cap-chain lemma, a proposition asserting the four overlaps, and the enclosure contradiction. The two adjacent overlaps are analytic. The two mixed overlaps are reduced to exact integer-polynomial signs and verified by global Bernstein identities in the Strategy 4 technical appendix, Subsection D.6. Thus

$$\Lambda(\text{conv}(\mathcal{D}_\eta \cup \{Q_-, Q_0, Q_+\})) \geq 1.$$

The nine-point set is compact. If it lay inside the open unit triangle U_C , positive clearance from the three sides would place it in a closed equilateral triangle of side strictly below one, contradicting the displayed lower bound.

Branches closed by direct forcing. Strategy 4 closes exactly the zero-gap, all-Vd0, $N_+ = 1$ rows of Table 1, for every center class. The reader-facing center-independent obstruction is Theorem 6.4. In CE0, a missed boundary point would create a positive center trace, so zero gaps are automatic. In CE1 or CE2, zero gaps say directly that the six vertex roles cover ∂H . Proposition 6.5 makes these two applications explicit.

2. STRUCTURAL REDUCTION TO FINITE CASES

Throughout, indices are taken in $\mathbb{Z}/6\mathbb{Z}$. Put

$$O = (0, 0), \quad V_i = \left(\cos \frac{i\pi}{3}, \sin \frac{i\pi}{3} \right), \quad H = \text{conv}\{V_0, \dots, V_5\}.$$

Thus H is the regular hexagon of side length one. We use the notation

$$e_{i,i+1} = [V_i, V_{i+1}], \quad r_i = [O, V_i], \quad M_i = \frac{1}{2}V_i,$$

and, when needed,

$$D = \bigcup_{i=0}^5 r_i, \quad S = \partial H \cup D, \quad S_{1/2} = \partial H \cup \{O, M_0, \dots, M_5\}.$$

The arms r_i meet only at O , up to sets of one-dimensional measure zero, and consequently

$$(4) \quad \mathcal{H}^1(\partial H) = 6, \quad \mathcal{H}^1(D) = 6, \quad \mathcal{H}^1(S) = 12.$$

For $L > 0$, $H_L := LH$ denotes the concentric regular hexagon of side length L . All symmetries used below belong to the dihedral group D_6 ; applying a symmetry always transports the indices, orientations, and incoming and outgoing labels together.

2.1. Open, closed, and scaled formulations.

Proposition 2.1 (Compactness and scaling equivalence). *The following assertions are equivalent.*

- (1) *The set H is covered by seven open equilateral triangles of side length 1.*
- (2) *For some $\varepsilon > 0$, the set H is covered by seven closed equilateral triangles of side length $1 - \varepsilon$.*
- (3) *For some $L > 1$, the set H_L is covered by seven closed equilateral triangles of side length 1.*

Proof. Suppose that open unit triangles $U_1, \dots, U_7$ cover H . Define

$$m_j(p) = \text{dist}(p, \mathbb{R}^2 \setminus U_j), \quad m(p) = \max_{1 \leq j \leq 7} m_j(p).$$

The continuous function m is positive on the compact set H , so $\eta := \min_{p \in H} m(p) > 0$. Choose

$$0 < \rho < \min \left\{ \eta, \frac{\sqrt{3}}{6} \right\}.$$

Moving each side of U_j inward through distance ρ gives a closed equilateral triangle K_j of side $1 - 2\sqrt{3}\rho$. Every $p \in H$ has $m_j(p) \geq \eta > \rho$ for some j , and hence belongs to K_j . This proves (1) $\Rightarrow$ (2), with $\varepsilon = 2\sqrt{3}\rho$.

Conversely, enlarging a closed triangle of side $1 - \varepsilon$ about its incenter to an open triangle of side 1 preserves containment, proving (2) $\Rightarrow$ (1). Finally, dilation by $(1 - \varepsilon)^{-1}$ identifies (2) with (3), with

$$L = (1 - \varepsilon)^{-1} > 1.$$

□

We conduct the geometric argument in the open side-one model. Denote the original open triangles by U_C and U_i for $0 \leq i \leq 5$, and put

$$T_C = \overline{U_C}, \quad T_i = \overline{U_i}.$$

The distinction between U_i and T_i will be maintained: closure may add endpoints or tangencies, although it does not change area or one-dimensional trace measure.

Lemma 2.2 (Distinct roles). *The seven triangles can be labelled so that*

$$O \in U_C, \quad V_i \in U_i \quad (0 \leq i \leq 5),$$

and these seven roles are distinct. Consequently

$$O \in \text{int}(T_C), \quad V_i \in \text{int}(T_i).$$

Proof. The seven distinguished points $O, V_0, \dots, V_5$ are pairwise at distance at least 1. Any two points of an open unit equilateral triangle are at distance strictly less than its diameter 1. Thus one open triangle contains at most one distinguished point. Selecting a covering triangle for each of the seven points gives an injection from seven points to seven triangles, hence a bijection and the asserted labelling. □

2.2. The center classification. For a closed center role define

$$N_E(T_C) = |\{i : T_C \cap e_{i,i+1} \text{ contains a positive-length interval}\}|.$$

Point contacts, endpoint contacts, and tangencies do not contribute. We say that T_C has type CE0, CE1, or CE2 according as $N_E(T_C)$ is 0, 1, or 2.

Proposition 2.3 (Exhaustive center types). *Every center role has exactly one of the types CE0, CE1, and CE2. Equivalently,*

$$N_E(T_C) \leq 2.$$

Proof. Among any three edges of the six-cycle ∂H , two have no common endpoint. It is therefore enough to show that relative-interior points of two nonadjacent edges are more than unit distance apart. Up to symmetry there are two cases. For $0 < t, s < 1$, put

$$x = (1 - t)V_0 + tV_1.$$

If $y = (1 - s)V_2 + sV_3$, direct calculation gives

$$\|x - y\|^2 - 1 = (1 - t)(2 - t) + s(s + t) > 0.$$

If instead $y = (1 - s)V_3 + sV_4$ and $u = t + s$, then

$$\|x - y\|^2 - 1 = (u - 1)^2 + 2 > 0.$$

A unit equilateral triangle has diameter 1. It therefore cannot contain relative-interior points on two nonadjacent hexagon edges. Three positive-length edge traces would supply such a pair, a contradiction. $\square$

2.3. Vertex types and the T3-like normalization. For an original vertex role T_i , let

$$o(T_i) = |\{\text{vertices of } T_i \text{ outside } H\}|$$

and let $n(T_i)$ be the number of adjacent arms r_{i-1}, r_{i+1} having positive-length intersection with T_i . We use the names

type	(o, n)
Vd0	$(1, 0), (2, 0), (3, 0)$
Vd1	$(1, 1)$
Vd2	$(1, 2)$
T3-like	$(2, 1)$.

Lemma 2.4 (Local wedge). *Every original vertex role has $1 \leq o \leq 3$. If it has positive-length intersection with one adjacent arm, at least one of its vertices belongs to H ; if it has positive-length intersection with both adjacent arms, at least two of its vertices belong to H .*

Proof. Normalize to $i = 0$ and write

$$X = V_0 + x(V_1 - V_0) + y(V_5 - V_0).$$

Then $V_0 = (0, 0)$,

$$\|(x, y)\|^2 = x^2 + y^2 - xy,$$

and, within the closed unit ball about V_0 , membership in H is equivalent to membership in the wedge

$$W = \{(x, y) : x \geq 0, y \geq 0\}.$$

Indeed,

$$x^2 + y^2 - xy = \left(y - \frac{x}{2}\right)^2 + \frac{3x^2}{4} = (x - y)^2 + xy$$

shows that $x, y \geq 0$ and norm at most one imply the remaining hexagon inequalities. The adjacent arms are

$$r_1 = \{(1, y) : 0 \leq y \leq 1\}, \quad r_5 = \{(x, 1) : 0 \leq x \leq 1\}.$$

Suppose $T_0 \cap r_1$ has positive length, and let m be the largest x -coordinate of a vertex of T_0 . If $m > 1$, a maximizing vertex (m, y) lies in the unit ball and hence satisfies $0 < y < 1$; it belongs to H . If $m = 1$, positive-length contact with the supporting line $x = 1$ forces a side of T_0 to lie there, and both endpoints satisfy $1 + y^2 - y \leq 1$, hence belong to H . The same argument applies to r_5 . If both arms are met but only one triangle vertex belonged to H , that vertex would have to have both $x > 1$ and $y > 1$. This is impossible because $(x - y)^2 + xy > 1$.

Finally, if all three triangle vertices belonged to the convex set H , then $T_0 \subset H$. This is incompatible with the extreme point V_0 lying in the interior of T_0 . Hence $o \geq 1$. $\square$

Proposition 2.5 (Exhaustive vertex types). *Every original vertex role has exactly one of the four types Vd0, Vd1, Vd2, and T3-like.*

Proof. We have $n \in \{0, 1, 2\}$. Lemma 2.4 gives $o \leq 2$ when $n \geq 1$, and $o \leq 1$ when $n = 2$. Together with $o \geq 1$, the possibilities are

n	possible o
0	1, 2, 3
1	1, 2
2	1,

which are precisely the four named classes. $\square$

The next normalization is useful for trace estimates, but its final warning is essential.

Proposition 2.6 (T3-like closed-trace domination). *Let T be the closure of an original T3-like V_i -role. There is a translate $\hat{T}$ of T , with the same orientation and side length, such that*

$$V_i \in \text{relint}(a \text{ side of } \hat{T}), \quad T \cap H \subsetneq \hat{T} \cap H,$$

and $\hat{T}$ remains of type $(o, n) = (2, 1)$. Every old open interior point in $H \setminus \{V_i\}$ remains an interior point of $\hat{T}$. The point V_i itself becomes a boundary point, so $\hat{T}$ is a closed-trace majorant and is not a replacement open role.

Proof. Use the wedge coordinates of Lemma 2.4. Set

$$D_t = 1 - t + t^2, \quad z = \sqrt{D_t}, \quad 0 \leq t \leq 1.$$

After relabelling triangle vertices, every orientation belongs to one of the following two families:

	first side vector	second side vector
I	$z^{-1}(1, t)$	$z^{-1}(1 - t, 1)$
II	$z^{-1}(t, -(1 - t))$	$z^{-1}(1, t)$.

For Type I write

$$U = \alpha + y - tx, \quad V = \beta + x - (1 - t)y, \quad T = \{U \geq 0, V \geq 0, U + V \leq z\}.$$

Here $\alpha, \beta > 0$ and $\alpha + \beta < z$. Direct inversion shows that if exactly two triangle vertices are outside W , the sole wedge vertex has both coordinates strictly below 1: in the two possible cases its coordinates are bounded respectively by $(1-t)/z$ and t/z , with the endpoint strictness supplied by $\alpha > 0$ or $\beta > 0$. The support argument in Lemma 2.4 then excludes positive-length contact with either $x = 1$ or $y = 1$. Thus Type I cannot be T3-like.

For Type II write

$$U = \alpha + (1-t)x + ty, \quad V = \beta + tx - y, \quad T = \{U \geq 0, V \geq 0, U + V \leq z\}.$$

Reflecting in $x = y$ if necessary, assume the supported adjacent arm is r_1 . The two outside vertices have respectively a negative x - and a negative y -coordinate, while the unique wedge vertex is

$$Q = \left(\frac{z - \alpha - t\beta}{D_t}, \frac{t(z - \alpha) + (1-t)\beta}{D_t} \right).$$

The endpoint cases $t = 0, 1$ give no positive adjacent interval, so $0 < t < 1$. The exact axis reaches are

$$A = \beta, \quad B = z - \alpha - \beta,$$

and T3-like support is equivalent to

$$Q_x > 1, \quad Q_y \leq 1.$$

Define $\hat{T}$ by replacing α with 0 and keeping t, β fixed. Since the linear map $(x, y) \mapsto (U, V)$ is nonsingular, this changes only the translation. At the origin, $\hat{U} = 0$ and $0 < \hat{V} = \beta < z$, so V_0 is in the relative interior of a side. For $(x, y) \in W$,

$$\hat{U} = (1-t)x + ty \geq 0, \quad \hat{V} = V, \quad \hat{U} + \hat{V} = U + V - \alpha.$$

Thus $T \cap W \subset \hat{T} \cap W$. The old x -axis reach is B and the new one is $B + \alpha$; because $B < 1$ and $\alpha > 0$, the inclusion is strict inside H .

The two outside vertices remain outside, whereas V_0 and the new wedge vertex lie in H , so $o = 2$. Moreover $\hat{Q}_x > Q_x > 1$. From $Q_x > 1$ one gets $t\beta < z - \alpha - D_t$, and hence

$$tD_t(1 - \hat{Q}_y) > D_t(1 - z) + (1-t)\alpha > 0.$$

Thus $\hat{Q}_y < 1$ and $n = 1$. Finally, for every nonzero point of W , $\hat{U} > 0$; the displayed slack relations show that every old interior point other than the origin remains interior. $\square$

2.4. Reach coordinates, supercritical rows, and gaps. For the original vertex role U_i , define its actual maximal incoming, outgoing, and radial reaches by

$$\begin{aligned} A_i &= \sup\{t \in [0, 1] : V_i + t(V_{i-1} - V_i) \in U_i\}, \\ B_i &= \sup\{t \in [0, 1] : V_i + t(V_{i+1} - V_i) \in U_i\}, \\ C_i &= \sup\{t \in [0, 1] : V_i + t(O - V_i) \in U_i\}. \end{aligned}$$

Thus the closure T_i contains the corresponding endpoints. We reserve lowercase (a_i, b_i, c_i) for selected or prescribed lower-bound demands whenever actual and selected values occur together. A realizing role then satisfies

$$A_i \geq a_i, \quad B_i \geq b_i, \quad C_i \geq c_i.$$

An actual row is supercritical if $A_i + B_i > 1$, and

$$(5) \quad N_+ = |\{i : A_i + B_i > 1\}|.$$

This definition never counts a smaller demand pair in place of the actual row.

Parametrize $e_{i,i+1}$ by

$$X_i(x) = V_i + x(V_{i+1} - V_i), \quad 0 \leq x \leq 1.$$

The two incident open traces are

$$U_i \cap e_{i,i+1} = [0, B_i), \quad U_{i+1} \cap e_{i,i+1} = (1 - A_{i+1}, 1]$$

in this coordinate. A *boundary gap* on this edge is the nonempty closed complement between these two traces,

$$[B_i, 1 - A_{i+1}] \quad \text{when } B_i \leq 1 - A_{i+1}.$$

In particular, equality gives a singleton gap. This convention records the actual open traces; a point missing from both open roles is not erased merely because both closures contain it.

Lemma 2.7 (Exhaustiveness of the gap split). *On each edge the complement of the two incident vertex-role traces is either empty or one closed interval; a singleton interval counts as a gap. Every edge having a gap has positive-length intersection with the open center role. Consequently a CE0 center permits no gaps, a CE1 center permits at most one, and a CE2 center permits at most two.*

Proof. The two incident traces are intervals issuing from the two endpoints, so their complement is exactly the interval displayed above when nonempty. A nonincident vertex role cannot contain a relative-interior point of the edge: that point is more than unit distance from the role's distinguished vertex. Thus every point of a gap lies in U_C . Because U_C is open, its intersection with that edge contains a relative neighborhood and hence has positive length. Different gap edges therefore give different edges counted by $N_E(T_C)$. Proposition 2.3 gives the three stated bounds, including singleton gaps. $\square$

Lemma 2.8 (T3-like rows are nonsupercritical). *Every original T3-like row satisfies*

$$(6) \quad A_i + B_i \leq 1.$$

In particular, it is not counted by N_+ .

Proof. The proof of Proposition 2.6 showed that a T3-like role has Type II orientation with

$$A_i = \beta, \quad B_i = z - \alpha - \beta, \quad z = \sqrt{1 - t + t^2} \leq 1.$$

Thus $A_i + B_i = z - \alpha \leq 1$. The same conclusion holds after reflection. $\square$

We record the precise transfer from actual rows to strict handoff demands. Its hypothesis that the six vertex roles themselves cover the boundary is part of the statement.

Proposition 2.9 (Strict boundary handoffs). *Assume that $U_0, \dots, U_5$ cover ∂H . On each edge choose*

$$x_i \in (1 - A_{i+1}, B_i)$$

and define

$$a_i = 1 - x_{i-1}, \quad b_i = x_i.$$

Then $0 < a_i < A_i$, $0 < b_i < B_i$, the two selected anchors lie in U_i , and

$$a_i^2 + a_i b_i + b_i^2 < 1.$$

Moreover:

- (1) if exactly one actual row, indexed by p , is supercritical, every strict selection has exactly one selected supercritical row, also indexed by p ;
- (2) if at least two actual rows are supercritical, some simultaneous strict selection has at least two selected supercritical rows.

Proof. Because V_i is interior to a diameter-one closed triangle,

$$0 < A_i < 1, \quad 0 < B_i < 1.$$

No nonincident vertex role contains a relative-interior point of $e_{i,i+1}$, since such a point is more than unit distance from its distinguished vertex. The two displayed incident traces therefore cover the edge and, being relatively open and nonempty, overlap. Hence each interval $(1 - A_{i+1}, B_i)$ is nonempty. The anchor and distance assertions follow from the choice and from the fact that two points of U_i are at distance strictly less than one.

Put $\ell_i = 1 - A_{i+1}$ and $u_i = B_i$. Then

$$(7) \quad A_i + B_i > 1 \iff \ell_{i-1} < u_i, \quad a_i + b_i > 1 \iff x_{i-1} < x_i.$$

If actual row i is nonsupercritical, then

$$x_i < u_i \leq \ell_{i-1} < x_{i-1},$$

so every selected version of it is strictly subcritical. Also

$$(8) \quad \sum_{i=0}^5 (a_i + b_i) = \sum_{i=0}^5 (1 - x_{i-1} + x_i) = 6.$$

If p is the sole actual supercritical index, the other five selected sums are strictly below one; equation (8) forces the p th sum above one. This proves (1).

For (2), first take two nonadjacent supercritical indices p, q . For each $r \in \{p, q\}$ choose

$$\ell_{r-1} < z_r < u_r,$$

then choose

$$x_{r-1} \in (\ell_{r-1}, \min\{u_{r-1}, z_r\}), \quad x_r \in (\max\{\ell_r, z_r\}, u_r).$$

The two pairs of variables are disjoint and give ascents at p, q . Any set of at least three indices on a six-cycle contains two nonadjacent indices. It remains to consider exactly two adjacent supercritical indices $p, p+1$. For the other four nonsupercritical rows, successive inequalities $\ell_i < u_i \leq \ell_{i-1}$ give

$$\ell_{p-1} < \ell_{p+1}.$$

Together with supercriticality at $p, p+1$, this implies

$$\max\{\ell_{p-1}, \ell_p\} < \min\{u_p, u_{p+1}\}.$$

Choose x_p between these bounds and then choose $x_{p-1} < x_p < x_{p+1}$ in their respective overlap intervals. By (7), both adjacent rows are selected supercritical. $\square$

2.5. Midpoint forcing. We first need a local fact for vertex roles. Take unit vectors

$$u = \left(\frac{1}{2}, \frac{\sqrt{3}}{2} \right), \quad v = \left(\frac{1}{2}, -\frac{\sqrt{3}}{2} \right).$$

Lemma 2.10 (Self-midpoint obstruction). *Let a closed unit triangle contain*

$$0, \quad au, \quad bv, \quad \frac{u+v}{2}.$$

Then $a + b \leq 1$. Consequently, for every vertex role,

$$M_i \in T_i \implies A_i + B_i \leq 1.$$

If the two demanded boundary points and the midpoint are contained with a positive margin, the conclusion is strict.

Proof. Suppose $a + b > 1$ and set

$$K = \text{conv} \left\{ 0, au, \frac{u+v}{2}, bv \right\}.$$

For outward unit normals n_0, n_1, n_2 separated by 120° , the least equilateral triangle with those normals and containing K has side length

$$\frac{2}{\sqrt{3}} \sum_{j=0}^2 h_K(n_j).$$

On an angular interval where the three support vertices are fixed, the support sum S satisfies $S'' = -S < 0$. Hence its angular minimum occurs at a breakpoint, namely at an outward normal to one of the four edges of K .

For the edge from 0 to au , direct projection gives

$$S_{0A} = \max \left\{ \frac{\sqrt{3}}{4}, \frac{\sqrt{3}a}{2} \right\} + \frac{\sqrt{3}b}{2} > \frac{\sqrt{3}}{2};$$

if $a \geq 1/2$ this is $(\sqrt{3}/2)(a+b)$, and if $a < 1/2$ then $b > 1/2$. The edge through bv is symmetric. For the edge from au to $(u+v)/2$, put $D_a = \sqrt{4a^2 - 2a + 1}$. Projection gives

$$S_{AM} = \begin{cases} \frac{\sqrt{3}(a+b)}{2D_a}, & a \leq \frac{1}{2}, \\ \frac{\sqrt{3}(2a^2+b)}{2D_a}, & a > \frac{1}{2}. \end{cases}$$

The first value is strictly above $\sqrt{3}/2$ because $D_a \leq 1$. In the second case, $b > 1 - a$ and

$$(2a^2 - a + 1)^2 - D_a^2 = a^2(2a - 1)^2 \geq 0,$$

so again $S_{AM} > \sqrt{3}/2$. The fourth edge is symmetric. Every enclosing equilateral triangle therefore has side greater than one, a contradiction.

If equality $a + b = 1$ occurred while all relevant points had positive margin, both boundary demands could be increased slightly; the closed conclusion would then be violated. $\square$

The corresponding center fact is exact and uses the interior condition at O .

Proposition 2.11 (Unique center midpoint). *Let T_C be a closed unit equilateral triangle with $O \in \text{int}(T_C)$. If T_C has positive-length intersection with a boundary edge of H , then it contains exactly one of $M_0, \dots, M_5$.*

Proof. Normalize a positive overlap to

$$T_C \cap e_{0,1} = [s, t], \quad 0 \leq s < t \leq 1,$$

where $e_{0,1}$ is parametrized from V_0 to V_1 . In affine coordinates

$$X = V_0 + b(V_1 - V_0) + a(V_5 - V_0),$$

reflect if necessary so that the ratio of the two cutting slopes satisfies $0 < \lambda < 1$, and put $\rho = \sqrt{1 - \lambda + \lambda^2}$. The triangle has the exact side-slack representation

$$\begin{aligned} F_1 &= \lambda b + (1 - \lambda)a - \lambda s, \\ F_2 &= -b + \lambda a + t, \\ F_0 &= (1 - \lambda)b - a + \rho + \lambda s - t, \end{aligned} \quad T_C = \{F_0, F_1, F_2 \geq 0\},$$

with $F_0 + F_1 + F_2 = \rho$. The case $\lambda = 1$ would give $F_0(O) = s - t < 0$; the reflected labelling supplies $0 < \lambda < 1$.

Set $C_j = F_j(O)$. Since O is interior,

$$C_0, C_1, C_2 > 0, \quad C_0 + C_1 + C_2 = \rho.$$

The positive overlap is equivalent to

$$C_0 + (1 - \lambda)C_2 < P, \quad P = \rho(1 - \rho).$$

Using $1 - \rho^2 = \lambda(1 - \lambda)$, we have

$$P = \frac{\lambda(1 - \lambda)\rho}{1 + \rho} < \frac{\lambda(1 - \lambda)}{2}, \quad P < \frac{1 - \lambda}{2}.$$

Substitution of the six midpoints gives the exact criteria

$$\begin{array}{l|l} M_0 & C_1 \geq \frac{1}{2} \\ M_1 & C_1 \geq \frac{1-\lambda}{2}, \quad C_2 \geq \frac{\lambda}{2} \\ M_2 & C_2 \geq \frac{1}{2} \\ M_3 & C_0 \geq \frac{\lambda}{2}, \quad C_2 \geq \frac{1-\lambda}{2} \\ M_4 & C_0 \geq \frac{1}{2} \\ M_5 & C_0 \geq \frac{1-\lambda}{2}, \quad C_1 \geq \frac{\lambda}{2}. \end{array}$$

Now

$$C_0 + C_2 < \frac{P}{1 - \lambda},$$

and therefore

$$C_1 > \rho - \frac{P}{1 - \lambda} = \frac{\rho(\rho - \lambda)}{1 - \lambda} > \frac{1}{2};$$

the last inequality follows from

$$2\rho(\rho - \lambda) - (1 - \lambda) = \frac{(1 - \lambda)(\rho - \lambda)}{\rho + \lambda} > 0.$$

Thus $M_0 \in T_C$. On the other hand,

$$C_2 < \frac{P}{1 - \lambda} < \frac{\lambda}{2}, \quad C_0 < P < \min \left\{ \frac{\lambda}{2}, \frac{1 - \lambda}{2}, \frac{1}{2} \right\}.$$

The table excludes $M_1, \dots, M_5$. Undoing the reflection and rotation proves the assertion. $\square$

Remark 2.12. The hypothesis $O \in \text{int}(T_C)$ cannot be weakened to $O \in T_C$: the triangle $\text{conv}\{O, V_0, V_1\}$ has a boundary-edge overlap and contains both M_0 and M_1 .

Proposition 2.13 (Exhaustive structural reduction). *Every hypothetical cover by seven open unit equilateral triangles admits the distinct center and vertex roles used in Table 1, and its actual reaches determine exactly one row of that table.*

Proof. Lemma 2.2 gives the seven roles. Propositions 2.3 and 2.5 give the exhaustive and mutually exclusive center and vertex types. The actual reaches determine exactly one of $N_+ = 0$, $N_+ = 1$, and $N_+ \geq 2$. In the all-Vd0, $N_+ = 1$ center-trace cases, Lemma 2.7 supplies the exhaustive gap refinement. The remaining vertex-type refinements in Table 1 are disjoint by construction. $\square$

3. TRACE BUDGETS AND LENGTH CONTRADICTIONS

This section develops the complete trace formulas and length contradictions. It includes all endpoint and strictness checks used by the length mechanism.

For a closed triangle T and a rectifiable target E , write

$$L_E(T) = \mathcal{H}^1(T \cap E).$$

The passage from an original open role U to its closure $T = \overline{U}$ can only enlarge the trace. Hence a cover of E by the open roles implies

$$(9) \quad \mathcal{H}^1(E) \leq L_E(T_C) + \sum_{i=0}^5 L_E(T_i).$$

All estimates in this section concern full traces. They therefore remain valid for any assigned measurable subsets of those traces.

The strategy uses three one-dimensional targets: ∂H , D , and $S = \partial H \cup D$, whose measures were recorded in (4). The skeleton estimates below are conditional on specific center and row hypotheses; no unconditional noncoverage assertion for S is made.

3.1. Exact neighboring-arm capacity. We first record the exact neighboring-arm capacity. Normalize at V_0 and put

$$A_a = V_0 + a(V_5 - V_0), \quad B_b = V_0 + b(V_1 - V_0), \quad X_c = (1 - c)V_1.$$

Let $C_+(a, b)$ be the largest $c \in [0, 1]$ for which a closed unit equilateral triangle contains V_0, A_a, B_b, X_c ; call it undefined when no such c exists. Reflection gives $C_-(a, b) = C_+(b, a)$ for the other neighboring arm.

Theorem 3.1 (Exact neighboring-arm capacity). *For $0 \leq a \leq 1/2$, let $p(a)$ be the unique zero in $[a, 1]$ of*

$$f_a(x) = x^3 - (a + 2)x^2 + 2(a + 1)x - 1,$$

and set

$$\sigma(a) = 1 - p(a), \quad \tau(a) = 1 - a - (p(a) - a)(1 - p(a)).$$

Then

$$C_+(a, b) = \begin{cases} \text{undefined}, & a + b > 1, \\ 1 - b, & a + b \leq 1, \ a > 1/2, \\ 1 - b, & 0 \leq a \leq 1/2, \ 0 \leq b \leq \sigma(a), \\ p(a), & 0 \leq a \leq 1/2, \ \sigma(a) \leq b \leq \tau(a), \\ a + \frac{1}{2} - \sqrt{(a+b)^2 - \frac{3}{4}}, & 0 \leq a \leq 1/2, \ \tau(a) < b \leq 1 - a. \end{cases}$$

At $b = \tau(a)$ the plateau value $p(a)$ is used. In particular, if a vertex role has actual reaches $A_i + B_i > 1$, it has no positive-length trace on either adjacent radial arm; in fact it contains no point of either adjacent arm.

Proof. Use the cone basis

$$E_- = V_5 - V_0, \quad E_+ = V_1 - V_0.$$

Then the four points have coordinates

$$(0, 0), \quad (a, 0), \quad (0, b), \quad (c, 1),$$

and $\|uE_- + vE_+\|^2 = u^2 + v^2 - uv$. For $R^2 + S^2 - RS = 1$, define

$$U = (R - S)u + Sv, \quad V = Su - Rv, \quad W = Ru + (S - R)v.$$

A translate of the unit equilateral triangle with this orientation contains the four points exactly when

$$(*) \quad \max W - \min U - \min V \leq 1.$$

As in the support argument in Lemma 2.10, on every angular cell with fixed support vertices the left side of $(*)$ is a sinusoid with negative second derivative at an interior minimum candidate. Thus a minimizing orientation lies at a support breakpoint, where a side contains one of the hull edges. Checking OA, AX, XB, BO against the three possible support sides leaves, up to dominated reflected patterns, precisely the following three configurations:

active support	candidate c
OA and BX	$1 - b$
AX, B inactive	$p(a)$
AX, B active	$a + \frac{1}{2} - \sqrt{(a+b)^2 - \frac{3}{4}}$

We give the algebra selecting the cells, which also verifies that no squared branch has been retained incorrectly.

For the first pattern one may take $R = S = 1$. Its side length is $b + c$, and A is contained exactly when $a \leq c$. Thus $c = 1 - b$ is feasible exactly when $a + b \leq 1$.

For the second and third patterns, the side AX gives, with $d = c - a$, $R = Sd$. When B is inactive, the opposite support conditions give

$$(d(c - 1) + 1)^2 = d^2 - d + 1,$$

which factors as

$$(c - a)f_a(c) = 0.$$

For $0 \leq a \leq 1/2$, one has $f_a(a) = 2a - 1 \leq 0$, $f_a(1) = a \geq 0$, and

$$f'_a(x) = 3x^2 - 2(a + 2)x + 2(a + 1) > 0$$

because its discriminant is $4(a^2 - 2a - 2) < 0$. Hence the required root $p(a)$ is unique. The remaining containment inequality is

$$b \leq 1 - p(a)(1 + a - p(a)) = \tau(a),$$

and this branch dominates $1 - b$ exactly when $b \geq \sigma(a)$.

When B is active, the side-length equation gives $S = -1/(a + b)$ and

$$d^2 - d + 1 = (a + b)^2.$$

The relevant root $0 \leq d \leq 1/2$ yields the last expression in (3.1). Its nonactive inequalities reduce to

$$0 \leq a \leq \frac{1}{2}, \quad \tau(a) \leq b \leq 1 - a.$$

The square root is real on this cell. Indeed, writing $p = p(a)$ gives

$$a = \frac{(1-p)(p^2 - p + 1)}{p(2-p)}, \quad a + \tau(a) = \frac{p^2 - p + 1}{p(2-p)} \geq \frac{\sqrt{3}}{2}.$$

Finally, for $s = a + b \in [\sqrt{3}/2, 1]$,

$$s - \frac{1}{2} - \sqrt{s^2 - \frac{3}{4}} \geq 0,$$

so this branch dominates $1 - b$. The three selected cells are ordered because $\tau(a) - \sigma(a) = (p(a) - a)p(a) \geq 0$ and $\tau(a) \leq 1 - a$. The support-edge enumeration shows that when $a + b > 1$ none is feasible, proving undefinedness. Reflection exchanges a, b and the two neighboring arms. $\square$

3.2. Conditional skeleton trace caps. We now develop the skeleton estimate used only when the center is CE1 or CE2. The first lemma is the center contribution.

Lemma 3.2 (Center skeleton cap). *If T_C is a CE1 or CE2 center role and contains exactly one radial midpoint, then*

$$L_S(T_C) < \frac{3}{2}.$$

Proof. Normalize the unique midpoint as M_0 and a positive boundary overlap to $e_{0,1}$. Use the side slacks from Proposition 2.11, with $0 < \lambda < 1$ and $\rho = \sqrt{1 - \lambda + \lambda^2}$. Put

$$X = F_0(O), \quad Y = F_2(O), \quad P = \rho(1 - \rho).$$

The positive-length condition $s < t$ is equivalent to

$$(10) \quad X + (1 - \lambda)Y < P.$$

The boundary trace on $e_{0,1}$ is

$$\ell_{01} = 1 - \lambda + Y - \frac{X + Y + 1 - \rho}{\lambda},$$

and the possible second trace on $e_{5,0}$ has length

$$\ell_{50} = \left[\frac{P - (\lambda X + Y)}{1 - \lambda} \right]_+.$$

Parametrizing the six radial arms by distance from O , direct substitution in the three side slacks gives the upper bounds

$$\frac{i}{\mathcal{H}^1(T_C \cap r_i)} \mid \begin{array}{cccccc} 0 & 1 & 2 & 3 & 4 & 5 \\ \rho - X - Y & Y/\lambda & Y & \min\{X/\lambda, Y/(1-\lambda)\} & X & X/(1-\lambda). \end{array}$$

Adding boundary and radial contributions yields

$$L_S(T_C) \leq K_\lambda + \mathcal{E}(X, Y),$$

where

$$K_\lambda = 1 - \lambda + \rho - \frac{1 - \rho}{\lambda}$$

and

$$\mathcal{E}(X, Y) = -\frac{X}{\lambda} + \frac{X}{1-\lambda} + Y + \min\left\{\frac{X}{\lambda}, \frac{Y}{1-\lambda}\right\} + \left[\frac{P - (\lambda X + Y)}{1-\lambda}\right]_+.$$

Splitting according to which term realizes the minimum and whether the positive part vanishes gives, in all four cases,

$$(11) \quad \mathcal{E}(X, Y) \leq \frac{P}{1-\lambda}.$$

For example, if $X/\lambda \leq Y/(1-\lambda)$, the expression without the positive part is $X/(1-\lambda) + Y$; if the positive part is active, the excess over $P/(1-\lambda)$ is $X - \lambda Y/(1-\lambda) \leq 0$. In the opposite order, the corresponding excess is $Y - (1-\lambda)X/\lambda \leq 0$. If the positive part vanishes, (10) gives the same bound strictly.

It remains only to estimate a one-variable expression:

$$\frac{3}{2} - K_\lambda - \frac{P}{1-\lambda} = \frac{1 + A - 2\rho A}{2\lambda(1-\lambda)}, \quad A = 1 + \lambda - \lambda^2.$$

Both sides in $1 + A > 2\rho A$ are positive, and

$$(1 + A)^2 - 4A^2\rho^2 = \lambda^2(1-\lambda)^2(5 + 4\lambda - 4\lambda^2) > 0.$$

Thus $K_\lambda + P/(1-\lambda) < 3/2$, proving the lemma. $\square$

Lemma 3.3 (Positive-support skeleton cap). *Let T_i be the closure of an original open vertex role, so $V_i \in \text{int}(T_i)$. If T_i has positive-length intersection with an adjacent radial arm, then*

$$L_S(T_i) < \frac{3}{2}.$$

Proof. Normalize to $i = 0$ and translate the hexagon by V_0 :

$$\tilde{H} = V_0 + H, \quad \tilde{O} = V_0, \quad W_j = V_0 + V_j.$$

Then $W_3 = O$, $W_2 = V_1$, and $W_4 = V_5$. The five original skeleton components on which a unit triangle containing V_0 can have positive-length trace are

$$[V_0, V_1], [V_0, V_5], [O, V_1], [O, V_0], [O, V_5].$$

All five belong to the skeleton $\tilde{S}$ of $\tilde{H}$. Since $V_0 = \tilde{O}$ is interior to T_0 , the triangle is a legitimate center role for $\tilde{H}$. Its positive adjacent-arm trace is a positive boundary-edge trace of $\tilde{H}$, so it is CE1 or CE2 by Proposition 2.3. Proposition 2.11 and Lemma 3.2 therefore give

$$L_{\tilde{S}}(T_0) < \frac{3}{2}.$$

A nonlocal component of S is more than unit distance from the interior point V_0 , apart from zero-measure endpoints. Hence $L_S(T_0) \leq L_{\tilde{S}}(T_0)$. $\square$

Lemma 3.4 (No-support skeleton cap). *If a vertex role has $A_i + B_i \leq 1$ and no positive-length trace on either adjacent radial arm, then*

$$L_S(T_i) \leq 2.$$

Proof. Its boundary contribution is $A_i + B_i \leq 1$ by (17). For $P = sV_j \in r_j \setminus \{O\}$,

$$\|V_i - P\|^2 = 1 + s^2 - 2s \cos \frac{(j-i)\pi}{3} > 1$$

when $j \notin \{i-1, i, i+1\}$. The two adjacent arms have zero-length trace by hypothesis, so only the own unit arm can contribute positive length. This adds at most one. $\square$

For the remaining cap we use one specialized part of the exact local equilateral enclosure criterion. In local directions u, v meeting at 120° , suppose a closed unit triangle contains

$$0, \quad au, \quad bv, \quad c(u+v),$$

with $0 \leq a \leq b \leq 1$ and $a + b > 1$. The support-line calculation gives the selected supercritical cell

$$(12) \quad \begin{aligned} a^2 + ab + b^2 &\leq 1, & c &\leq \frac{1}{2}, \\ (a^2 - 1)c^2 + (2ab^2 + b)c + (b^4 - b^2) &\leq 0, \end{aligned}$$

where c lies on the connected component containing $c = 0$. To see the selector directly, the quadratic is negative at $c = 0$, positive at $c = 1/2$, and opens downward. For the middle assertion, four times its value at $1/2$ is increasing in a and, at $a = 1 - b$, equals $b^2(2b - 1)^2$; it is therefore positive when $a + b > 1$. Hence $c \leq 1/2$ selects its smaller-root component. Formula (12) follows by fixing in the support sum the three active contacts $\{B\}$, $\{B, c(u+v)\}$, and $\{A\}$, and squaring the positive unit-side equation.

Lemma 3.5 (Supercritical skeleton cap). *If $A_i + B_i > 1$, then*

$$L_S(T_i) < \frac{3}{2}.$$

Proof. Theorem 3.1 excludes both adjacent arms. Thus, after normalizing and writing $a = A_i$, $b = B_i$, the skeleton trace has length $a + b + c$, where c is the own-arm reach. Reflect so that $a \leq b$, and put

$$s = a + b, \quad \delta = b - a, \quad c_0 = \frac{3}{2} - s.$$

The diameter constraint gives

$$1 < s \leq \frac{2}{\sqrt{3}}, \quad 0 \leq \delta \leq \sqrt{4 - 3s^2}.$$

Let $P(c)$ denote the quadratic in (12). Substitution and multiplication by 16 give

$$\begin{aligned} 16P(c_0) = Q(s, \delta) &= \delta^4 + 8\delta^3s - 6\delta^3 + 14\delta^2s^2 - 18\delta^2s + 5\delta^2 \\ &\quad - 8\delta s^3 + 30\delta s^2 - 34\delta s + 12\delta \\ &\quad + s^4 - 6s^3 - 19s^2 + 60s - 36. \end{aligned}$$

Its derivative factors as

$$\frac{\partial Q}{\partial \delta} = 2(2\delta + 4s - 3)(\delta^2 + \delta(4s - 3) + (s - 1)(2 - s)) \geq 0.$$

Therefore

$$Q(s, \delta) \geq Q(s, 0) = (s - 1)(s^3 - 5s^2 - 24s + 36) > 0.$$

Indeed, the cubic is decreasing on $[1, 2/\sqrt{3}]$ and at its right endpoint equals $88/3 - 136\sqrt{3}/9 > 0$. Hence $P(c_0) > 0$. Since (12) selects the smaller-root component, $P(c) \leq 0$ forces $c < c_0$. Thus $a + b + c < 3/2$. $\square$

Lemma 3.6 (A positive-support rescuer is forced). *Suppose T_C is CE1 or CE2 and, after symmetry, contains exactly M_0 . If at least two vertex rows are supercritical, then a vertex role distinct from a chosen supercritical role has positive-length adjacent-arm support.*

Proof. Choose a supercritical index $s \neq 0$. Lemma 2.10 gives $M_s \notin T_s$, and $M_s \notin T_C$ by the unique-midpoint property. In the original open cover, M_s is therefore contained by some other open vertex role. Diameter one restricts this role to U_{s-1} or U_{s+1} . Because it contains an interior point of the adjacent arm, its intersection with that arm has positive length. $\square$

Theorem 3.7 (Conditional skeleton obstruction). *If T_C is CE1 or CE2, $O \in \text{int}(T_C)$, and $N_+ \geq 2$, the seven roles cannot cover S .*

Proof. Proposition 2.11 supplies the exact-one-midpoint hypothesis. Two supercritical roles, the forced positive-support rescuer, and the center each contribute strictly less than $3/2$, by Lemmas 3.5, 3.6, 3.3, and 3.2. The remaining three rows contribute at most 2 each; a further positive-support or supercritical row only improves this estimate. Therefore

$$L_S(T_C) + \sum_{i=0}^5 L_S(T_i) < \frac{3}{2} + 2\left(\frac{3}{2}\right) + \frac{3}{2} + 3 \cdot 2 = 12,$$

contradicting (9) and $\mathcal{H}^1(S) = 12$. $\square$

Proposition 3.8 (CE2 mixed positive-support obstruction). *Assume T_C is CE2, $N_+ = 1$, exactly one vertex role is Vd1 or Vd2, and $k \geq 1$ additional vertex roles have positive-length adjacent-arm support. Then the seven roles cannot cover S .*

Proof. The unique supercritical role has no adjacent support by Theorem 3.1; it is therefore distinct from the Vd1/Vd2 role and the k additional positive-support roles. The center, supercritical role, Vd1/Vd2 role, and the k additional roles all have skeleton trace strictly below $3/2$. Each of the remaining $4 - k$ roles is nonsupercritical and has no adjacent support, so its trace is at most 2. Consequently

$$L_S(T_C) + \sum_{i=0}^5 L_S(T_i) < \frac{3}{2}(3 + k) + 2(4 - k) = \frac{25 - k}{2} \leq 12.$$

For $k = 1$ the numerical right side equals 12, but the preceding trace bounds are strict, so the total is still strictly below 12. $\square$

3.3. Diagonal trace caps.

Theorem 3.9 (Diagonal trace table). *For closures of original open roles,*

role	hypothesis	L_D
T_C	CE1/CE2, $O \in \text{int}(T_C)$	$< \frac{3}{2}$
T_i	Vd0, $A_i + B_i \leq 1$	≤ 1
T_i	Vd0, $A_i + B_i > 1$	$< \frac{1}{2}$
T_i	T3-like	$< \frac{1}{2}$.

Proof. A vertex role T_i can meet only r_{i-1}, r_i, r_{i+1} in positive length. Indeed, for $P = sV_j \in r_j \setminus \{O\}$,

$$\|V_i - P\|^2 = 1 + s^2 - 2s \cos \frac{(j-i)\pi}{3} > 1$$

when $j \notin \{i-1, i, i+1\}$. A Vd0 role therefore meets only its own unit arm, proving the first Vd0 cap. If it is supercritical, the self-midpoint obstruction gives $M_i \notin T_i$. Since $T_i \cap r_i$ is an interval containing V_i , its length is then strictly below $1/2$.

For the center cap, use the notation $\lambda, \rho, C_0, C_1, C_2$ from Proposition 2.11. The positive boundary overlap gives

$$C_0 + (1 - \lambda)C_2 < \rho(1 - \rho).$$

Direct radial exit bounds are

i	0	1	2	3	4	5
$\mathcal{H}^1(T_C \cap r_i)$ is at most	C_1	C_2/λ	C_2	C_0/λ	C_0	$C_0/(1 - \lambda)$.

Thus, using $C_0 + C_1 + C_2 = \rho$,

$$\begin{aligned} L_D(T_C) &\leq \rho + \frac{C_0 + (1 - \lambda)C_2}{\lambda(1 - \lambda)} \\ &< \rho + \frac{\rho(1 - \rho)}{1 - \rho^2} = \rho + \frac{\rho}{1 + \rho} < \frac{3}{2}. \end{aligned}$$

The last inequality follows from

$$\frac{3}{2} - \rho - \frac{\rho}{1 + \rho} = \frac{(1 - \rho)(2\rho + 3)}{2(1 + \rho)} > 0.$$

It remains to treat T3-like roles. Apply the closed-trace majorant $\widehat{T}$ from Proposition 2.6, reflected so its supported adjacent arm is r_1 . In its Type II parameters,

$$0 < t < 1, \quad \Delta = 1 - t + t^2, \quad z = \sqrt{\Delta}, \quad 0 < \beta < z,$$

and positive support is equivalent to

$$\beta < \frac{z - \Delta}{t} = \frac{(1 - t)z}{1 + z}.$$

The own-arm length and adjacent-arm length are respectively

$$c = \frac{\beta}{1 - t}, \quad d = \frac{z - \Delta - t\beta}{1 - t}.$$

No other arm contributes. Hence

$$L_D(T_i) \leq L_D(\widehat{T}) = \beta + \frac{z - \Delta}{1 - t} < \frac{z - \Delta}{t(1 - t)} = \frac{z}{1 + z} < \frac{1}{2}.$$

□

3.4. Vd1/Vd2 corner estimates. We next isolate the much stronger trace loss caused by Vd1 and Vd2 geometry. For the following two results use the temporary coordinates

$$X = V_0 + x(V_5 - V_0) + y(V_1 - V_0).$$

Thus $V_0 = (0, 0)$, $V_5 = (1, 0)$, $V_1 = (0, 1)$, $O = (1, 1)$, and $\|(x, y)\|^2 = x^2 + y^2 - xy$.

Proposition 3.10 (Vd1/Vd2 corner normal form). *Let T be the closure of an original Vd1 or Vd2 role at V_0 , and let a, b be its exact reaches toward V_5, V_1 . There is a unique $t > 0$ such that, with $d = \sqrt{t^2 + t + 1}$,*

$$T = \left\{ (x, y) : \begin{array}{l} x - (t+1)y \leq a, \\ ty - (t+1)x \leq tb, \\ tx + y \leq d - a - tb \end{array} \right\}.$$

The raw traces, in coordinates measured from the indicated outer vertex, are

$$\begin{array}{l} r_0 \left| \begin{array}{l} 0 \leq s \leq \frac{d - a - tb}{t + 1} \\ \frac{t(1 - b)}{t + 1} \leq s \leq \frac{d - a - tb - 1}{t} \\ \frac{1 - a}{t + 1} \leq s \leq d - a - tb - t. \end{array} \right. \\ r_1 \\ r_5 \end{array}$$

Intersecting with $[0, 1]$ gives the actual traces. Moreover,

$$(13) \quad a + b < \frac{1}{2}.$$

Proof. A Vd1/Vd2 triangle has one vertex W outside H and two vertices U, Z in H . Since V_0 is an interior convex combination of them, both coordinates of W are negative. The two axis endpoints $(a, 0), (0, b)$ lie on distinct sides incident with W . Put $p = U - W$, $q = Z - W$. Both vectors have positive coordinates and satisfy

$$\|p\| = \|q\| = \|p - q\| = 1.$$

The order in which the two sides cross the positive axes selects the 60° rotation

$$q = (p_x - p_y, p_x), \quad p_x > p_y > 0.$$

Writing $t = (p_x - p_y)/p_y$ gives uniquely

$$p = \frac{(t+1, 1)}{d}, \quad q = \frac{(t, t+1)}{d}.$$

The two side lines through the axis endpoints and the parallel third side are exactly those in (3.10). Substitution of $(s, s), (s, 1), (1, s)$ gives (3.10).

If the r_1 trace has positive length, its lower endpoint is smaller than its upper endpoint. Clearing denominators gives

$$(t+1)a + tb < (t+1)(d-1) - t^2.$$

Since the left side is at least $t(a+b)$,

$$a + b < \frac{(t+1)(d-1) - t^2}{t} < \frac{1}{2}.$$

For the last inequality, both sides are positive and

$$\left(t^2 + \frac{3t}{2} + 1\right)^2 - (t+1)^2(t^2 + t + 1) = \frac{t^2}{4} > 0.$$

Reflection proves the same conclusion when the supported arm is r_5 . $\square$

Lemma 3.11 (Vd2 neighboring-midpoint cap). *If an original Vd2 role contains either neighboring midpoint, then its exact boundary reaches satisfy*

$$a + b < \frac{1}{3}.$$

Proof. In Proposition 3.10, reflection sends (t, a, b, M_1, M_5) to $(1/t, b, a, M_5, M_1)$, so assume $t \geq 1$ and put $U = a + tb$.

If $M_5 = (1, 1/2)$ is contained, the third half-plane in (3.10) gives

$$a + b \leq U \leq d - t - \frac{1}{2} \leq \sqrt{3} - \frac{3}{2} < \frac{1}{3};$$

the middle expression decreases for $t \geq 1$.

If $M_1 = (1/2, 1)$ is contained, the second and third half-planes give

$$b \geq \frac{t-1}{2t}, \quad a + b \leq S(t) := d - t - \frac{1}{2t}.$$

Positive-length contact with the other adjacent arm r_5 gives, by (3.10),

$$a + b < R(t) := \frac{2(t+1)d - 3t^2 - t - 2}{2t}.$$

Direct differentiation shows that S is increasing and R decreasing on $[1, \infty)$. For example, the sign calculation for S' reduces to

$$(2t+1)^2 t^4 - (t^2 + t + 1)(2t^2 - 1)^2 = (t+1)(t^3 + 3t^2 - 1) > 0,$$

while $R' < 0$ follows from $d > t + 1/2$. Splitting at $t = 4/3$ yields

$$\begin{aligned} 1 \leq t \leq \frac{4}{3} : \quad a + b &\leq S(4/3) = \frac{8\sqrt{37} - 41}{24} < \frac{1}{3}, \\ t \geq \frac{4}{3} : \quad a + b &< R(4/3) = \frac{7\sqrt{37} - 39}{12} < \frac{1}{3}. \end{aligned}$$

The final inequalities follow respectively from $8\sqrt{37} < 49$ and $7\sqrt{37} < 43$. $\square$

3.5. Exact center intervals and boundary trace caps. The next two lemmas turn the unique-midpoint normalization from Proposition 2.11 into exact boundary data.

Lemma 3.12 (CE1 maximal interval). *Normalize an exact- M_0 CE1 center triangle so that its only positive-length boundary trace is*

$$T_C \cap e_{0,1} = [s, t].$$

Then $0 < s < t < 1$ and

$$(14) \quad t \leq f(s) := \sqrt{s^2 - s + 1} - (1 - s)^2.$$

In the closed relaxation $O \in T_C$, the maximal intervals under inclusion are exactly $[s, f(s)]$, $0 < s < 1$. They are mutually incomparable.

Proof. Use the side slacks in the proof of Proposition 2.11. Exact M_0 and positive overlap force $0 < \lambda < 1$. The center condition gives

$$t \leq \rho - \lambda(1 - s), \quad \rho = \sqrt{1 - \lambda + \lambda^2}.$$

Since CE1 has no positive-length trace on $e_{5,0}$, restricting the slacks to that edge gives

$$t \geq \rho - \frac{\lambda^2}{1 - \lambda} s.$$

Compatibility implies $\lambda \geq 1 - s$. Put $q = 1 - s$ and write $\lambda = q + d$, $d \geq 0$. Then

$$t \leq \sqrt{1 - \lambda + \lambda^2} - q\lambda \leq \sqrt{1 - q + q^2} - q^2 = f(s).$$

For completeness, the second inequality follows after squaring from

$$d(1 - q^2) \leq 1 - 2q + 2q\sqrt{1 - q + q^2};$$

its left side is at most $(1 - q)(1 - q^2)$, and the difference is

$$q\sqrt{1 - q + q^2} \left(2 - \sqrt{1 - q + q^2}\right) > 0.$$

For $0 < s < 1$, choose $\lambda = 1 - s$ and $t = f(s)$. Both displayed center and no- $e_{5,0}$ inequalities are equalities. The target length is

$$t - s = \rho - \rho^2 = \rho(1 - \rho) > 0, \quad \rho = \sqrt{s^2 - s + 1},$$

and direct substitution gives the strict exclusions of M_1 and M_5 ; the remaining midpoints are then excluded by the same side slacks. Thus the frontier is feasible in the closed relaxation. Finally,

$$f'(s) = 2 - 2s + \frac{2s - 1}{2\sqrt{s^2 - s + 1}} > 0,$$

so two different frontier intervals have both left and right endpoints strictly ordered and neither contains the other. $\square$

Lemma 3.13 (CE2 interval-pair characterization). *Normalize an exact- M_0 CE2 center triangle so that*

$$T_C \cap e_{5,0} = [x, u], \quad T_C \cap e_{0,1} = [y, v].$$

Put $S_0 = x + y$ and $\Delta = \sqrt{x^2 + xy + y^2}$. The feasible interval pairs are exactly those satisfying

$$0 < x < u < 1, \quad 0 < y < v < 1,$$

$$(15) \quad (u + v)S_0 - xy = \Delta,$$

$$(16) \quad uS_0 \geq x, \quad vS_0 \geq y, \quad uS_0 < x + \frac{y}{2}, \quad vS_0 < y + \frac{x}{2}.$$

Every such pair is maximal under product interval inclusion. This is the exact maximal closed interval-pair domain. When the pair is the closure of an original open center role, the two center inequalities retain the strict form

$$uS_0 > x, \quad vS_0 > y.$$

Proof. The point V_0 cannot belong to T_C : otherwise O, V_0 , at distance one, would be two vertices of the unit triangle, and its third vertex would be V_1 or V_5 , allowing only one of the two stated boundary overlaps. The unique side violated at V_0 therefore cuts the two initial endpoints x, y . Its outward normal and the two rotations by 120° are

$$n_2 = \frac{1}{2\Delta}(\sqrt{3}(x+y), y-x), \quad n_0 = R_{2\pi/3}n_2, \quad n_1 = R_{-2\pi/3}n_2.$$

Writing the triangle as $n_j \cdot X \leq \alpha_j$, the four endpoints give

$$\begin{aligned} \alpha_2 &= \frac{\sqrt{3}(S_0 - xy)}{2\Delta}, \\ \alpha_0 &= \frac{\sqrt{3}(vS_0 - y)}{2\Delta}, \\ \alpha_1 &= \frac{\sqrt{3}(uS_0 - x)}{2\Delta}. \end{aligned}$$

The unit side condition $\alpha_0 + \alpha_1 + \alpha_2 = \sqrt{3}/2$ is precisely (15). The conditions $O \in T_C$ are the first two weak inequalities in (16). Substitution of M_1 and M_5 shows that their exclusion is exactly the last two strict inequalities; these also exclude M_2, M_3, M_4 , while the weak inequalities imply $M_0 \in T_C$. Restricting the three side slacks to the two target edges gives exactly $[x, u]$ and $[y, v]$. On each remaining edge one of the following necessary slack sums is negative:

$$\Delta + xy - (2x + y), \quad \Delta + xy - (x + 2y),$$

because

$$\begin{aligned} (2x + y - xy)^2 - \Delta^2 &= x(x(1-y)(3-y) + y(3-2y)) > 0, \\ (x + 2y - xy)^2 - \Delta^2 &= y(x(3-2x) + y(1-x)(3-x)) > 0. \end{aligned}$$

Thus there are no further positive boundary traces.

For maximality define

$$F(x, y) = \frac{\Delta + xy}{x + y} = u + v.$$

The weak center inequalities imply the start domain $2x + 2y - xy \geq 1$. On it,

$$\frac{\partial F}{\partial x} = \frac{y(x - y + 2y\Delta)}{2\Delta(x + y)^2} > 0, \quad \frac{\partial F}{\partial y} = \frac{x(y - x + 2x\Delta)}{2\Delta(x + y)^2} > 0.$$

If another feasible pair product-dominated the given one, its initial coordinates would be no larger and its terminal coordinates no smaller. Monotonicity would make its terminal sum no larger, while domination makes it no smaller. Equality forces all four endpoints to agree. $\square$

Theorem 3.14 (Boundary trace table). *For closures of original open roles, the following bounds hold:*

<i>role</i>	<i>hypothesis</i>	$L_{\partial H}$
T_C	CE0	0
T_C	CE1	$\leq \frac{\sqrt{3}}{2} - \frac{3}{4}$
T_C	CE2	$< \frac{1}{2}$
T_i	Vd0, $A_i + B_i \leq 1$	$\leq \frac{1}{2}$
T_i	Vd0, $A_i + B_i > 1$	$\leq \frac{2}{\sqrt{3}}$
T_i	Vd1/Vd2	$< \frac{1}{2}$
T_i	T3-like	< 1 .

Proof. For every vertex role,

$$(17) \quad L_{\partial H}(T_i) = A_i + B_i.$$

Indeed, the two incident traces are initial intervals of these lengths. A relative-interior point of any nonincident edge is more than unit distance from V_i , whereas V_i is interior to the diameter-one triangle T_i . This excludes any other positive-length boundary trace.

The CE0 row is the definition. In the CE1 row put

$$\rho = \sqrt{s^2 - s + 1} \in [\sqrt{3}/2, 1).$$

Lemma 3.12 gives

$$L_{\partial H}(T_C) = t - s \leq \rho(1 - \rho) \leq \frac{\sqrt{3}}{2} - \frac{3}{4}.$$

For CE2, Lemma 3.13 gives

$$\begin{aligned} L_{\partial H}(T_C) &= (u - x) + (v - y) \\ &= \frac{\Delta + xy}{S_0} - S_0 = \frac{\Delta(1 - \Delta)}{S_0}. \end{aligned}$$

The center inequalities imply $\Delta + xy \geq S_0$, whence $S_0 \geq (1 + xy)/2 > 1/2$ after squaring. Since the trace is positive, $0 < \Delta < 1$, and therefore

$$L_{\partial H}(T_C) \leq \frac{1}{4S_0} < \frac{1}{2}.$$

For a nonsupercritical Vd0 row, (17) is enough. For a supercritical row, its incident endpoints are at squared distance $A_i^2 + A_i B_i + B_i^2 \leq 1$, and hence

$$\frac{3}{4}(A_i + B_i)^2 \leq 1.$$

This proves the $2/\sqrt{3}$ cap. The Vd1/Vd2 row is (13). Finally, the Type II calculation in Proposition 2.6 gives for an original T3-like role

$$A_i + B_i = z - \alpha < z < 1$$

because $0 < t < 1$ and $\alpha > 0$. □

3.6. Terminal length contradictions.

Proposition 3.15 (Boundary-length branches). *The following branches are impossible:*

- (1) *CE0 with $N_+ = 0$;*
- (2) *CE0 with $N_+ = 1$ and at least one Vd1/Vd2 role;*
- (3) *CE1 or CE2 with $N_+ = 0$ and at least one Vd1/Vd2 role;*
- (4) *CE1 with $N_+ = 1$ and at least one Vd1/Vd2 role;*
- (5) *CE2 with $N_+ = 1$ and at least two Vd1/Vd2 roles.*

Proof. In CE0, the center open role contains no relative-interior boundary point. No nonincident vertex role can contain such a point either. Hence on every edge the two incident open vertex traces form two nonempty relatively open sets covering a connected interval; they overlap in positive length. Thus

$$1 < B_i + A_{i+1}$$

on every edge. Summing gives $6 < \sum_i (A_i + B_i)$, contradicting $N_+ = 0$. This proves (1).

For the other cases apply Theorem 3.14. Put

$$\gamma_1 = \frac{\sqrt{3}}{2} - \frac{3}{4}.$$

Vd1, Vd2, and T3-like roles are nonsupercritical, so whenever $N_+ = 1$ the unique supercritical role in these cases is Vd0. The respective total trace bounds are

case	$L_{\partial H}(T_C) + \sum_i L_{\partial H}(T_i)$
CE0, $N_+ = 1, \geq 1$ Vd1/Vd2	$< \frac{1}{2} + \frac{2}{\sqrt{3}} + 4 < 6$
CE1, $N_+ = 0, \geq 1$ Vd1/Vd2	$< \gamma_1 + \frac{1}{2} + 5 < 6$
CE2, $N_+ = 0, \geq 1$ Vd1/Vd2	$< \frac{1}{2} + \frac{1}{2} + 5 = 6$
CE1, $N_+ = 1, \geq 1$ Vd1/Vd2	$< \gamma_1 + \frac{1}{2} + \frac{2}{\sqrt{3}} + 4 < 6$
CE2, $N_+ = 1, \geq 2$ Vd1/Vd2	$< \frac{1}{2} + 2\left(\frac{1}{2}\right) + \frac{2}{\sqrt{3}} + 3 < 6.$

Each contradicts (9) for the length-six boundary. The numerical inequalities follow, for example, from $9/2 + 2/\sqrt{3} < 6$ and $15/4 + 7/(2\sqrt{3}) < 6$. $\square$

Proposition 3.16 (Vd2 neighboring-midpoint branch). *Assume a CE2, $N_+ = 1$ configuration has exactly one Vd1/Vd2 role, that role is Vd2 and contains a neighboring midpoint, and all other vertex roles are Vd0. Then the boundary cannot be covered.*

Proof. Lemma 3.11 and Theorem 3.14 give

$$L_{\partial H}(T_C) + \sum_i L_{\partial H}(T_i) < \frac{1}{2} + \frac{1}{3} + \frac{2}{\sqrt{3}} + 4 < 6.$$

The last inequality is equivalent to $12 < 7\sqrt{3}$, whose square is $144 < 147$. $\square$

Proposition 3.17 (At least two T3-like roles). *Assume the center is CE1 or CE2, $N_+ = 1$, no vertex role is Vd1 or Vd2, and at least two vertex roles are T3-like. Then the diagonal target D cannot be covered.*

Proof. Let $k \geq 2$ be the number of T3-like roles. By Lemma 2.8, the unique supercritical role is Vd0; $2 \leq k \leq 5$, and the remaining $5 - k$ roles are nonsupercritical Vd0. Theorem 3.9 gives

$$L_D(T_C) + \sum_i L_D(T_i) < \frac{3}{2} + \frac{1}{2} + \frac{k}{2} + (5 - k) = 7 - \frac{k}{2} \leq 6,$$

contradicting $\mathcal{H}^1(D) = 6$ and (9). $\square$

Proposition 3.18 (Branches closed by Strategy 1). *Direct length-sum obstructions close precisely the following rows of the final routing table:*

- (1) CE0, $N_+ = 0$;
- (2) CE0, $N_+ = 1$, with a Vd1/Vd2 role;
- (3) CE1/CE2, $N_+ = 0$, with a Vd1/Vd2 role;
- (4) CE1, $N_+ = 1$, with a Vd1/Vd2 role;
- (5) CE2, $N_+ = 1$, with at least two Vd1/Vd2 roles;
- (6) CE1/CE2, $N_+ = 1$, with no Vd1/Vd2 role and at least two T3-like roles;
- (7) CE1/CE2, $N_+ \geq 2$;
- (8) within the CE2, $N_+ = 1$ package with exactly one Vd1/Vd2 role, the Vd2 neighboring-midpoint subcase and every subcase with an additional positive-support role.

Proof. Items (1)–(5) are Proposition 3.15; item (6) is Proposition 3.17; item (7) is Theorem 3.7; and the two subfamilies in item (8) are Propositions 3.16 and 3.8. These are only the indicated subcases of the exceptional CE2 row; its remaining placements are handled by the exact-demand strategy. $\square$

4. STRATEGY 2: EXACT DEMAND PROPAGATION

This section gives the reusable geometric mechanism and the shortened proofs that drive Strategy 2. The exact admissible-set classification, all radical comparisons, the complete four-label tables, and the unchanged terminal polynomial estimates are recorded in Appendix A and its following Strategy 2 sections. In the body, every formal map is connected explicitly to an actual nonsupercritical row.

4.1. The local propagation interface. Let

$$u = \left(\frac{1}{2}, \frac{\sqrt{3}}{2} \right), \quad v = \left(\frac{1}{2}, -\frac{\sqrt{3}}{2} \right),$$

and for $0 \leq a, b, c \leq 1$ put

$$K(a, b, c) = \text{conv}\{0, au, bv, c(u + v)\}.$$

The local admissible set is

$$\mathcal{A} = \{(a, b, c) : K(a, b, c) \text{ lies in a closed unit equilateral triangle}\}.$$

For $a, c \in [0, 1]$ define

$$B_c(a) = \max\{b : (a, b, c) \in \mathcal{A}\}, \quad F_c(a) = \min\{B_c(a), 1 - a\}, \quad G_c(a) = 1 - F_c(a).$$

Propositions A.2 and A.3 prove that $\mathcal{A}$ is coordinatewise down-closed, F_c is nonincreasing, and G_c is nondecreasing and extensive:

$$(18) \quad G_c(z) \geq z.$$

They also prove the exact reflection duality

$$(19) \quad G_c(a) \leq z \iff G_c(1-z) \leq 1-a.$$

Suppose an actual row has maximal reaches (A_i, B_i, C_i) and is nonsupercritical. If

$$A_i \geq z, \quad C_i \geq c_i, \quad A_i + B_i \leq 1,$$

then the same role realizes the lower-bound demand triple (z, B_i, c_i) and therefore

$$B_i \leq F_{c_i}(z).$$

If the next boundary handoff gives $A_{i+1} \geq 1 - B_i$, then

$$(20) \quad A_{i+1} \geq 1 - B_i \geq 1 - F_{c_i}(z) = G_{c_i}(z).$$

Every composition below is shorthand for repeated applications of (20) to the stated actual rows.

4.2. A universal normal form for every selected T_+ arc. The exact capped map has four genuine labels: Full, L , T_- , and selected T_+ . The nonlinear part of the last label is independent of the radial deficit.

Proposition 4.1 (Universal selected- T_+ curve). *Fix $0 < d < 1/2$, put $c = 1 - d$, and suppose an input p belongs to a selected T_+ arc of G_c . Write*

$$q = G_c(p), \quad \nu = q - p, \quad x = \frac{q - d}{1 - d}.$$

Then

$$(21) \quad \nu = \psi(x) := 1 - \sqrt{1 - x + x^2},$$

and

$$(22) \quad q = d + (1 - d)x, \quad p = d + (1 - d)x - \psi(x).$$

Consequently $p \mapsto G_{1-d}(p)$ is increasing and strictly concave on every selected T_+ arc.

Proof. The selected quadratic in Proposition A.3 is

$$(1 - q)(q - d) = (1 - d)^2 \nu(2 - \nu).$$

Substituting $q = d + (1 - d)x$ and cancelling $(1 - d)^2$ gives

$$x(1 - x) = \nu(2 - \nu).$$

The selected component uses the smaller root, which is (21). Put $r(x) = 1 - x + x^2$.

Then

$$\psi'(x) = \frac{1 - 2x}{2\sqrt{r(x)}}, \quad \psi''(x) = -\frac{3}{4r(x)^{3/2}} < 0.$$

The input function

$$p_d(x) = d + (1 - d)x - \psi(x)$$

satisfies

$$p'_d(x) = 1 - d - \psi'(x) > 0, \quad p''_d(x) = -\psi''(x) > 0.$$

Thus p_d is increasing and strictly convex. Its inverse is increasing and strictly concave, and $q = d + (1 - d)x$ is affine in that inverse. $\square$

Two chord bounds will be used repeatedly. In the high-radial range put $e(d) = \ell(1-d)$. The selected arc has endpoints

$$(p, q) = (d, d), \quad (p, q) = (e(d), d + e(d)),$$

so

$$(23) \quad G_{1-d}(p) \geq p + \frac{d}{e(d)-d}(p-d).$$

In the low-radial range, if $t = h(1-d)$ is the order-transition value, the endpoints are (d, d) and $(t, 1-t)$, and therefore

$$(24) \quad G_{1-d}(p) \geq p + \frac{1-2t}{t-d}(p-d).$$

For algebraic calculations the same curve has a rational parameter. With $z = \psi(x)/x$,

$$(25) \quad x = \frac{1-2z}{1-z^2}, \quad \psi(x) = \frac{z(1-2z)}{1-z^2}.$$

In the historical high-sheet notation $x = 1-\beta$ and $m_\beta = \sqrt{\beta^2 - \beta + 1}$, this becomes

$$(26) \quad \beta = \frac{z(2-z)}{1-z^2}, \quad m_\beta = \frac{1-z+z^2}{1-z^2}.$$

This is the normalization used in the shortened 407X high-sheet proof.

4.3. Threshold routing. For $0 < d < 1 - \sqrt{3}/2$, the exact high-demand threshold is

$$(27) \quad z > e(d) \implies G_{1-d}(z) \geq 1 - e(d).$$

The following lemma isolates the only logical use of a long map composition after such a threshold has been crossed.

Lemma 4.2 (Threshold routing). *Let $z_j = G_j(z_{j-1})$, where every G_j is nondecreasing and extensive. If $G_k = G_{1-d}$ and $z_{k-1} > e(d)$, then*

$$z_j \geq 1 - e(d) \quad (j \geq k).$$

In particular, if $e(d) < Q$, then every later output is strictly larger than $1 - Q$.

Suppose instead that a chain contains later occurrences of both G_{1-d_1} and G_{1-d_2} , that

$$z_0 > e(d_1), \quad \min\{e(d_1), e(d_2)\} < Q.$$

Then one of the two threshold rows forces every subsequent output above $1 - Q$.

Proof. Equation (27) gives $z_k \geq 1 - e(d)$, and extensivity preserves this bound. For the two-threshold statement, if $e(d_1) < Q$, route the initial strict inequality to the G_{1-d_1} row. Otherwise

$$z_0 > e(d_1) \geq Q > e(d_2),$$

so route it to the G_{1-d_2} row. The first triggered row gives the desired strict output, and all later maps preserve it. $\square$

The lemma does not replace the actual-row argument. It is applied only after (20) has identified the formal iterates as lower bounds for the actual incoming reaches.

4.4. The CE1 one-gap chain with the shortened $X > 1/2$ proof. We give the full logical route and isolate the unchanged terminal estimate in one appendix lemma. Use the exact CE1 variables of Proposition A.4:

$$\begin{aligned} R = \lambda, \quad w = 1 - R, \quad E = \sqrt{1 - R + R^2} = \sqrt{1 - Rw}, \\ \eta = 1 - E, \quad P = \eta E, \quad A = C_0, \quad D = C_2. \end{aligned}$$

They satisfy

$$(28) \quad A, D > 0, \quad A + wD < P, \quad D + RA \geq P, \quad A < \frac{w}{2}, \quad D < \frac{R}{2}.$$

Put

$$X = R - D, \quad m = \frac{A}{R}, \quad H = \frac{\eta + A + D}{2R}.$$

The five high radial demands are

$$c_1 = 1 - \frac{D}{R}, \quad c_2 = 1 - D, \quad c_3 = 1 - m, \quad c_4 = 1 - A, \quad c_5 = 1 - \frac{A}{w}.$$

A gap in the center interval $[s, t] \subset e_{0,1}$ gives $a_1 \geq X = 1 - t$ and $b_0 \geq s$. Applying (20) to rows 1, 2, 3, 4, 5 gives

$$(29) \quad a_0 \geq Z := (G_{c_5} \circ G_{c_4} \circ G_{c_3} \circ G_{c_2} \circ G_{c_1})(X).$$

The supercritical row realizes (s, a_0, c_0) after reflection and coordinatewise relaxation, so

$$(30) \quad a_0 \leq B_{c_0}(s).$$

Duality and extensivity reduce the lower bound to the three-row suffix

$$(31) \quad (G_{c_2} \circ G_{c_3} \circ G_{c_4})(H) > 1 - X.$$

The elementary low-root estimates give

$$(32) \quad e(A) < X, \quad H > A.$$

For completeness, the first inequality follows from $A + wD < P$ and $D = R - X$. Indeed, with

$$f(X) = wX - \frac{X}{2(1+X)},$$

convexity and the endpoint values at $X = R/2, R$ give $f(X) < \eta$, hence

$$A < \frac{X}{2(1+X)}.$$

Together with $e(A) < 2A/(1 - 2A)$ this gives $e(A) < X$. Also

$$2R(H - A) = \eta + D + (1 - 2R)A > 0;$$

when $R > 1/2$, use $A < P = \eta E$ to obtain the strict sign.

At row 4, the Full labels are impossible. The L and T_- labels give

$$G_{1-A}(H) \geq 1 - e(A) > 1 - X.$$

It remains to consider the selected T_+ label.

Lemma 4.3 (The selected row-4 branch forces $X > 1/2$). *Under (28), if $H \leq e(A)$, then*

$$X > \frac{1}{2}.$$

Proof. Suppose $X \leq 1/2$. The selected label and the strict low-root estimate give

$$H \leq e(A) < \frac{2A}{1-2A}.$$

The other center inequality gives

$$\begin{aligned} 2RH &= \eta + A + D \\ &\geq \eta + A + P - RA \\ &= \eta + P + wA = w(R + A). \end{aligned}$$

Consequently

$$f_R(A) < 0, \quad f_R(z) := w(R + z)(1 - 2z) - 4Rz.$$

But $f_R''(z) = -4w < 0$ and $f_R(0) = Rw > 0$.

If $R \leq 1/2$, then $0 < A < P$. Using $P = (1 - E)E$ and $E^2 = 1 - R + R^2$,

$$f_R(P) = (1 - E)(4E^3 - 2E^2 + 1 - R(2E^3 - 2E^2 + 5E)).$$

The coefficient of R is positive, so

$$f_R(P) \geq \frac{1-E}{2}\phi(E), \quad \phi(E) = 6E^3 - 2E^2 - 5E + 2.$$

On $\sqrt{3}/2 \leq E < 1$,

$$\phi'(E) = 18E^2 - 4E - 5 > 0, \quad \phi(\sqrt{3}/2) = \frac{2 - \sqrt{3}}{4} > 0.$$

Thus $f_R(P) > 0$. Strict concavity places f_R above the positive chord joining its values at 0 and P , contradicting $f_R(A) < 0$.

If $R > 1/2$, then $D = R - X \geq R - 1/2$, and the first center inequality gives

$$0 < A < A_0, \quad A_0 = P - w\left(R - \frac{1}{2}\right) = \frac{w}{2} - \eta.$$

Again using $E^2 = 1 - R + R^2$,

$$f_R(A_0) = \frac{1-E}{2}(E + 11R - 3ER - 5).$$

Since $11 - 3E > 0$ and $R > 1/2$,

$$E + 11R - 3ER - 5 > \frac{11 - 3E}{2} + E - 5 = \frac{1 - E}{2} > 0.$$

Thus $f_R(A_0) > 0$, and concavity gives the same contradiction on $(0, A_0)$. Therefore $X > 1/2$. $\square$

Assume now that rows 4 and 3 are both selected T_+ . Put

$$p_1 = G_{1-A}(H), \quad p_2 = G_{1-m}(p_1).$$

The high-radial chord (23) and the low-root bounds give

$$(33) \quad p_1 > L_1 := (2 - 4A)H - (1 - 4A)A.$$

Since $X > 1/2$, one has $R > 1/2$ and $m < w/2 < 1/4$. Applying the appropriate high- or low-radial chord from (23)–(24) gives

$$(34) \quad p_2 > L_2 := (2 - 5m)L_1 - (1 - 5m)m.$$

The coefficient comparison is elementary: for $m \leq 1/8$ use $e(m) \leq 2m + 5m^2$; for $1/8 < m < 1 - \sqrt{3}/2$ use $e(m) < (1 - m)/2$; and on the low-radial arc use $h(1 - m) < 7/16 < (3 + m)/7$.

The unchanged terminal calculation in Appendix B proves from (28), the two selected labels, and (34) that

$$(35) \quad D < \frac{1}{10}, \quad p_2 > L_2 > 2D + 3D^2 > e(D).$$

Now threshold routing, rather than a new five-label table, finishes the argument. The row-2 map is G_{1-D} , so

$$G_{1-D}(p_2) \geq 1 - e(D) > 1 - X,$$

where the last inequality follows from

$$e(D) < 2D + 3D^2 < \frac{23}{100} < \frac{1}{2} < X.$$

This is (31). Duality and the extensive outer maps give

$$Z > 1 - H = 1 - \frac{s}{2}.$$

Finally the boundary diameter condition gives

$$B_{c_0}(s) \leq \frac{-s + \sqrt{4 - 3s^2}}{2} < 1 - \frac{s}{2}.$$

Together with (29) and (30), this is the contradiction

$$a_0 \geq Z > B_{c_0}(s) \geq a_0.$$

Proposition 4.4 (Simplified CE1 one-gap obstruction). *The CE1, $N_+ = 1$, all-Vd0, exactly-one-gap state is impossible.*

Proof. The actual-row induction, the suffix duality, the branch split above, and Lemma 4.3 prove the required terminal contradiction in every selected row-4 label. $\square$

4.5. CE2 one-gap routing. The CE2 proof is an especially clean use of Lemma 4.2. Normalize the exact center parameters by

$$R = \frac{x}{x+y}, \quad w = 1 - R, \quad E = \sqrt{1 - Rw}, \quad \eta = 1 - E, \quad P = \eta E,$$

and put

$$\alpha = u - R, \quad \delta = v - w, \quad k = \eta + \alpha + \delta.$$

The strict center domain is

$$\alpha, \delta > 0, \quad \delta + R\alpha < P, \quad \alpha + w\delta < P.$$

For a right-hand gap the initial input is

$$z_0 = 1 - v = R - \delta,$$

and the decisive target is

$$Q = \frac{y}{2} = \frac{k}{2R}.$$

The exact center inequalities, proved in Appendix B, give

$$(36) \quad z_0 > e(\alpha), \quad \min\{e(\alpha), e(\delta)\} < Q.$$

The five actual nonsupercritical rows give the formal chain by (20). If $e(\alpha) < Q$, the row with radial deficit α forces the output above $1 - Q$. Otherwise

$$z_0 > e(\alpha) \geq Q > e(\delta),$$

and the row with deficit δ does so. Every subsequent map is extensive. Thus the return demand satisfies

$$Z_{\text{CE2}}^R > 1 - \frac{y}{2}.$$

The diameter endpoint gives

$$B_{c_0}(y) < 1 - \frac{y}{2},$$

contradicting the two actual bounds on row 0. Reflection exchanges (x, u, α, R) with (y, v, δ, w) and proves the left-gap orientation. This is Proposition B.2.

4.6. The 407X high-sheet package. The T3-like $N_+ = 0$ branch reduces to one strict two-coordinate capped-loss inequality. Its difficult first-coordinate label is the selected high sheet. The local radical formerly written

$$m_\beta = \sqrt{\beta^2 - \beta + 1}$$

is exactly the universal curve (21) with $x = 1 - \beta$, and may be replaced by the rational coordinate (26). The center radical $\rho = \sqrt{r^2 - r + 1}$ is independent and is retained.

The exact high-sheet estimates in Appendix C prove

$$y_* < \frac{1}{10}, \quad 3y_* \leq 1 - B_5, \quad S > 3y, \quad A_C > 3y,$$

and the analytic left-high threshold needed for the right T_- label. With the universal normal form, these are the only branch-specific estimates: no separate selected- T_+ curvature calculation remains. They assemble the four genuine labels into Proposition C.5.

4.7. Strategy 2 branch assembly.

Proposition 4.5 (Branches closed by exact demand). *Strategy 2 closes all routing-table rows assigned to exact demand propagation:*

- (1) the $N_+ = 0$ all-Vd0 and T3-like-without-Vd1/Vd2 branches for CE1 and CE2;
- (2) every positive-gap $N_+ = 1$ all-Vd0 state;
- (3) the exactly-one-T3-like $N_+ = 1$ state with no Vd1/Vd2 role;
- (4) the complementary exact placements in the CE2, $N_+ = 1$, exactly-one-Vd1/Vd2 branch.

Singleton gaps are included.

Proof. The first item is Propositions C.2 and C.5. For the second item, the CE1 one-gap state is Proposition 4.4; the two CE2 one-gap orientations are Proposition B.2; the CE2 two-gap and all remaining gap states are assembled in Proposition B.3. The third item is Proposition C.1. The fourth is Proposition C.16. In every positive-gap chain, Equation (20) supplies the passage from scalar maps to actual role reaches. $\square$

5. AREA-LOSS OBSTRUCTIONS

This section proves the local inequalities and cyclic estimates used in the area-loss argument.

Let

$$A_{\Delta} = \frac{\sqrt{3}}{4}$$

be the area of a unit equilateral triangle. Throughout this section areas are divided by A_{Δ} . Thus a unit triangle has normalized area one and H has normalized area six.

5.1. The local square-loss inequalities. Fix a vertex V_i . Let P_a and P_b be the points at distances a and b from V_i on its incoming and outgoing boundary edges. For a closed unit equilateral triangle T containing V_i, P_a, P_b , put

$$G(T) = \frac{\text{area}(T \setminus H)}{A_{\Delta}}.$$

Equivalently, if

$$f(a, b) = \max_T \frac{\text{area}(T \cap H)}{A_{\Delta}}, \quad G(a, b) = 1 - f(a, b),$$

then $G(a, b)$ is the least possible normalized loss. Reflection in the radial axis interchanges a and b , and increasing either demand can only decrease f .

Lemma 5.1 (Local wedge reduction). *After rotating indices to $i = 0$, write*

$$X = V_0 + x(V_1 - V_0) + y(V_5 - V_0), \quad W = \{(x, y) : x \geq 0, y \geq 0\}.$$

If a closed unit equilateral triangle T contains V_0 , then

$$T \cap H = T \cap W.$$

Normalized Euclidean area is twice the corresponding (x, y) -coordinate area.

Proof. The two basis vectors have length one and angle 120° , so

$$\|(x, y)\|^2 = x^2 + y^2 - xy, \quad |\det(V_1 - V_0, V_5 - V_0)| = \frac{\sqrt{3}}{2}.$$

The hexagon is

$$H = \{0 \leq x, y \leq 2, |x - y| \leq 1\} \subset W.$$

Conversely, if $(x, y) \in T \cap W$, then the diameter-one condition gives $x^2 + y^2 - xy \leq 1$. Hence

$$|x - y|^2 \leq x^2 + y^2 - xy \leq 1, \quad \frac{3}{4}x^2 \leq x^2 + y^2 - xy,$$

and the same estimate holds for y . Thus $|x - y| \leq 1$ and $0 \leq x, y \leq 2/\sqrt{3} < 2$, proving $(x, y) \in H$. The determinant identity proves the area assertion. $\square$

Theorem 5.2 (Unconditional local square loss). *For every feasible pair $0 \leq a, b \leq 1$ and every realizing closed unit equilateral triangle T ,*

$$(37) \quad G(T) \geq \min(a, b)^2.$$

If $a + b > 1$, then

$$(38) \quad G(T) \geq \max(a, b)^2.$$

Consequently the same inequalities hold for $G(a, b) = 1 - f(a, b)$.

Proof. In the coordinates of Lemma 5.1, physical rotation through 60° is $R(x, y) = (x - y, x)$. Put

$$0 \leq t \leq 1, \quad D = 1 - t + t^2, \quad z = \sqrt{D}.$$

After choosing one of the six directed sides, every orientation has one of the following two forms:

$$(39) \quad \begin{array}{c|cc} & d & Rd \\ \hline \text{I} & z^{-1}(1, t) & z^{-1}(1 - t, 1) \\ \text{II} & z^{-1}(t, -(1 - t)) & z^{-1}(1, t). \end{array}$$

Indeed these forms cover, respectively, the physical direction sectors $[120^\circ, 180^\circ]$ and $[60^\circ, 120^\circ]$; their endpoints cover the common boundary orientations.

For Type I introduce

$$U = \alpha + y - tx, \quad V = \beta + x - (1 - t)y.$$

Then T is the simplex $U, V \geq 0, U + V \leq z$. Containment of $(0, 0), (0, a), (b, 0)$ gives

$$(40) \quad \alpha \geq tb, \quad \beta \geq (1 - t)a, \quad \alpha + \beta + (1 - t)b \leq z, \quad \alpha + \beta + ta \leq z.$$

The two inequalities defining W become

$$(1 - t)U + V \geq (1 - t)\alpha + \beta, \quad U + tV \geq \alpha + t\beta.$$

For fixed t , lowering α or β therefore enlarges $T \cap W$. The outside loss is minimized at $\alpha = tb, \beta = (1 - t)a$. At that placement the portion outside W has vertices, in the (U, V) -plane,

$$(0, 0), \quad (a + tb, 0), \quad (tb, (1 - t)a), \quad (0, b + (1 - t)a).$$

Since the Jacobian of $(x, y) \mapsto (U, V)$ has absolute value D ,

$$(41) \quad G(T) \geq G_I(a, b; t) = \frac{(1 - t)a^2 + tb^2 + 2t(1 - t)ab}{D}.$$

If $a \leq b$, then

$$D(G_I - a^2) = t\{b^2 - a^2 + (1 - t)(a^2 + 2ab)\} \geq 0;$$

the reflected factorization proves $G_I \geq b^2$ when $b \leq a$. This proves (37) in Type I.

Suppose now that $a + b > 1$ and $a \leq b$. Set

$$r = \frac{a}{b}, \quad t_0 = \frac{1 - r^2}{1 + 2r}.$$

Since $b > 1/(1 + r)$, feasibility in (41) and (40) excludes $0 \leq t < t_0$: for $0 < t < t_0$,

$$\left(\frac{1 + r(1 - t)}{1 + r}\right)^2 - D = \frac{t(1 + 2r)(t_0 - t)}{(1 + r)^2} > 0,$$

so $b + (1 - t)a > z$, while $t = 0$ would give $a + b \leq 1$. Hence $t \geq t_0$, and

$$D(G_I - b^2) = (1 - t)b^2\{r^2 - 1 + t(2r + 1)\} \geq 0.$$

If $b \leq a$, put $r = b/a$ and $t_1 = (r^2 + 2r)/(1 + 2r)$. The identical reflected calculation excludes $t > t_1$ and gives

$$D(G_I - a^2) = ta^2\{r^2 + 2r - t(1 + 2r)\} \geq 0.$$

Thus (38) holds in Type I.

For Type II put

$$U = \alpha + (1-t)x + ty, \quad V = \beta + tx - y.$$

Again T is the simplex $U, V \geq 0, U + V \leq z$. Its exact reaches on the positive y - and x -axes are

$$(42) \quad A = \beta, \quad B = z - \alpha - \beta, \quad A + B \leq z \leq 1.$$

In particular Type II is impossible when $a + b > 1$. Its wedge intersection is the quadrilateral with vertices

$$(0, 0), \quad (B, 0), \quad \left(\frac{B + (1-t)A}{D}, \frac{A + tB}{D} \right), \quad (0, A).$$

Hence

$$(43) \quad G(T) = G_{\text{II}}(A, B; t) = 1 - \frac{(1-t)A^2 + tB^2 + 2AB}{D}.$$

Let $S = A + B$. If $A \leq B$, write $A = rS$, $B = (1-r)S$ with $0 \leq r \leq 1/2$. Since $S^2 \leq D$, and with

$$C = (1-t)r^2 + t(1-r)^2 + 2r(1-r),$$

we have $G_{\text{II}} \geq 1 - C$. The exact factorization

$$1 - C - r^2D = (1-t)(1 - 2r + tr^2) \geq 0$$

gives $G_{\text{II}} \geq r^2D \geq A^2$. Reflection gives $G_{\text{II}} \geq B^2$ when $B \leq A$. Since $A \geq a$ and $B \geq b$, this proves (37). The two orientation types are exhaustive, so the theorem follows. $\square$

5.2. The direct T3-like loss. The next theorem uses the classification data (o, n) from Section 2: o counts triangle vertices outside H , and n counts adjacent radial arms met in positive length. Thus a T3-like role has $(o, n) = (2, 1)$.

Theorem 5.3 (Direct T3-like loss). *Let T be a closed unit V_i -triangle of T3-like type, and let a, b be lower bounds for its adjacent boundary reaches. Then*

$$(44) \quad a + b \leq 1.$$

Moreover, for $0 \leq m \leq 1/2$,

$$(45) \quad a, b \geq m \implies G(T) \geq 2m - 4m^2.$$

Proof. Use the two orientation forms (39). In Type I the three simplex vertices are

$$\begin{aligned} Q_0 &= D^{-1}(-(1-t)\alpha - \beta, -\alpha - t\beta), \\ Q_1 &= D^{-1}(z - (1-t)\alpha - \beta, tz - \alpha - t\beta), \\ Q_2 &= D^{-1}((1-t)z - (1-t)\alpha - \beta, z - \alpha - t\beta). \end{aligned}$$

The first coordinate of Q_1 and the second coordinate of Q_2 are nonnegative. If two vertices lie outside the wedge, then Q_0 and exactly one of Q_1, Q_2 do. If Q_1 is outside, the sole wedge vertex Q_2 has both coordinates strictly below one; this follows from $(1-t)/z \leq 1$, with strictness either from $t > 0$ or from $\alpha > 0$ at $t = 0$. If Q_2 is outside, the reflected estimate $t/z \leq 1$ gives the same conclusion for Q_1 . Axis intersection points are also at coordinate at most one by the diameter bound. Hence neither adjacent radial arm is met in positive length. A Type I triangle with $o = 2$ has $n = 0$, so it cannot be T3-like.

For Type II retain the exact reaches A, B in (42), put $S = A + B$, and let

$$Q = \left(\frac{B + (1-t)A}{D}, \frac{A + tB}{D} \right)$$

be the unique triangle vertex in the wedge. Equation (42) immediately gives $a + b \leq A + B \leq 1$, proving (44). The endpoint orientations $t = 0, 1$ cannot have $(o, n) = (2, 1)$; hence $0 < t < 1$. Reflecting if necessary, assume that $Q_x > 1$ and $Q_y \leq 1$. The first inequality is

$$S - tA > D.$$

Since $S \leq z$, it follows that

$$(46) \quad A < L := \frac{z - D}{t} = \frac{z(1-z)}{t}.$$

The normalized inside area is

$$I(S, A) = \frac{(1-t)A^2 + t(S-A)^2 + 2A(S-A)}{D}.$$

For fixed t, A , this increases with S , so the loss is bounded below by its value at $S = z$. There

$$G(z, A) = \frac{tA^2 + (1-t)(z-A)^2}{D},$$

and this decreases for $A < L < (1-t)z$. Thus

$$(47) \quad G(T) \geq G(z, L).$$

Set $r = (1-z)/t$. Then $0 < r < 1/2$, and eliminating t, z gives

$$t = \frac{1-2r}{1-r^2}, \quad z = \frac{r^2-r+1}{1-r^2},$$

$$L = A_0(r) := \frac{r(r^2-r+1)}{1-r^2}, \quad z-L = B_0(r) := \frac{r^2-r+1}{1+r}.$$

The limiting normalized inside area and loss are

$$F_0(r) = \frac{(r^2-r+1)^2}{1-r^2}, \quad G_0(r) = 1 - F_0(r).$$

Since the exact reach A is at least the required demand $a \geq m$, (46) gives $A_0(r) > m$. Direct simplification yields

$$G_0(r) - \{2A_0(r) - 4A_0(r)^2\} = \frac{r^2(5r^4 - 8r^3 + 13r^2 - 8r + 2)}{(1-r^2)^2} \geq 0,$$

because the last polynomial is at least $9r^2 - 8r + 2 \geq 2/9$ on $[0, 1/2]$. Also $A_0(r) \leq r$, and

$$G_0(r) - \frac{1}{4} = \frac{(1-2r)(2r^3 - 3r^2 + 6r - 1)}{4(1-r^2)} \geq 0 \quad \text{when } A_0(r) \geq \frac{1}{4};$$

the cubic is increasing and has value $11/32$ at $r = 1/4$.

Finally $\psi(u) = 2u - 4u^2$ is increasing on $[0, 1/4]$ and never exceeds $1/4$ on $[0, 1/2]$. If $A_0(r) \leq 1/4$, then $G_0(r) \geq \psi(A_0(r)) \geq \psi(m)$; if $A_0(r) \geq 1/4$, then $G_0(r) \geq 1/4 \geq \psi(m)$. Together with (47), this proves (45). The reflected crossing is identical. $\square$

5.3. The cyclic area certificates. The selected handoffs of Theorem 2.9 have the form

$$(a_i, b_i) = (1 - x_{i-1}, x_i), \quad 0 < x_i < 1,$$

with cyclic indices. Row i is selected supercritical exactly when $x_i > x_{i-1}$.

Lemma 5.4 (At least two selected ascents). *Suppose every selected pair is feasible and at least two rows are supercritical. Then*

$$\sum_{i=0}^5 f(1 - x_{i-1}, x_i) < \frac{99}{20} < 5.$$

Proof. Put $m = \min_i x_i$ and $M = \max_i x_i$. The reflection $y_i = 1 - x_{i-1}$ swaps each row's two coordinates and preserves the sum and the number of ascents. Hence assume $m \leq 1 - M$. Every row coordinate is at least m , so (37) gives loss at least m^2 .

Choose an ascent out of a minimum plateau, say $x_{p-1} = m < x_p$. Its row contains the coordinate $1 - m$ and (38) gives loss at least $(1 - m)^2$. At a second ascent $q \neq p$, one row coordinate is strictly larger than $1/2$, so its loss is strictly larger than $1/4$. The other four rows lose at least m^2 each. Therefore the total loss is strictly larger than

$$4m^2 + (1 - m)^2 + \frac{1}{4} = 5 \left(m - \frac{1}{5} \right)^2 + \frac{21}{20} \geq \frac{21}{20}.$$

Subtracting from six proves the assertion. $\square$

Lemma 5.5 (One ascent and a T3-like row). *Suppose there is exactly one selected supercritical row, all roles are Vd0 or T3-like, and at least one is T3-like. Then the total normalized loss of the six actual vertex triangles is strictly greater than one.*

Proof. Rotate so that the unique ascent is row 0. Then

$$x_0 > x_5, \quad x_0 \geq x_1 \geq \cdots \geq x_5.$$

Put $M = x_0$, $m = x_5$, and reflect as above so that $m \leq 1 - M$. Every row coordinate is at least m , and $0 < m \leq 1/2$. The supercritical row has pair $(1 - m, M)$, so its loss is at least $(1 - m)^2$. By (44), a T3-like row is not row 0; by Theorem 5.3 it loses at least $2m - 4m^2$. Each of the remaining four rows loses at least m^2 . The total is therefore at least

$$(1 - m)^2 + (2m - 4m^2) + 4m^2 = 1 + m^2 > 1.$$

$\square$

Proposition 5.6 (Branches closed by area). *The following branches are impossible:*

- (1) T_C is CE0, $N_+ \geq 2$, with arbitrary vertex-role types;
- (2) T_C is CE0, $N_+ = 1$, no role is Vd1 or Vd2, and at least one role is T3-like.

Proof. In the CE0 class, the center role has no positive-length boundary trace. If the six vertex roles missed a boundary point, openness of the center role would give such a trace. Hence the vertex roles cover ∂H , so Theorem 2.9 applies.

If $N_+ \geq 2$, its at-least-two clause selects handoffs with at least two supercritical rows. Lemma 5.4 bounds the six vertex triangles' total normalized inside area by

less than $99/20$. The center triangle contributes at most one, so the total is less than

$$\frac{99}{20} + 1 = \frac{119}{20} < 6,$$

which is insufficient to cover H .

If $N_+ = 1$, the exact-one clause preserves the unique actual supercritical index for every strict selection. The exhaustive vertex-type hypothesis gives the assumptions of Lemma 5.5; the six vertex triangles contribute less than five normalized area, and the center triangle contributes at most one. Again the sum is strictly below the normalized area six of H . $\square$

6. STRATEGY 4: DIRECT NINE-POINT FORCING

This section gives the geometric proof of the direct nine-point obstruction. The exact frontier calculations, moving-circle signs, adjacent-overlap inequalities, rational radial envelopes, and global Bernstein identities are proved in Appendix D. The body of the proof uses those verified inputs only through the forcing lemmas and the four-overlap proposition below.

Assume throughout that all six vertex roles are Vd0, that they cover ∂H , and that exactly one actual boundary row is supercritical. Rotate its index to 4. Lemma D.3 gives strict handoffs

$$(48) \quad x_3 < x_2 < x_1 < x_0 < x_5 < x_4,$$

and parameters

$$(49) \quad a = 1 - x_3, \quad b = x_4, \quad p = 1 - b, \quad q = 1 - a$$

with

$$(50) \quad 0 < a, b < 1, \quad a + b > 1, \quad a^2 + ab + b^2 < 1.$$

Every selected row has incoming demand at least p and outgoing demand at least q .

6.1. Direct forcing of the disk and three frontier points. Let

$$c_* = c_{\max}(p, q), \quad h = \frac{\sqrt{3}}{2}, \quad \eta = h(1 - c_*).$$

The six radial points

$$D_i = (1 - c_*)V_i, \quad 0 \leq i \leq 5,$$

are excluded from their matching roles by maximality, from the two adjacent roles by Vd0 locality, and from the other three roles by distance from their distinguished vertices. Lemma D.4 therefore gives

$$(51) \quad \mathcal{D}_\eta := \{X : \|X\| \leq \eta\} \subset U_C.$$

At the supercritical corner V_4 , the strict AB -frontier consists of two circle pieces and two line pieces. The two line pieces meet at Q_0 ; their first intersections with the two actual adjacent-handoff unit circles are Q_- and Q_+ . The exact moving-circle signs in Lemma D.5 exclude all three points from the two adjacent roles. Direct vertex-distance inequalities exclude the remaining three nondistinguished roles, and exact frontier membership excludes the supercritical role. Lemma D.6 therefore gives

$$(52) \quad Q_-, Q_0, Q_+ \in U_C.$$

Consequently the center role contains the compact witness set

$$(53) \quad K_{\text{wit}} = \mathcal{D}_\eta \cup \{Q_-, Q_0, Q_+\}.$$

6.2. Newton inner points. The exact outer points Q_- and Q_+ contain two independent quadratic radicals. We remove them without weakening the obstruction by taking one Newton step inward along each frontier line.

Put

$$\rho = a^2 + ab + b^2, \quad D = \sqrt{4\rho - 3},$$

and define the V_4 -corner frontier coefficients

$$\begin{aligned} \alpha &= \frac{h(b + 2a - bD)}{2\rho}, & \beta &= \frac{h(b - a + (a + b)D)}{2\rho}, \\ \gamma &= \frac{h(-b + a + (a + b)D)}{2\rho}, & \delta &= \frac{h(2b + a - aD)}{2\rho}. \end{aligned}$$

The two line pieces meet in the local point

$$Q_0^{\text{loc}} = \left(\frac{2b + a - aD}{D + 3}, \frac{b + 2a - bD}{D + 3} \right).$$

Normalize

$$\bar{\alpha} = \alpha/h, \quad \bar{\beta} = \beta/h, \quad \bar{\gamma} = \gamma/h, \quad \bar{\delta} = \delta/h,$$

and use the dimensionless line parameters

$$\xi = \lambda/h, \quad \zeta = \mu/h, \quad \xi_* = \zeta_* = \frac{8\rho}{3(D + 3)}.$$

Let g_- and g_+ be the rescaled quadratic restrictions of the two adjacent-handoff circles to the two frontier lines. Appendix D proves

$$g_-(\xi_*) > 0 > g'_-(\xi_*), \quad g_+(\zeta_*) > 0 > g'_+(\zeta_*).$$

Define

$$\hat{\xi}_- = \xi_* - \frac{g_-(\xi_*)}{g'_-(\xi_*)}, \quad \hat{\zeta}_+ = \zeta_* - \frac{g_+(\zeta_*)}{g'_+(\zeta_*)}.$$

For a strictly convex quadratic that is positive and decreasing before its first root, the tangent zero lies strictly between the base point and that root. Hence the local points

$$\begin{aligned} A^{\text{loc}} &= \left(b - \frac{3}{4}\bar{\beta}\hat{\xi}_-, \frac{3}{4}\bar{\alpha}\hat{\xi}_- \right), \\ B^{\text{loc}} &= Q_0^{\text{loc}}, \\ C^{\text{loc}} &= \left(\frac{3}{4}\bar{\delta}\hat{\zeta}_+, a - \frac{3}{4}\bar{\gamma}\hat{\zeta}_+ \right) \end{aligned}$$

satisfy

$$A^{\text{loc}} \in (Q_0^{\text{loc}}, Q_-^{\text{loc}}), \quad C^{\text{loc}} \in (Q_0^{\text{loc}}, Q_+^{\text{loc}}).$$

Convert a local point (u, v) to the plane by

$$V_4 + u(V_5 - V_4) + v(V_3 - V_4),$$

and denote the resulting points by A, B, C . Then

$$(54) \quad \hat{K} := \text{conv}(\mathcal{D}_\eta \cup \{A, B, C\}) \subseteq \text{conv}(K_{\text{wit}}), \quad \Lambda(K_{\text{wit}}) \geq \Lambda(\hat{K}).$$

All coordinates of A, B, C are rational functions of a, b, D .

6.3. The cap-chain lemma. For a compact set K , let

$$h_K(n) = \max_{X \in K} \langle X, n \rangle,$$

and let R denote rotation through $2\pi/3$. The least equilateral side containing K is

$$(55) \quad \Lambda(K) = \frac{1}{h} \min_{\|n\|=1} \sum_{j=0}^2 h_K(R^j n).$$

For a disk $\mathbb{D}(X, r)$ define its level- h cap

$$I(X, r) = \{n \in S^1 : \langle X, n \rangle + r \geq h\}.$$

Lemma 6.1 (Cyclic cap chain). *Let K be compact and put*

$$S = K + RK + R^2K.$$

Suppose S contains the full R -orbits of

$$\mathbb{D}(A, 2\eta), \quad \mathbb{D}(B, 2\eta), \quad \mathbb{D}(C, 2\eta), \quad \mathbb{D}(W, \eta), \quad W = C + RA.$$

Assume that the rays

$$A, B, C, W, RA$$

occur in this cyclic order, each corresponding cap is a connected arc of half-width less than $\pi/2$, and the four consecutive pairs

$$\begin{aligned} I(A, 2\eta) &\leftrightarrow I(B, 2\eta), & I(B, 2\eta) &\leftrightarrow I(C, 2\eta), \\ I(C, 2\eta) &\leftrightarrow I(W, \eta), & I(W, \eta) &\leftrightarrow I(RA, 2\eta) \end{aligned}$$

overlap across their intervening short sectors. Then $\Lambda(K) \geq 1$.

Proof. The four overlaps and the cyclic ray order make the five caps cover the sector from the ray of A to the ray of RA . Their R -orbits cover the unit circle. For every unit normal n , one of the displayed disks therefore has support at least h in direction n . Since all those disks lie in S ,

$$h_S(n) \geq h.$$

On the other hand,

$$h_S(n) = \sum_{j=0}^2 h_K(R^j n).$$

Equation (55) now gives $\Lambda(K) \geq 1$. □

6.4. The four verified overlaps. For $K = \widehat{K}$, the Minkowski sum in the cap-chain lemma contains the four required disk orbits: one witness point and two copies of the centered disk give the three radius- 2η orbits, while $C + RA$ and one disk copy give the radius- η orbit.

Proposition 6.2 (Four-overlap certificate). *Assume (50) and $c_* > 2/3$. The Newton points above satisfy*

$$\|A\|, \|C\| > \eta,$$

the rays A, B, C, W, RA occur in that cyclic order, and all four consecutive cap overlaps in Lemma 6.1 hold.

Proof. The exact ray-order, determinant, and radius inequalities are proved in Appendix D. The two equal-radius overlaps are the analytic adjacent-line inequalities there. The two mixed overlaps are reduced to rational radial envelopes and exact Gram residuals in Subsection D.6; the resulting eight polynomial signs are proved by the twenty global Bernstein identities recorded in that appendix. These statements are precisely the four overlaps asserted here. $\square$

6.5. Enclosure and the center-independent contradiction.

Theorem 6.3 (Reader-facing witness enclosure). *For every pair (a, b) satisfying (50),*

$$\Lambda(K_{\text{wit}}(a, b)) \geq 1.$$

Proof. The disk alone gives

$$\Lambda(K_{\text{wit}}) \geq \frac{3\eta}{h} = 3(1 - c_*),$$

so the result is immediate when $c_* \leq 2/3$. Assume $c_* > 2/3$. The Newton reduction (54), Proposition 6.2, and Lemma 6.1 give

$$\Lambda(K_{\text{wit}}) \geq \Lambda(\hat{K}) \geq 1.$$

$\square$

Theorem 6.4 (Center-independent direct obstruction). *There is no cover of H by open unit equilateral triangles $U_C, U_0, \dots, U_5$ such that every U_i is a Vd0 role, the six vertex roles cover ∂H , and exactly one actual boundary row is supercritical.*

Proof. Direct forcing gives the compact containment $K_{\text{wit}} \subset U_C$, whereas Theorem 6.3 gives $\Lambda(K_{\text{wit}}) \geq 1$. A compact subset of an open unit equilateral triangle has positive clearance from all three sides and is therefore contained in a closed equilateral triangle of side strictly below one. This would give $\Lambda(K_{\text{wit}}) < 1$, a contradiction. $\square$

Proposition 6.5 (Branches closed by Strategy 4). *Theorem 6.4 closes*

- (1) *the CE0, $N_+ = 1$, all-Vd0 branch;*
- (2) *the zero-gap CE1 and CE2, $N_+ = 1$, all-Vd0 branches.*

Proof. In the CE0 branch, a boundary point missed by the six vertex roles would belong to the open center role and create a positive center trace, contrary to CE0. Thus the vertex roles cover ∂H . In the CE1 and CE2 branches, zero gaps mean exactly that the adjacent vertex traces cover every boundary edge. The exact-one hypothesis supplies the unique supercritical row in all three cases, so Theorem 6.4 applies. $\square$

7. EXHAUSTIVE ASSEMBLY

We now follow Table 1. Its center split is exhaustive by Proposition 2.3; its vertex refinements are exhaustive by Proposition 2.5; and the gap counts are exhaustive by Lemma 2.7. Every use of N_+ refers to the actual reaches (A_i, B_i) , while Proposition 2.9 supplies the smaller demands used by Strategies 3 and 4.

Proof of Theorem 1.1. Assume that seven open unit equilateral triangles cover H and label their distinct center and vertex roles as in Lemma 2.2.

CE0 with $N_+ = 0$ or $N_+ \geq 2$. The first state contradicts the terminal length theorem, Proposition 3.18; the second contradicts the terminal area theorem, Proposition 5.6.

CE0 with $N_+ = 1$. The exhaustive vertex trichotomy is: all roles Vd0; at least one Vd1/Vd2 role; or no Vd1/Vd2 role and at least one T3-like role. These are excluded, respectively, by Propositions 6.5, 3.18, and 5.6.

CE1/CE2 with $N_+ = 0$ or $N_+ \geq 2$. For $N_+ = 0$, the all-Vd0 and T3-like-without-Vd1/Vd2 refinements contradict Proposition 4.5, while the Vd1/Vd2 refinement contradicts Proposition 3.18. For $N_+ \geq 2$, the conditional skeleton part of Proposition 3.18 applies.

CE1/CE2 with $N_+ = 1$ and all roles Vd0. Lemma 2.7 gives zero or one gap for CE1 and zero, one, or two gaps for CE2. Zero gaps contradict Proposition 6.5; every positive-gap state contradicts Proposition 4.5.

CE1/CE2 with $N_+ = 1$ and mixed vertex types. A Vd1/Vd2 role in CE1, or at least two such roles in CE2, is excluded by Proposition 3.18. The remaining CE2 state has exactly one Vd1/Vd2 role: Strategy 1 closes the neighboring-midpoint and additional-positive-support subcases, and Strategy 2 closes the complementary exact placements. If no Vd1/Vd2 role occurs, then the non-Vd0 roles are precisely T3-like. Exactly one is excluded by Proposition 4.5, and at least two by Proposition 3.18.

Every row of Table 1 is impossible. Hence the assumed open cover does not exist. $\square$

Corollary 1.2 follows from Proposition 2.1.

APPENDIX A. EXACT LOCAL DEMAND CALCULUS

This section develops the local calculus for configurations in which a coarse length sum does not by itself record enough information. The mechanism is always the same. A center trace fixes boundary inputs and complementary radial demands; an exact local map propagates these data through nonsupercritical rows; and the returning demand exceeds the remaining boundary or radial capacity.

Actual maximal reaches continue to be denoted by (A_i, B_i, C_i) , while lowercase letters denote prescribed demands. Every estimate in this section is proved analytically. In particular, no empirical branch catalogue, machine-assisted certificate, or formal root from a discarded algebraic component is used.

A.1. The exact local feasible set. Put

$$u = \left(\frac{1}{2}, \frac{\sqrt{3}}{2} \right), \quad v = \left(\frac{1}{2}, -\frac{\sqrt{3}}{2} \right).$$

For $0 \leq a, b, c \leq 1$ define

$$K(a, b, c) = \text{conv}\{0, au, bv, c(u+v)\}.$$

Thus a, b are lower demands on the two incident boundary rays and c is a lower demand on the inward radial ray.

Definition A.1. The local admissible set is

$$\mathcal{A} = \{(a, b, c) \in [0, 1]^3 : K(a, b, c) \text{ is contained} \\ \text{in a closed unit equilateral triangle}\}.$$

For $a, c \in [0, 1]$ put

$$\begin{aligned} B_c(a) &= \max\{b : (a, b, c) \in \mathcal{A}\}, \\ F_c(a) &= \min\{B_c(a), 1 - a\}, \\ G_c(a) &= 1 - F_c(a). \end{aligned}$$

The distinction between demands and actual reaches is important. If a role has actual reaches (A, B, C) and $A \geq a$, $C \geq c$, then (a, B, c) is a demand triple realized by the same role. If in addition $A + B \leq 1$, then

$$(56) \quad B \leq F_c(a).$$

No assertion about the type of a maximizing triangle is required.

Proposition A.2 (Exact admissible set and radial envelope). *Put*

$$s = a + b, \quad d = a^2 + ab + b^2, \quad m = \min\{a, b\}, \quad M = \max\{a, b\},$$

and

$$q = s^4 - s^2 + ab.$$

Define

$$\begin{aligned} F_L(a, b, c) &= c^4 - c^2 + mc - m^2, \\ F_T(a, b, c) &= (s^2 - 1)c^2 + Mc - M^2, \\ F_S(a, b, c) &= (m^2 - 1)c^2 + (2mM^2 + M)c + M^4 - M^2. \end{aligned}$$

Then $\mathcal{A} = \mathcal{A}_L \cup \mathcal{A}_T \cup \mathcal{A}_S$, where

$$\begin{aligned} \mathcal{A}_L &= \{d \leq 1, s \leq 1, q \leq 0, F_L \leq 0\}, \\ \mathcal{A}_T &= \{d \leq 1, s \leq 1, q \geq 0, F_T \leq 0, c \leq 2M\}, \\ \mathcal{A}_S &= \{d \leq 1, s > 1, F_S \leq 0, c \leq \tfrac{1}{2}\}. \end{aligned}$$

All variables in these three sets are also constrained to $[0, 1]$.

For $d \leq 1$, the maximal radial demand is

$$(57) \quad c_{\max}(a, b) = \begin{cases} 1, & a = b = 0, \\ C_L(m), & s \leq 1, q \leq 0, (a, b) \neq (0, 0), \\ C_T(m, M), & s \leq 1, q > 0, \\ C_S(m, M), & s > 1, \end{cases}$$

where $C_L(m)$ is the unique root in $[\sqrt{3}/2, 1]$ of

$$z^4 - z^2 + mz - m^2 = 0,$$

and

$$C_T(m, M) = \frac{2M}{1 + \sqrt{4s^2 - 3}}, \quad C_S(m, M) = \frac{M(2mM + 1) - M\sqrt{4d - 3}}{2(1 - m^2)}.$$

If $d > 1$, the radial envelope is undefined. At $q = 0$ the two subcritical formulas agree and equal s .

Proof. If $s > 0$ and $cs \leq ab$, then $c/a + c/b \leq 1$ (with the zero-coordinate cases obtained by continuity), so $c(u+v)$ lies in $\text{conv}\{0, au, bv\}$. Its minimum enclosing equilateral side is the diameter

$$\sqrt{a^2 + ab + b^2}.$$

Assume now that $cs > ab$. The four points occur cyclically as $0, bv, c(u+v), au$. For three outward unit normals separated by 120° , the required side equals $2/\sqrt{3}$ times the sum of the three support values. On an angular cell with fixed support points this sum has second derivative equal to its negative. Hence an angular minimum occurs when a normal is perpendicular to one of the four hull edges. The four resulting side lengths are

$$\begin{aligned} L_{0A} &= b + \max\{a, c\}, & L_{B0} &= a + \max\{b, c\}, \\ L_{AC} &= \begin{cases} (a^2 + bc)/\sqrt{a^2 - ac + c^2}, & a \geq c, \\ c(a+b)/\sqrt{a^2 - ac + c^2}, & a < c \leq a+b, \\ c^2/\sqrt{a^2 - ac + c^2}, & c > a+b, \end{cases} \end{aligned}$$

and L_{CB} is obtained by interchanging a, b . Thus the minimum side is the minimum of these four expressions.

Ordering a, b as $m \leq M$, squaring the positive support equations gives $F_L = 0, F_T = 0, F_S = 0$ in the three respective contact regimes; the inactive support inequalities give $q \leq 0, q \geq 0, s > 1$. Squaring creates one spurious component in each of the last two cells. In the T cell the two roots are

$$\frac{2M}{1 + \sqrt{4s^2 - 3}}, \quad \frac{2M}{1 - \sqrt{4s^2 - 3}},$$

and $c \leq 2M$ selects the first. In the S cell the smaller root lies below $1/2$ and the other above $1/2$; hence $c \leq 1/2$ is exactly the geometric selector. The L fiber is the interval from zero to its unique selected root. Solving the selected equations gives (57). This also proves that $\mathcal{A}$ is coordinatewise down-closed. $\square$

A.2. The exact capped map. For $1/2 < c < 1$ put

$$\ell(c) = \frac{c}{2} \left(1 - \sqrt{4c^2 - 3} \right), \quad r(c) = c - \ell(c)$$

when $c \geq \sqrt{3}/2$, and

$$h(c) = \frac{2c}{1 + \sqrt{16c^2 - 3}}.$$

Also define

$$\begin{aligned} T_-(a, c) &= \frac{\sqrt{a^2 - ac + c^2}}{c} - a, \\ T_+(a, c) &= \frac{2(1 - a^2)c^2}{2ac^2 + c + \sqrt{(2ac^2 + c)^2 - 4(1 - c^2)(1 - a^2)c^2}}. \end{aligned}$$

Proposition A.3 (Four-label capped map). *For $1/2 < c \leq \sqrt{3}/2$,*

$$(58) \quad F_c(a) = \begin{cases} 1-a, & 0 \leq a \leq 1-c, \\ T_+(a, c), & 1-c < a < h(c), \\ h(c), & a = h(c), \\ T_-(a, c), & h(c) < a < c, \\ 1-a, & c \leq a \leq 1. \end{cases}$$

For $\sqrt{3}/2 < c < 1$,

$$(59) \quad F_c(a) = \begin{cases} 1-a, & 0 \leq a \leq 1-c, \\ T_+(a, c), & 1-c < a \leq \ell(c), \\ \ell(c), & \ell(c) < a < r(c), \\ T_-(a, c), & r(c) \leq a < c, \\ 1-a, & c \leq a \leq 1. \end{cases}$$

Thus there are exactly four labels

$$\text{Full}, \quad L, \quad T_-, \quad T_+^{\text{hi}}.$$

At $a = \ell(c)$ in (59), $F_c(a) = r(c)$; immediately to its right the value is $\ell(c)$. This downward jump is genuine. The formal other root of the T_+ quadratic is not feasible because it violates $c \leq 2 \max\{a, F_c(a)\}$.

The maps F_c are nonincreasing, the maps $G_c = 1 - F_c$ are nondecreasing and extensive, and

$$(60) \quad G_c(a) \leq z \iff G_c(1-z) \leq 1-a.$$

Proof. The cap $a + b \leq 1$ eliminates the S cell of Proposition A.2. On the cap $b = 1 - a$, the T inequality reduces to $M(c - M) \leq 0$; hence the Full face is feasible exactly for $a \leq 1 - c$ or $a \geq c$. Off that face a maximal point lies on $F_L = 0$ or $F_T = 0$. The L equation has roots $\ell(c), r(c)$ and the selector $q \leq 0$ keeps only the smaller coordinate. The ordered T halves give respectively the displayed lower quadratic root T_+ and the root T_- . Their order boundary is $a = b = h(c)$, while their $q = 0$ boundaries are $(\ell(c), r(c))$ and $(r(c), \ell(c))$. These observations give (58)–(59), including all equality assignments.

Down-closedness gives monotonicity, while $F_c(a) \leq 1 - a$ gives $G_c(a) \geq a$. Finally,

$$G_c(a) \leq z \iff F_c(a) \geq 1 - z \iff (a, 1 - z, c) \in \mathcal{A}, \quad a + 1 - z \leq 1.$$

Reflection $a \leftrightarrow b$ gives (60). □

Put

$$\mu = \frac{2\sqrt{3} - 3}{4},$$

and, for $0 < X < 1 - \sqrt{3}/2$, write

$$e(X) = \ell(1 - X).$$

The root equation gives

$$(61) \quad \begin{aligned} X &< e(X) && \left(0 < X < 1 - \frac{\sqrt{3}}{2}\right), \\ e(X) &< \frac{2X}{1-2X} && (0 < X \leq \mu), \\ e(X) &\leq 2X + 5X^2 && (0 < X \leq 1/8). \end{aligned}$$

Moreover,

$$(62) \quad a > e(X) \implies G_{1-X}(a) \geq 1 - e(X) \quad \left(0 < X < 1 - \frac{\sqrt{3}}{2}\right).$$

Indeed, for

$$p_X(z) = z^2 - (1-X)z + (1-X)^2 - (1-X)^4,$$

one has

$$p_X(X) = X(1 - 3X + 4X^2 - X^3) > 0.$$

Since $X < (1-X)/2$, this puts X strictly below the smaller root $e(X)$. For $U = 2X/(1-2X)$, direct substitution gives

$$p_X(U) = -\frac{X^2}{(1-2X)^2}(4X^4 - 20X^3 + 37X^2 - 28X + 3).$$

The polynomial in parentheses is decreasing on $[0, \mu]$ and at the right endpoint equals $8217/64 - 74\sqrt{3} > 0$. Indeed, its derivative is at most $16/512 + 74/8 - 28 < 0$, and $(8217/64)^2 - 3 \cdot 74^2 = 230001/4096 > 0$. Hence $p_X(U) < 0$, which puts U strictly between the roots. Finally,

$$p_X(2X + 5X^2) = X^2(3X + 4)(8X - 1) \leq 0,$$

and $X < 2X + 5X^2 < (1-X)/2$ on the stated range. Thus the trial point lies between the two roots, proving the third comparison. The implication (62) is read directly from (59). We shall also use $e(X) < 2X + 3X^2$ for $0 < X \leq 1/10$, obtained from $p_X(2X + 3X^2) = X^2(8X^2 + 19X - 2) < 0$; here too the trial point lies strictly between X and $(1-X)/2$.

A.3. Exact center traces and complementary demands. The normalization in Proposition 2.11 lets us assume that the unique center midpoint is M_0 .

Proposition A.4 (Exact CE1 and CE2 formulas). *In the CE1 case there are $0 < \lambda < 1$, $0 \leq s < t \leq 1$, and $\rho = \sqrt{1 - \lambda + \lambda^2}$ for which*

$$T_C \cap e_{0,1} = [s, t]$$

and

$$C_0 = \rho + \lambda s - t - \lambda, \quad C_1 = 1 - \lambda s, \quad C_2 = t + \lambda - 1.$$

The exact domain is

$$\begin{aligned} C_0 > 0, \quad C_2 > 0, \quad \lambda s \leq \frac{1}{2}, \quad t < 1 - \frac{\lambda}{2}, \\ t > \rho + \lambda s - \frac{1 + \lambda}{2}, \quad t \geq \rho - \frac{\lambda^2}{1 - \lambda} s. \end{aligned}$$

The center exits on the six arms are

$$\begin{aligned} d_0 &= C_1, & d_1 &= C_2/\lambda, \\ d_2 &= C_2, & d_3 &= \min\{C_0/\lambda, C_2/(1-\lambda)\}, \\ d_4 &= C_0, & d_5 &= C_0/(1-\lambda). \end{aligned}$$

In the CE2 case write

$$T_C \cap e_{5,0} = [x, u], \quad T_C \cap e_{0,1} = [y, v], \quad S = x + y, \quad D = \sqrt{x^2 + xy + y^2}.$$

Then

$$\begin{aligned} 0 &< x < u < 1, & 0 &< y < v < 1, \\ (u+v)S - xy &= D, & uS &\geq x, \quad vS \geq y, \\ uS &< x + y/2, & vS &< y + x/2, \end{aligned}$$

where the two weak center inequalities are strict for closures of original open center roles. The six exits are

$$\begin{aligned} d_0 &= 1 - \frac{xy}{S}, & d_1 &= \frac{vS - y}{x}, & d_2 &= \frac{vS - y}{S}, \\ d_3 &= \min\left\{\frac{vS - y}{y}, \frac{uS - x}{x}\right\}, & d_4 &= \frac{uS - x}{S}, & d_5 &= \frac{uS - x}{y}. \end{aligned}$$

In either case the complementary demand on row i is $c_i = 1 - d_i$; $c_0 \leq 1/2$ and $c_i > 1/2$ for $i \neq 0$.

Proof. In the affine coordinates $X = V_0 + b(V_1 - V_0) + a(V_5 - V_0)$, the CE1 triangle is cut out by

$$\begin{aligned} F_1 &= \lambda b + (1 - \lambda)a - \lambda s, \\ F_2 &= -b + \lambda a + t, \\ F_0 &= (1 - \lambda)b - a + \rho + \lambda s - t. \end{aligned}$$

Substitution of $O, M_0, \dots, M_5$ and of the six radial parametrizations gives the displayed domain and exits. The last domain inequality is exactly the absence of a positive interval on $e_{5,0}$.

For CE2, the side through the initial endpoints has outward normal $(\sqrt{3}S, y - x)/(2D)$. Rotating it by $\pm 120^\circ$, the three support constants are

$$\frac{\sqrt{3}(S - xy)}{2D}, \quad \frac{\sqrt{3}(vS - y)}{2D}, \quad \frac{\sqrt{3}(uS - x)}{2D}.$$

Their sum is $\sqrt{3}/2$ exactly when $(u + v)S - xy = D$. Substitution of the midpoints gives the four remaining inequalities, and substitution of the radial rays gives the six exits. The inequalities $d_0 \geq 1/2 > d_i$ for $i \neq 0$ are precisely the exact-midpoint conditions. $\square$

APPENDIX B. CYCLIC PROPAGATION IN THE ALL-Vd0 CASES

B.1. The exactly-one-supercritical all-Vd0 branch. For a center interval, call the nonempty closed set missed by its two open endpoint roles a *V-gap*. It may be a singleton. If the maximal closed center trace is $[s, t]$, open coverage gives the weak bounds $B_i \geq s$ and $A_{i+1} \geq 1 - t$; thus none of the arguments below discards a singleton gap.

Proposition B.1 (CE1 one-gap five-map obstruction). *Assume that every vertex role is Vd0, $N_+ = 1$, and the center role is CE1. If its boundary trace contains a V-gap, the roles do not cover H .*

Proof. By midpoint locality, rows $1, \dots, 5$ contain their own radial midpoints; hence they are nonsupercritical and row 0 is the unique supercritical row. Put

$$R = \lambda, \quad w = 1 - R, \quad E = \sqrt{1 - Rw}, \quad \eta = 1 - E, \quad P = \eta E.$$

Let A and D denote the center exits called C_0 and C_2 , respectively, in Proposition A.4; they are not the actual vertex-role radial reaches denoted by C_i . Then, with $X = R - D$, $m = A/R$ and $H = (\eta + A + D)/(2R) = s/2$, the exact strict domain is

$$(63) \quad \begin{aligned} &A > 0, \quad D > 0, \quad A + wD < P, \quad D + RA \geq P, \\ &A < \frac{w}{2}, \quad D < \frac{R}{2}, \quad RD - wA > 0, \end{aligned}$$

and the six complementary demands are

$$c_0 = \eta + A + D, \quad c_1 = 1 - \frac{D}{R}, \quad c_2 = 1 - D, \quad c_3 = 1 - m, \quad c_4 = 1 - A, \quad c_5 = 1 - \frac{A}{w}.$$

The gap gives $B_0 \geq s$ and $A_1 \geq X$. Define

$$z_0 = X, \quad z_j = G_{c_j}(z_{j-1}) \quad (1 \leq j \leq 5).$$

Radial coverage gives row j actual radial reach at least c_j , and rows $1, \dots, 5$ are nonsupercritical. Hence (56) and the monotonicity of F_{c_j} apply at each step. There is no V-gap on the next edge: for $j = 1, \dots, 4$ that edge has no center trace, and for $j = 5$ the CE1 center has no trace on $e_{5,0}$. The five actual-row transitions are therefore

$$\begin{aligned} A_1 &\geq z_0, & B_1 &\leq F_{c_1}(z_0), & A_2 &\geq 1 - B_1 \geq G_{c_1}(z_0) = z_1, \\ A_2 &\geq z_1, & B_2 &\leq F_{c_2}(z_1), & A_3 &\geq 1 - B_2 \geq G_{c_2}(z_1) = z_2, \\ A_3 &\geq z_2, & B_3 &\leq F_{c_3}(z_2), & A_4 &\geq 1 - B_3 \geq G_{c_3}(z_2) = z_3, \\ A_4 &\geq z_3, & B_4 &\leq F_{c_4}(z_3), & A_5 &\geq 1 - B_4 \geq G_{c_4}(z_3) = z_4, \\ A_5 &\geq z_4, & B_5 &\leq F_{c_5}(z_4), & A_0 &\geq 1 - B_5 \geq G_{c_5}(z_4) = z_5. \end{aligned}$$

Thus

$$(64) \quad A_0 \geq z_5 =: Z = (G_{c_5} \circ G_{c_4} \circ G_{c_3} \circ G_{c_2} \circ G_{c_1})(X).$$

On row 0, the inequalities $B_0 \geq s$ and actual radial reach at least c_0 , followed by reflection and down-closedness of $\mathcal{A}$, show that $(s, A_0, c_0) \in \mathcal{A}$. Therefore

$$(65) \quad A_0 \leq B_{c_0}(s).$$

It remains to prove $Z > B_{c_0}(s)$. Since G_{c_1} and G_{c_5} are extensive and the middle maps are nondecreasing, it is enough first to prove

$$(G_{c_4} \circ G_{c_3} \circ G_{c_2})(X) > 1 - H.$$

The negation of this inequality is, by three explicit applications of (60), equivalent to

$$\begin{aligned} (G_{c_4} \circ G_{c_3} \circ G_{c_2})(X) &\leq 1 - H \\ \iff G_{c_4}(H) &\leq 1 - (G_{c_3} \circ G_{c_2})(X) \\ \iff (G_{c_3} \circ G_{c_4})(H) &\leq 1 - G_{c_2}(X) \\ \iff (G_{c_2} \circ G_{c_3} \circ G_{c_4})(H) &\leq 1 - X. \end{aligned}$$

We shall prove the reverse strict inequality

$$(66) \quad (G_{c_2} \circ G_{c_3} \circ G_{c_4})(H) > 1 - X.$$

Here $A < P \leq \mu$, because $P = E(1 - E)$ and $E \geq \sqrt{3}/2$. We first prove the two input comparisons used at row 4. Since $D = R - X$, the inequality $A + wD < P$ and the identity $Rw = \eta + P$ give

$$A < wX - \eta.$$

Put

$$\varphi(q) = wq - \frac{q}{2(1+q)}.$$

The bound $0 < D < R/2$ gives $R/2 < X < R$, and φ is convex. At the endpoints,

$$\varphi(R/2) < \frac{Rw}{2} < \eta,$$

while $\varphi(R) < \eta$ is equivalent to

$$E(1 - 2R^2) < 1,$$

which follows from $E < 1$. Convexity therefore gives $\varphi(X) < \eta$, and hence

$$A < \frac{X}{2(1+X)}.$$

Using the rational low-root bound in its valid range,

$$(67) \quad e(A) < \frac{2A}{1-2A} < X.$$

We also have

$$2R(H - A) = \eta + D + (1 - 2R)A.$$

This is positive if $R \leq 1/2$. If $R > 1/2$, then $A < P = \eta E$ gives

$$2R(H - A) > \eta(1 - (2R - 1)E) > 0.$$

Thus $H > A$. Since $H = s/2$ and $0 < s < t \leq 1$, we also have $H < 1/2 < 1 - A$.

Now consult the exact high-radial catalog for $c_4 = 1 - A$. The lower and upper Full ranges would require $H \leq A$ and $H \geq 1 - A$, respectively, so both are impossible. On the L or T_- range,

$$F_{1-A}(H) \leq e(A),$$

and therefore

$$G_{1-A}(H) \geq 1 - e(A) > 1 - X$$

by (67). Thus only the T_+^{hi} range remains. Its selector gives

$$H \leq e(A) \leq 2A + 5A^2.$$

We claim that this range forces $X > 1/2$. Indeed, if $X \leq 1/2$, the selector and the two center inequalities give

$$Q(A) := \eta + P + wA - 2R(2A + 5A^2) \leq 0, \quad 0 < A < A_\star := P - w(R - \tfrac{1}{2}).$$

The quadratic Q is strictly concave and $Q(0) = \eta + P = Rw > 0$. Rationalize by

$$k = \frac{R}{1+E}, \quad R = \frac{k(2-k)}{1-k^2}, \quad E = \frac{1-k+k^2}{1-k^2}, \quad \eta = \frac{k(1-2k)}{1-k^2}.$$

When $R \leq 1/2$, one has $k \leq 2 - \sqrt{3} < 27/100$ and $A_\star \geq P$, while

$$Q(P) = \frac{k(1-2k)(-f_8(k))}{(1-k^2)^5},$$

where

$$f_8(k) = 16k^8 - 77k^7 + 175k^6 - 244k^5 + 199k^4 - 101k^3 + 13k^2 + 12k - 3.$$

Put $p = -f_8$. Direct differentiation and regrouping give

$$p'(k) = -12 + g_1(k) + g_2(k) + g_3(k),$$

where

$$g_1(k) = -k(4k-1)(199k-26),$$

$$g_2(k) = 10k^4(122-105k),$$

$$g_3(k) = k^6(539-128k).$$

The first term is positive only for $26/199 < k < 1/4$. On that interval,

$$g_1(k) \leq \frac{1}{4} \max_k (1-4k)(199k-26) = \frac{9025}{12736} < \frac{3}{4},$$

and the same upper bound is automatic when $g_1(k) \leq 0$. The functions g_2 and g_3 are increasing on $[0, 27/100]$, and direct rational evaluation gives

$$g_2\left(\frac{27}{100}\right) < 5, \quad g_3\left(\frac{27}{100}\right) < \frac{1}{5}.$$

Consequently

$$p'(k) < -12 + \frac{3}{4} + 5 + \frac{1}{5} < 0.$$

Since

$$p\left(\frac{27}{100}\right) = \frac{81581849832351}{2500000000000000} > 0,$$

one has $-f_8(k) = p(k) > 0$ throughout the required interval, and so $Q(P) > 0$. When $R > 1/2$, $0 < A_\star < P$ and

$$Q(A_\star) = \frac{(1-2k)(-g_5(k))}{2(1-k^2)^3}, \quad g_5(k) = 32k^5 - 118k^4 + 150k^3 - 86k^2 + 18k - 1.$$

At $k_0 = 2 - \sqrt{3}$, $g_5(k_0) = 3471 - 2004\sqrt{3} < 0$ because $3 \cdot 2004^2 - 3471^2 = 207$. If $y = 4k - 1 \in [0, 1]$, then

$$-g'_5(k) = \frac{29 - 5y^4 + y(39(y-1)^2 + 12)}{8} > 0.$$

Thus g_5 decreases and remains negative for $k > k_0 > 1/4$, so $Q(A_\star) > 0$. In either range concavity makes $Q(A) > 0$ throughout the allowed interval, contradicting $Q(A) \leq 0$ and proving $X > 1/2$.

Put $p_1 = G_{1-A}(H)$. If row 3 is not T_+^{hi} , then $p_1 \leq A + e(A) < 3A + 5A^2 < 29/64 < 1/2$, and $2R(H - m) = \eta + D - A > \eta - P > 0$. Extensivity gives $p_1 \geq H > m$, while $p_1 < 1/2 < 1 - m$; thus both Full ranges are impossible. On L , T_- , or the low- c tie one has $F_{1-m}(p_1) \leq p_1$, so

$$G_{1-m}(p_1) \geq 1 - p_1 > 1 - X.$$

Extensivity of the final reverse map G_{1-D} preserves this strict bound and proves (66). It remains to treat the selected row-3 T_+ branch. We use the following elementary property of a selected T_+ arc. Put $d = 1 - c$, let $q = G_c(p)$, and write $u = q - p$. The selected equation is

$$f(q) := (1 - q)(q - d) = g(u) := (1 - d)^2 u(2 - u).$$

On the interior of the arc, $f', g' > 0$. If

$$A = f'(q) = 1 + d - 2q, \quad B = g'(u) = 2(1 - d)^2(1 - u),$$

then

$$u' = \frac{A}{B}, \quad u'' = \frac{2((1 - d)^2 A^2 - B^2)}{B^3}.$$

Moreover

$$B - (1 - d)A = (1 - d)(1 - 3d + 2p + 2du) > 0$$

and

$$B - A \geq (1 - d)(1 - 2d) > 0.$$

Thus $p = q - u$ is increasing and strictly convex as a function of q . Consequently $q = G_{1-d}(p)$ is increasing and strictly concave on each selected arc; a tie endpoint follows by continuity.

When $d < 1 - \sqrt{3}/2$, the arc endpoints are (d, d) and $(e(d), d + e(d))$, so concavity gives

$$(68) \quad q \geq p + \frac{d}{e(d) - d}(p - d).$$

For $0 < d \leq \mu$, the valid rational low-root bound gives

$$\frac{d}{e(d) - d} > \frac{1 - 2d}{1 + 2d} > 1 - 4d > 1 - 5d.$$

This covers row 4, because its parameter is $d = A < P \leq \mu$. For the row-3 parameter $d = m$, which need not be at most μ , the remaining high-radial range is handled without the rational estimate. If $\mu < d \leq 1/8$, then

$$\frac{d}{e(d) - d} \geq \frac{1}{1 + 5d} > 1 - 5d$$

by $e(d) \leq 2d + 5d^2$. If $1/8 < d < 1 - \sqrt{3}/2$, then $e(d) < (1 - d)/2$ and hence

$$\frac{d}{e(d) - d} > \frac{2d}{1 - 3d} > 1 - 5d.$$

The last comparison is equivalent to $15d^2 - 10d + 1 < 0$; its smaller root $(5 - \sqrt{10})/15$ is less than $1/8$. In the low- c row-3 range $1 - \sqrt{3}/2 \leq d < 1/5$, the other endpoint is $(h(1 - d), 1 - h(1 - d))$, and the chord coefficient is

$$\frac{1 - 2h(1 - d)}{h(1 - d) - d}.$$

On $3/4 \leq c \leq \sqrt{3}/2$, direct differentiation of the formula for h shows that $h(c)$ is nonincreasing. Moreover,

$$h\left(\frac{3}{4}\right) = \frac{3}{2(\sqrt{6}+1)} < \frac{7}{16}.$$

Here low c gives $d \geq 1 - \sqrt{3}/2 > 2/15 > 1/16$, and hence

$$h(1-d) < \frac{7}{16} < \frac{3+d}{7}.$$

Thus the displayed chord coefficient is greater than $1/3$, while $1 - 5d < 1/3$. For $d \geq 1/5$, extensivity alone is stronger than the row-3 estimate.

Applying these chord bounds first with $d = A$ and then with $d = m$ proves, without an omitted selected branch,

$$\begin{aligned} p_1 &> L_1 := (2 - 4A)H - (1 - 4A)A, \\ p_2 &:= G_{1-m}(p_1) > L_2 := (2 - 5m)L_1 - (1 - 5m)m. \end{aligned}$$

Finally let

$$D_0 = P - RA, \quad D_h = A(4R - 1) + 10RA^2 - \eta, \quad x = Rw.$$

Feasibility puts $D_0 \leq D \leq D_h$. We first prove that the actual feasible value satisfies $D < 1/10$. Since

$$P = \frac{xE}{1+E} < \frac{x}{2}, \quad \eta = \frac{x}{1+E} > \frac{x}{2},$$

the contrary assumption $D \geq 1/10$ and $A + wD < P$ give

$$A < \bar{A} := \frac{w(5R - 1)}{10}.$$

The function $D_h(A)$ is increasing, so

$$D \leq D_h(A) < D_h(\bar{A}) < U(R) := \bar{A}(4R - 1) + 10R\bar{A}^2 - \frac{x}{2}.$$

Direct expansion yields

$$\frac{1}{10} - U(R) = -\frac{R}{10}P_4(R), \quad P_4(R) = 25R^4 - 60R^3 + 26R^2 + 22R - 14.$$

For $y = 2R - 1 \in [0, 1]$,

$$16P_4(R) = 25y^4 - 20y^3 - 106y^2 + 124y - 39 \leq -101y^2 + 124y - 39 < 0.$$

The last quadratic has discriminant -380 . Hence $U(R) < 1/10$, a contradiction, and therefore $D < 1/10$.

It remains to compare L_2 with $2D + 3D^2$. Put

$$J(D) = L_2(D) - \frac{23}{10}D.$$

Since

$$L_2 - 2D - 3D^2 - J(D) = 3D\left(\frac{1}{10} - D\right) > 0,$$

it is enough to prove $J(D) > 0$. Exact differentiation gives

$$J'(D) = \frac{S(R, A)}{10R^2}, \quad S(R, A) = 100A^2 - (40R + 50)A + R(20 - 23R).$$

The inequality $D_0 \leq D_h$ is equivalent to

$$x \leq A(5R - 1 + 10RA).$$

Because $A < P < x/2 \leq 1/8$,

$$A > L := \frac{4x}{25R - 4}.$$

Also $\partial_A S = 200A - 40R - 50 < -45$, and direct substitution gives

$$S(R, L) = -\frac{Rq_3(R)}{(25R - 4)^2},$$

where

$$q_3(R) = 8775R^3 - 14260R^2 + 7928R - 1120.$$

For $y = 2R - 1$,

$$8q_3(R) = 8775y^3 - 2195y^2 + 997y + 3007 \geq 812 > 0.$$

Thus $J'(D) < 0$ and $J(D) \geq J(D_h)$.

At $D = D_h$ one has $H = 2A + 5A^2$. Write $L_{2,h} = L_2(D_h)$. Expansion gives

$$L_{2,h} - A(1 + 4R) = \frac{A}{R^2} \mathcal{Q}(R, A),$$

where

$$\begin{aligned} \mathcal{Q}(R, A) &= 100A^3R - 40A^2R^2 - 30A^2R + 12AR^2 - 15AR + 5A \\ &\quad - 4R^3 + 5R^2 - R. \end{aligned}$$

On $0 \leq A \leq x/2$,

$$\partial_A^2 \mathcal{Q} = 20R(30A - 4R - 3) < 0.$$

Its two endpoint values are

$$\mathcal{Q}(R, 0) = x(4R - 1) > 0, \quad \mathcal{Q}\left(R, \frac{x}{2}\right) = \frac{x}{2}p(R),$$

where

$$p(R) = 25R^5 - 30R^4 + 20R^3 - 3R^2 - 7R + 3.$$

For $y = 2R - 1$,

$$32p(R) = 25y^5 + 65y^4 + 90y^3 + 106y^2 - 35y + 5 > 0,$$

because the terms of degree at least three are nonnegative and $106y^2 - 35y + 5$ has discriminant -895 . Concavity therefore gives $L_{2,h} > A(1 + 4R)$.

On the other hand, $A < P = \eta E$ and $A < P < x/2$ imply

$$\frac{D_h}{A} = 4R - 1 + 10RA - \frac{\eta}{A} < C_0 := 4R - 2 + 5Rx - \frac{x}{2}.$$

Here we used

$$\frac{1}{E} > 1 + \frac{x}{2}, \quad (1 - x) \left(1 + \frac{x}{2}\right)^2 = 1 - \frac{x^2(3 + x)}{4} < 1.$$

Finally,

$$1 + 4R - \frac{23}{10}C_0 = \frac{h_3(R)}{20}, \quad h_3(R) = 230R^3 - 253R^2 - 81R + 112.$$

Since $h_3''(R) = 46(30R - 11) \geq 184$, strong convexity at $R_0 = 20/23$ gives

$$h_3(R) \geq \frac{788}{529} - \frac{(17/23)^2}{368} = \frac{289695}{194672} > 0.$$

Consequently

$$J(D_h) > A \left(1 + 4R - \frac{23}{10}C_0 \right) > 0,$$

and hence

$$p_2 > L_2 > 2D + 3D^2.$$

The low root $e(D)$ is the smaller root of

$$H_D(z) = z^2 - (1 - D)z + (1 - D)^2 - (1 - D)^4,$$

and direct substitution gives $H_D(2D + 3D^2) = D^2(8D^2 + 19D - 2) < 0$. Hence $e(D) < 2D + 3D^2 < p_2$. To check every row-2 label explicitly, put

$$e_D = e(D) = \ell(1 - D), \quad r_D = 1 - D - e_D.$$

Since $D < 1/10$, one has $c_2 = 1 - D > \sqrt{3}/2$, and the exact catalog is

input range	label	$F_{1-D}(p)$
$0 \leq p \leq D$	lower Full	$1 - p$
$D < p \leq e_D$	T_+	$T_+(p, 1 - D)$
$e_D < p < r_D$	L	e_D
$r_D \leq p < 1 - D$	T_-	$T_-(p, 1 - D)$
$1 - D \leq p \leq 1$	upper Full	$1 - p$

The inequality $p_2 > e_D$ excludes the first two ranges. On the L range $F_{1-D}(p_2) = e_D$; on the T_- range, monotonicity and $F_{1-D}(r_D) = e_D$ give $F_{1-D}(p_2) \leq e_D$. Hence in both middle ranges

$$G_{1-D}(p_2) \geq 1 - e_D > 1 - X,$$

because

$$e_D < 2D + 3D^2 < \frac{23}{100} < \frac{1}{2} < X.$$

Finally, on upper Full,

$$G_{1-D}(p_2) = p_2 \geq 1 - D > 1 - X$$

because $X > 1/2 > D$. Thus every row-2 label gives the strict inequality (66).

By the suffix reduction and duality established above,

$$Z > 1 - H = 1 - \frac{s}{2}.$$

For any admissible pair of boundary reaches s, b , the diameter constraint is $s^2 + sb + b^2 \leq 1$. Its positive endpoint

$$\beta(s) = \frac{-s + \sqrt{4 - 3s^2}}{2}$$

therefore satisfies $B_{c_0}(s) \leq \beta(s)$. Since $s > 0$, $\sqrt{4 - 3s^2} < 2$, and so $\beta(s) < 1 - s/2$. Thus $Z > B_{c_0}(s)$, contradicting (64) and (65). $\square$

Proposition B.2 (CE2 one-gap five-map obstruction). *Assume that every vertex role is Vd0, $N_+ = 1$, and the center role is CE2. If exactly one of its two boundary traces contains a V-gap, the roles do not cover H .*

Proof. Again row 0 is uniquely supercritical. Put

$$R = \frac{x}{x+y}, \quad w = 1 - R, \quad E = \sqrt{1 - Rw}, \quad \eta = 1 - E, \quad P = \eta E,$$

and

$$\alpha = u - R, \quad \delta = v - w, \quad k = \eta + \alpha + \delta.$$

The exact CE2 domain becomes

$$(69) \quad \alpha, \delta > 0, \quad \delta + R\alpha < P, \quad \alpha + w\delta < P,$$

with $x = k/w$, $y = k/R$, and demands

$$\begin{aligned} c_0 &= k, & c_1 &= 1 - \delta/R, \\ c_2 &= 1 - \delta, & c_3 &= 1 - \min\{\delta/w, \alpha/R\}, \\ c_4 &= 1 - \alpha, & c_5 &= 1 - \alpha/w. \end{aligned}$$

Write $F_i = F_{c_i}$ and $G_i = G_{c_i}$ for $1 \leq i \leq 5$. Since $E \geq \sqrt{3}/2$ and $E(1-E)$ decreases on this range,

$$(70) \quad 0 < \alpha, \delta < P \leq \mu < 1 - \frac{\sqrt{3}}{2}.$$

Thus all low-root estimates used below are within their stated ranges.

Suppose first that the gap is in $[y, v]$; then $[x, u]$ contains no V-gap. The endpoint bounds give

$$B_0 \geq y, \quad A_1 \geq z_0 := 1 - v = R - \delta.$$

Define all five successive map inputs before using them:

$$(71) \quad z_i = G_i(z_{i-1}) \quad (1 \leq i \leq 5), \quad Z_R = z_5 = (G_5 \circ G_4 \circ G_3 \circ G_2 \circ G_1)(1 - v).$$

Radial coverage, the nonsupercritical cap, and monotonicity bound each actual outgoing reach. For rows $1, \dots, 4$ the next edge has no center trace, while after row 5 the other center interval $[x, u]$ contains no V-gap. All five handoffs are therefore strict. The complete actual-row chain is

$$\begin{aligned} A_1 &\geq z_0, & B_1 &\leq F_{c_1}(z_0), & A_2 &> 1 - B_1 \geq G_1(z_0) = z_1, \\ A_2 &> z_1, & B_2 &\leq F_{c_2}(z_1), & A_3 &> 1 - B_2 \geq G_2(z_1) = z_2, \\ A_3 &> z_2, & B_3 &\leq F_{c_3}(z_2), & A_4 &> 1 - B_3 \geq G_3(z_2) = z_3, \\ A_4 &> z_3, & B_4 &\leq F_{c_4}(z_3), & A_5 &> 1 - B_4 \geq G_4(z_3) = z_4, \\ A_5 &> z_4, & B_5 &\leq F_{c_5}(z_4), & A_0 &> 1 - B_5 \geq G_5(z_4) = z_5. \end{aligned}$$

In particular,

$$(72) \quad A_0 > Z_R.$$

On the supercritical row, $B_0 \geq y$ and the actual radial reach is at least c_0 . Reflection and down-closedness of $\mathcal{A}$ therefore give

$$(73) \quad A_0 \leq B_{c_0}(y).$$

We next estimate the formal chain. The second inequality in (69) and the identity $Rw = \eta + P$ give

$$wz_0 = Rw - w\delta > \eta + \alpha.$$

The function

$$\pi(q) = (\eta + q)(1 - 2q) - 2qw$$

is concave, and direct simplification gives

$$\pi(0) = \eta > 0, \quad \pi(P) = \eta(\eta + 2ER^2) > 0.$$

Because $0 < \alpha < P$, concavity gives $\pi(\alpha) > 0$. Together with (61) and (70), this proves

$$(74) \quad z_0 > \frac{\eta + \alpha}{w} > \frac{2\alpha}{1 - 2\alpha} > e(\alpha).$$

Put $Q = y/2 = k/(2R)$. We also have

$$(75) \quad \min\{e(\alpha), e(\delta)\} < Q.$$

Suppose instead that both roots are at least Q . The strict rational low-root bounds then imply

$$\alpha > L, \quad \delta > L, \quad L = \frac{Q}{2(1 + Q)}.$$

Since $k = 2RQ = \eta + \alpha + \delta$, we obtain

$$\eta < Q \left(2R - \frac{1}{1 + Q} \right).$$

On the other hand, the first inequality in (69) gives

$$\begin{aligned} \eta E &> \delta + R\alpha = R(\alpha + \delta) + w\delta \\ &> R(2RQ - \eta) + wL, \end{aligned}$$

and hence

$$\eta(E + R) > Q \left(2R^2 + \frac{w}{2(1 + Q)} \right).$$

Combining the strict upper and lower bounds for η gives $H(Q) > 0$, where

$$H(q) = \left(2R - \frac{1}{1 + q} \right) (E + R) - 2R^2 - \frac{w}{2(1 + q)}.$$

But

$$H'(q) = \frac{E + R + w/2}{(1 + q)^2} > 0.$$

Moreover $x = k/w < 1$ gives $Q < w/(2R)$, and

$$H\left(\frac{w}{2R}\right) = \frac{R(2RE - R - 1)}{R + 1} < 0,$$

the last sign following from $(R + 1)^2 - 4R^2E^2 = (1 - R)(4R^3 + 3R + 1) > 0$. Thus $H(Q) < H(w/(2R)) < 0$, a contradiction. This proves (75).

Every G_i is extensive. If $e(\alpha) < Q$, then $z_3 \geq z_0 > e(\alpha)$, and $c_4 = 1 - \alpha$ together with (62) gives

$$z_4 \geq 1 - e(\alpha) > 1 - Q.$$

Thus $z_5 > 1 - Q$. Otherwise $e(\alpha) \geq Q$, so (75) and (74) give

$$z_1 \geq z_0 > e(\alpha) \geq Q > e(\delta).$$

Since $c_2 = 1 - \delta$, the threshold lemma gives

$$z_2 \geq 1 - e(\delta) > 1 - Q,$$

and rows 3, 4, 5 preserve this strict inequality. Hence in both cases

$$(76) \quad Z_R > 1 - Q = 1 - \frac{y}{2}.$$

Define again

$$\beta(q) = \frac{-q + \sqrt{4 - 3q^2}}{2}.$$

The universal diameter bound gives

$$B_{c_0}(y) \leq \beta(y) < 1 - \frac{y}{2}.$$

This contradicts (72) and (73).

It remains to record the other orientation explicitly. Write primes for the data after reflection across the axis through V_0 . The normalized center variables exchange as

$$(x, u, R, \alpha) \longleftrightarrow (y, v, w, \delta)$$

and the actual boundary reaches transform as

$$A'_0 = B_0, \quad B'_0 = A_0, \quad A'_i = B_{6-i}, \quad B'_i = A_{6-i} \quad (1 \leq i \leq 5).$$

The radial demands and capped maps transform in the exact order

$$c'_0 = c_0, \quad c'_i = c_{6-i}, \quad F'_i = F_{6-i}, \quad G'_i = G_{6-i} \quad (1 \leq i \leq 5).$$

Suppose now that the gap is in $[x, u]$, so $[y, v]$ contains no V-gap. Its endpoint bounds give

$$A_0 \geq x, \quad B_5 \geq \tilde{z}_0 := 1 - u.$$

Define the reflected chain by

$$\tilde{z}_1 = G_5(\tilde{z}_0), \quad \tilde{z}_2 = G_4(\tilde{z}_1), \quad \tilde{z}_3 = G_3(\tilde{z}_2), \quad \tilde{z}_4 = G_2(\tilde{z}_3), \quad \tilde{z}_5 = G_1(\tilde{z}_4).$$

Starting from the actual reach B_5 and traversing rows 5, 4, 3, 2, 1, the five reverse transitions are

$$\begin{array}{lll} B_5 \geq \tilde{z}_0, & A_5 \leq F_{c_5}(\tilde{z}_0), & B_4 > 1 - A_5 \geq G_5(\tilde{z}_0) = \tilde{z}_1, \\ B_4 > \tilde{z}_1, & A_4 \leq F_{c_4}(\tilde{z}_1), & B_3 > 1 - A_4 \geq G_4(\tilde{z}_1) = \tilde{z}_2, \\ B_3 > \tilde{z}_2, & A_3 \leq F_{c_3}(\tilde{z}_2), & B_2 > 1 - A_3 \geq G_3(\tilde{z}_2) = \tilde{z}_3, \\ B_2 > \tilde{z}_3, & A_2 \leq F_{c_2}(\tilde{z}_3), & B_1 > 1 - A_2 \geq G_2(\tilde{z}_3) = \tilde{z}_4, \\ B_1 > \tilde{z}_4, & A_1 \leq F_{c_1}(\tilde{z}_4), & B_0 > 1 - A_1 \geq G_1(\tilde{z}_4) = \tilde{z}_5. \end{array}$$

The last strict handoff is valid because the other center interval $[y, v]$ has no V-gap. In particular,

$$(77) \quad B_0 > Z_L := \tilde{z}_5 = (G_1 \circ G_2 \circ G_3 \circ G_4 \circ G_5)(1 - u).$$

Meanwhile the actual row-0 reaches satisfy $A_0 \geq x$ and $C_0 \geq c_0$. Coordinatewise down-closedness gives $(x, B_0, c_0) \in \mathcal{A}$, and hence

$$(78) \quad B_0 \leq B_{c_0}(x).$$

The normalized domain, initial-threshold proof, and two-threshold proof are invariant under the displayed exchange. Applying the right-chain argument to the reflected variables therefore gives

$$Z_L > 1 - \frac{x}{2} > B_{c_0}(x),$$

where the last inequality is the same diameter estimate with x in place of y . This contradicts (77) and (78), completing both orientations. $\square$

Proposition B.3 (Positive-gap all-Vd0, $N_+ = 1$). *If the center role is CE1 or CE2, every vertex role is Vd0, and $N_+ = 1$, then no cover of H can have a boundary gap in the actual open vertex-role traces.*

Proof. The midpoint argument used above makes T_0 uniquely supercritical. In an assumed cover, every boundary gap in the vertex-role traces must lie in a center trace. CE1 therefore has the one-gap state excluded by Proposition B.1. For CE2, the exactly-one-gap state is Proposition B.2; if both center intervals contain gaps, the endpoint bounds and exact exits are those of Lemma C.8, so the two far outgoing reaches sum to less than one. Rows 2, 3, 4 would then have to cover more than three boundary units, contradicting their three nonsupercritical caps. This exhausts the one- and two-gap states, including singleton gaps. The zero-gap state is intentionally owned by the center-independent obstruction of Strategy 4. $\square$

APPENDIX C. NONSUPERCRITICAL, T3-LIKE, AND VD1/VD2 DEMAND BRANCHES

C.1. The exactly-one-T3 branch.

Proposition C.1 (Exactly one T3-like row). *Assume the center role is CE1 or CE2, $N_+ = 1$, exactly one vertex role is T3-like, and no role is Vd1 or Vd2. Then the roles do not cover H .*

Proof. Normalize the center midpoint to M_0 . A T3-like row misses its own midpoint, a supercritical row misses all its local midpoints, and a Vd0 row cannot rescue a neighboring midpoint. Consequently, after reflection,

$$T_0 \text{ is T3-like, } M_1 \in T_0, \quad T_1 \text{ is uniquely supercritical,}$$

and $T_2, \dots, T_5$ are nonsupercritical Vd0 rows.

In V_0 wedge coordinates put

$$1 \leq D \leq \frac{2}{\sqrt{3}}, \quad E = \sqrt{4 - 3D^2}, \quad R = \frac{D + E}{2},$$

and let a be the $e_{5,0}$ reach of T_0 . Its interval on r_1 is

$$[c, u], \quad c = \frac{D(1 + a) - 1}{R}, \quad u = 1 - \frac{R}{D} + a,$$

where $E/(2D) \leq a \leq (4 - 3D + E)/(4D)$. Its own-radial exit is $x_0 = aD/R$. For $B(c) = \mathcal{B}(c)$ and $A(c) = 1 - B(c)$, direct substitution gives

$$c - \frac{x_0(2 - x_0)}{1 + x_0} = \frac{2N(D, a)}{(D + E)(2Da + D + E)},$$

where

$$N = 4D^2a^2 + D^2a + D^2 - DEa + DE - 2Da - D - E.$$

This convex quadratic is nonnegative: on $1 \leq D \leq 8/7$ its minimum on the allowed interval is at $a = E/(2D)$, and squaring reduces the sign to $4(D - 1)(7D^3 - 10D^2 - 8D + 12) \geq 0$; on $8/7 \leq D \leq 2/\sqrt{3}$ its critical value is positive because $(9D - 10)E + 9D^2 - 6D - 4 > 0$. Before using monotonicity, note explicitly that

$$x_0 = \frac{aD}{R} \leq \frac{4 - 3D + E}{4R} \leq \frac{1}{2};$$

the second inequality is equivalent, using $2R = D + E$, to $D \geq 1$. Also $c \leq 1/2$ because $M_1 \in [c, u]$, and hence $0 \leq A(c) = 1 - B(c) \leq 1/2$. Since $X(2 - X)/(1 + X)$ is increasing on $[0, 1/2]$, this proves

$$(79) \quad x_0 \leq A(c).$$

Radial coverage forces the supercritical row's own demand $c_1 \geq c$. Lemma C.6 gives $b_1 < B(c_1) \leq B(c)$. On $e_{5,0}$ the far row T_5 is forced to reach at least $h \geq B(c)$. Indeed this is $h \geq 1 - a$ unless a CE2 interval hides the boundary gap. In the hiding case the side model gives $T \leq x_0$, and (79) again gives $h \geq 1 - T \geq B(c)$. Therefore

$$\sum_{i=2}^5 (a_i + b_i) \geq (1 - b_1) + 1 + 1 + 1 + h > 4,$$

contrary to the four row caps. The case $c = 1/2$ is already impossible because the strict supercritical feasible set is empty there. $\square$

C.2. The nonsupercritical CE1/CE2 boundary packages. Figure 7 gives a common schematic for the package proved below.




FIGURE 7. Schematic view of the common CE1/CE2, $N_+ = 0$, all-Vd0 boundary package, not to scale. The center role and three representative Vd0 vertex roles are shown; the remaining three roles are omitted for readability. The simultaneous placement is not a candidate cover, and no proof depends on visual inspection.

Proposition C.2 (All-Vd0 boundary loss). *Assume that T_C is CE1 or CE2, every vertex role is Vd0, and*

$$A_i + B_i \leq 1 \quad (0 \leq i \leq 5).$$

Then the seven roles do not cover H .

Proof. If the center boundary traces are already covered by the open vertex roles, then those six roles cover all of ∂H . Put $W_i = U_i \cap \partial H$. These are relatively open subsets of the connected circle ∂H , each of length at most one. At least two are nonempty, and a finite relatively open cover of a connected space cannot have its nonempty members pairwise disjoint. Thus two overlap on a positive-length relatively open set, whence

$$6 = \mathcal{H}^1(\partial H) < \sum_{i=0}^5 \mathcal{H}^1(W_i) \leq \sum_{i=0}^5 (A_i + B_i) \leq 6,$$

a contradiction.

Suppose next that T_C is CE1 and its trace $[s, t] \subset e_{0,1}$ contains a boundary gap. Put $u = 1 - t$ and $w = t - s$. Boundary coverage gives $B_0 \geq s$ and $A_1 \geq u$. Since the center has no trace on $e_{5,0}$,

$$A_0 + B_5 \geq 1, \quad A_0 + B_0 \leq 1,$$

and hence $B_5 \geq B_0 \geq s$. The exact CE1 formulas, after putting $R = \lambda$ in the orientation selected there, have the form

$$c_1 = u/R, \quad c_5 = 1 - \left(u - \frac{R}{1 + \sqrt{1 - R + R^2}} - \frac{R}{1 - R} w \right).$$

All hypotheses of Lemma C.7 hold. Therefore the outgoing reaches of rows 1 and 5 satisfy $\hat{b}_1 + \hat{b}_5 < 1$, where

$$\hat{b}_1 \leq F_{c_1}(u), \quad \hat{b}_5 \leq F_{c_5}(s).$$

Rows 2, 3, 4 must then supply boundary length at least

$$(1 - \hat{b}_1) + 1 + 1 + (1 - \hat{b}_5) > 3,$$

contrary to their three row caps.

Now let T_C be CE2. If exactly one of $[x, u]$, $[y, v]$ contains a gap, reflect so that it is $[y, v]$. The interval $[x, u]$ is covered by its endpoint roles, so the same endpoint propagation gives inputs y and $1 - v$. With

$$R = \frac{x}{x + y}, \quad s = y, \quad u' = 1 - v, \quad w' = v - y,$$

the CE2 coupling and exit formulas are exactly those of Lemma C.7. The preceding three-row contradiction applies unchanged. If both center intervals contain gaps, put

$$p = 1 - u, \quad q = 1 - v, \quad r = \frac{x}{x + y}, \quad w = \frac{y}{x + y}.$$

The far endpoint roles have inputs at least p and q , with radial demands p/w and q/r . By Lemma C.8, their two outgoing reaches sum to less than one, and the same middle three-row budget is impossible. These are the no-, one-, and two-active-interval states, and are exhaustive. $\square$

We next treat the additional T3-like rows in the $N_+ = 0$ branch. The following local facts also reappear later.

Lemma C.3 (T3-like midpoint and side tradeoff). *Let T_i be T3-like. Then $M_i \notin T_i$. After applying the closed-trace domination of Proposition 2.6, suppose the selected adjacent support is on r_{i+1} and M_{i+1} belongs to the dominated trace. If b is its boundary reach toward V_{i+1} , c its exit on its own arm, and ℓ its entry on r_{i+1} , all measured from the relevant vertex, then*

$$(80) \quad b + c < 1, \quad b + \ell > 1.$$

Consequently two adjacent T3-like traces that cover each other's midpoint, whose selected supports point toward one another, cannot cover both outer half-arms if no other role has positive support there.

Proof. Normalize at V_0 with wedge coordinates (x, y) and metric $x^2 + y^2 - xy$. Suppose first, toward the own-midpoint claim, that a unit triangle with two outside vertices A, B contains both $(0, 0)$ and $(1/2, 1/2)$. Diameter one implies that an outside vertex has a negative coordinate and that each of its coordinates is strictly below one. Order A, B and write

$$B - A = (r, s), \quad C - A = (s, s - r), \quad r^2 + s^2 - rs = 1.$$

Since $(0, 0)$ lies in the triangle, for some $\lambda, \mu \geq 0$ with $\lambda + \mu \leq 1$,

$$A = -\lambda(r, s) - \mu(s, s - r).$$

The midpoint condition is exactly

$$\lambda + \frac{r}{2} \geq 0, \quad \mu + \frac{s - r}{2} \geq 0, \quad \lambda + \mu + \frac{s}{2} \leq 1.$$

Put $t = s - r$. The four choices of negative coordinates of A, B give the required bounds on the third vertex as follows.

- (1) If $A_x, B_x < 0$, then $C_x = A_x + s < 1$ when $s \leq 1$. When $s > 1$, the unit equation gives $r, t > 0$, and $B_x < 0$ gives $C_x < t < 1$. For the other coordinate, $C_y = B_y - r < 1$ when $r \geq 0$; when $r < 0$, put $q = -r + s$ to obtain

$$C_y \leq q - \frac{q^2 - 1}{2} \leq 1.$$

- (2) If $A_y, B_y < 0$, the reflected argument gives the same two bounds.
- (3) If $A_x < 0$ and $B_y < 0$, the alternative $s > 1$ would imply $\mu > 1 - \lambda$, which is impossible. Hence $C_x < 1$. Also, $r < -1$ would imply $s < 0$ and then $A_x \geq 0$; consequently $C_y = B_y - r \leq 1$.
- (4) If $A_y < 0$ and $B_x < 0$, reflection of the preceding case applies.

In the $r < 0$ entry, $q = -r + s$ and $(-r)^2 + s^2 - rs = 1$ give $(-r)s = q^2 - 1$, while $\lambda \geq -r/2$ gives the displayed bound. Thus every vertex satisfies $x \leq 1, y \leq 1$, and only C can meet either equality line. Convexity makes both adjacent-arm intersections zero-length, contradicting the T3-like condition. Hence $M_0 \notin T_0$.

For the tradeoff, the translated T3-like trace has the exhaustive Type-II form

$$\begin{aligned} (1 - t)x + ty &\geq 0, \\ \beta + tx - y &\geq 0, \quad 0 < t < 1, \quad z = \sqrt{1 - t + t^2}, \quad 0 < \beta < z. \\ \beta + x - (1 - t)y &\leq z, \end{aligned}$$

For the selected support on r_1 , substitution in the Type-II half-planes gives

$$b = z - \beta, \quad c = \frac{\beta}{1 - t}, \quad \ell = \frac{\beta + 1 - z}{1 - t}.$$

The condition $M_1 \in T$ is $\beta \leq z - (1+t)/2$. Therefore

$$b + c \leq z + \frac{t}{1-t} \left(z - \frac{1+t}{2} \right) = 1 - \frac{(1-z)^2}{2(1-t)} < 1,$$

whereas

$$b + \ell - 1 = \frac{t(1-z+\beta)}{1-t} > 0.$$

For a crossed adjacent pair, coverage forces $c_i \geq \ell_{i+1}$ and $c_{i+1} \geq \ell_i$. Combining these two inequalities with (80) gives both $b_i < b_{i+1}$ and $b_{i+1} < b_i$, a contradiction. $\square$

Lemma C.4 (The support-isolated four-label inequality). *Assume T_C is CE1 or CE2, $N_+ = 0$, no role is Vd1 or Vd2, and, after normalization and reflection, T_0 is T3-like with selected adjacent support on r_1 . Then the two endpoint rows in the isolated configuration satisfy*

$$(81) \quad F_{c_5^*}(a_5^*) + F_{c_1^*}(a_1^*) < 1,$$

where a_1^*, a_5^* are the exact boundary inputs and c_1^*, c_5^* the exact radial demands defined below.

Proof. Use the CE1 side variables $0 < \lambda < 1$, $\rho = \sqrt{1 - \lambda + \lambda^2}$ and center slacks

$$X = F_0(O), \quad Y = F_2(O).$$

Then

$$a_C = \lambda - Y, \quad \lambda s = X + Y + 1 - \rho,$$

and

$$\gamma_1 = \min \left\{ \frac{Y}{\lambda}, \frac{\rho - X - Y}{1 - \lambda} \right\}, \quad \gamma_5 = \min \left\{ \frac{X}{1 - \lambda}, \frac{\rho - X - Y}{\lambda} \right\}.$$

For CE2 the second boundary interval is $[S, T]$, where

$$S = \frac{X + Y + 1 - \rho}{1 - \lambda}, \quad T = X + \lambda.$$

The T3-like normal form supplies $D > 1$, $0 < R < 1$, and $\alpha, p \geq 0$ with

$$R^2 - DR + D^2 = 1, \quad \alpha + p = 1/D, \quad q = 1 - p = \alpha + (D - 1)/D < 1/2.$$

Its interval on r_1 , measured from V_1 , is

$$[c, u], \quad c = Dq/R, \quad u = q + (1 - R)/D.$$

For an initial trace $[0, p]$ and a center interval $J = [L, U]$, define

$$\Theta(p, J) = \begin{cases} 1 - p, & J = \emptyset \text{ or } p < L, \\ 1 - \max\{p, U\}, & p \geq L. \end{cases}$$

Then

$$a_1^* = \Theta(p, [s, t]), \quad a_5^* = \Theta(\alpha, [S, T]),$$

with the second interval empty in CE1, and

$$c_5^* = 1 - \gamma_5, \quad c_1^* = \begin{cases} c, & 1 - u \leq \gamma_1 \leq 1 - c, \\ 1 - \gamma_1, & \text{otherwise.} \end{cases}$$

If $a_1^* + a_5^* > 1$, the two caps already imply $F_{c_5^*}(a_5^*) + F_{c_1^*}(a_1^*) < 1$. Assume the hard region $a_1^* + a_5^* \leq 1$. The following table is an exhaustive audit of the four possible first labels; the middle column records the exact decisive inequality.

first label	right labels	conclusion
L	Full	$e(\gamma_5) < a_1^*$
L	L, T_-	$F_{c_5^*}(a_5^*) + F_{c_1^*}(a_1^*) < 1$
L	T_+^{hi}	$e(\gamma_5) < a_1^*$
T_-	L, T_-	$F_{c_5^*}(a_5^*) + F_{c_1^*}(a_1^*) < 1$
T_-	Full, T_+^{hi}	$F_{c_5^*}(a_5^*) < a_1^*$
Full	all	impossible
T_+^{hi}	Full, T_+^{hi}	impossible
T_+^{hi}	L, T_-	$F_{c_5^*}(a_5^*) + F_{c_1^*}(a_1^*) < 1$.

We verify the nonformal entries. Put $\kappa = (1 - \rho)/(1 - \lambda)$. From $X + (1 - \lambda)Y < \rho(1 - \rho)$ one obtains

$$(82) \quad \gamma_1 \geq a_C \implies e(\gamma_5) < a_C,$$

$$(83) \quad S \leq Y/\lambda \implies e(\gamma_5) < S.$$

For (82), one has $\gamma_5 < a_C - \kappa$ and $a_C - \kappa < 1/8$; then (61) applies, and the remaining comparison is

$$(a_C - \kappa) + 5(a_C - \kappa)^2 \leq \kappa.$$

After squaring its positive sides, the difference is $\lambda(23\lambda^3 + 58\lambda^2 + 7\lambda + 72) > 0$. For (83), combine the premise with the center inequality and put $x = \lambda$. This gives $0 < x < 3/8$ and

$$X < X_* = \frac{(1 - \rho)(\rho(1 - 2x) - x(1 - x))}{1 - x - x^2}.$$

Set

$$\delta = \frac{1 - \rho}{1 - x} = \frac{x}{1 + \rho}, \quad \eta = \frac{X}{1 - x}, \quad f = \rho(1 - 2x) - x(1 - x), \quad d = 1 - x - x^2.$$

Then $\gamma_5 \leq \eta < \delta f/d$. Moreover

$$\frac{f}{d} < 1 - 2x,$$

because, after multiplication by positive denominators, this is just

$$\rho < d + \frac{x(1 - x)}{1 - 2x} = 1 + \frac{2x^3}{1 - 2x}.$$

Hence $\eta < \delta(1 - 2x) < 1/8$. If $\eta = 0$, the assertion is immediate. Otherwise,

$$S \geq \frac{X + 1 - \rho}{1 - 2x} = \frac{1 - x}{1 - 2x}(\eta + \delta) > 2\eta \left(\frac{1 - x}{1 - 2x} \right)^2.$$

On the other hand,

$$5\eta < 5\delta(1 - 2x) < 2 \left[\left(\frac{1 - x}{1 - 2x} \right)^2 - 1 \right].$$

For the middle sign use $\rho > 1 - x$ and

$$2(2 - 3x)(2 - x) - 5(1 - 2x)^3 = (4x - 3)(10x^2 - 6x - 1) > 0 \quad (0 < x < 3/8).$$

Indeed both factors are negative; for the quadratic use $10x^2 \leq 15x/4 < 6x+1$. The low-root estimate (61) now gives

$$e(\gamma_5) \leq 2\gamma_5 + 5\gamma_5^2 \leq 2\eta + 5\eta^2 < S,$$

proving (83) without any polynomial coefficient list. The L separation follows from its exact polynomial: if its output is $z = e(\eta)$ at input $1 - \beta$, then

$$\beta \geq z + \eta.$$

Indeed the selector polynomial $P(x, z) = x^4 - 4x^3 + 5x^2 - (2+z)x + z - z^2$ is strictly decreasing on the low range and vanishes at $x = \eta$. Equations (82)–(C.2), with the three possibilities for Θ , prove every first- L entry. For a first T_- label the exact root test

$$T_-(a, c) < z \iff c^2(a+z)^2 - (a^2 - ac + c^2) > 0$$

is increasing in z . Substitution of a_1^* , followed by the center inequality, proves $F_{c_5^*}(a_5^*) < a_1^*$ except in the low-low entries, which are immediate from $T_-(a, c) \leq \min\{a, 1-a\}$ and $L < 1/2$. First-coordinate Full would force either $X \geq (1-\lambda)\alpha$ and $Y \geq \lambda - \alpha$, contradicting $s \leq p$, or, in the CE2 overlap case, $S \geq 1/2$, contradicting $S \leq \alpha < 1/2$.

For a first T_+^{hi} label put $r = 1 - \lambda$, $y = Y/\lambda$ and choose β so that

$$m = \sqrt{\beta^2 - \beta + 1}, \quad 1 - \gamma_5 = \frac{m}{r + \beta}, \quad F_{c_5^*}(a_5^*) = \frac{\beta m}{r + \beta}.$$

The exact selector is

$$(84) \quad \beta \geq \max\left\{r, \frac{1}{2}, \frac{1-r^2}{1+2r}\right\}, \quad r \geq (1-\beta)(r+\beta)^2.$$

The center inequalities give $y < 1/10$ and, with

$$y_* = \frac{r}{1-r} \left(\frac{m}{r+\beta} - \frac{1}{2} - \frac{1-r}{1+\sqrt{r^2-r+1}} \right),$$

give $y < y_*$ and

$$3y_* \leq 1 - \frac{\beta m}{r + \beta}.$$

This proves the right- L entry by $e(y) < 3y$. It excludes right Full and right high by the exact high-input filter. For right T_- with input a_C , direct substitution gives a sum below one. For input q , put $c = 1 - y$, $q = tc$, and let $\hat{b}_1 = F_{c_1^*}(q)$. The selected T_- equations give

$$\hat{b}_1 \leq \kappa := \frac{\sqrt{13}-1}{6}, \quad q \leq \tau < \frac{93}{200},$$

where $\tau \in (0, 1/2)$ is the unique root of $\tau^4 - 3\tau^3 + 3\tau^2 - 3\tau + 1 = 0$. In this CE2 overlap subcase the support ordering gives $q > S$. The common left-high identities are

$$S = S_0 + \frac{1-r}{r}y, \quad S_0 = 1 - \frac{m}{r+\beta} + \frac{1-r}{1+\sqrt{r^2-r+1}}.$$

Since $y > 0$, we have the complete chain

$$S_0 < S < q \leq \tau < \frac{93}{200}.$$

We now replace the former interval subdivision by a one-variable comparison. Write $s = r + \beta$ and $\rho_r = \sqrt{r^2 - r + 1}$. The selected Cell 2 inequality factors as

$$r - (1 - \beta)(r + \beta)^2 = (s - 1)(\beta^2 + r\beta - r).$$

The second factor is nonnegative because $r \leq \beta$ and $\beta \geq 1/2$; if it vanishes, then $r = \beta = 1/2$. Hence $s \geq 1$. It follows that $\rho_r \leq m$, since $\rho_r^2 - m^2 = (r - \beta)(s - 1) \leq 0$.

Define the one-variable lower bound

$$\bar{S}(\beta) = 1 - m + \frac{\beta}{1 + m}.$$

Then

$$S_0 \geq 1 - \frac{m}{s} + \frac{1 - r}{1 + m} \geq \bar{S}(\beta).$$

Indeed, the second difference is

$$(s - 1) \left(\frac{m}{s} - \frac{1}{1 + m} \right) \geq 0.$$

For the last sign use $s \leq 2\beta \leq m(1 + m)$: if $g = 3\beta - \beta^2 - 1$, then $g > 0$ and

$$m^2 - g^2 = -\beta(\beta - 1)(\beta^2 - 5\beta + 5) \geq 0, \quad m^2 + g = 2\beta.$$

Put $B_0 = 5657/10000$, $T = 93/200$, and $\beta_* = 161/250$. If $\beta m/s \geq B_0$, then $\beta m \geq B_0$. The function $\phi(\beta) = \beta m$ is strictly increasing, and the exact comparison

$$B_0^2 - \beta_*^2 m(\beta_*)^2 = \frac{22782769}{62500000000} > 0$$

therefore gives $\beta > \beta_*$. On $[\beta_*, 1]$ put

$$H(\beta) = \frac{107}{200} - Tm - (1 - \beta)^2.$$

Since $H'' = -3T/(4m^3) - 2 < 0$, the function H is concave. Its endpoint values are positive: $H(1) = 7/100$, while, for $a_* = 107/200 - (1 - \beta_*)^2 = 51033/125000$,

$$a_*^2 - T^2 m(\beta_*)^2 = \frac{1693881}{62500000000} > 0.$$

Thus $H > 0$ throughout the interval. The identity

$$(\bar{S}(\beta) - T)(1 + m) = H(\beta)$$

shows that $\beta m/s \geq B_0$ would force $S_0 > T$, a contradiction. Consequently

$$F_{c_5^*}(a_5^*) < \frac{5657}{10000} < \frac{7 - \sqrt{13}}{6}.$$

The final strict comparison follows by squaring $\sqrt{13} < 18029/5000$, whose squared difference is $44841/25000000 > 0$. Thus the final sum is also below one. The table is exhaustive, proving (81). $\square$

Proposition C.5 (The $N_+ = 0$ T3-like branch). *Assume that T_C is CE1 or CE2, $N_+ = 0$, no vertex role is Vd1 or Vd2, and at least one vertex role is T3-like. Then the seven roles do not cover H .*

Proof. Normalize the unique center midpoint to M_0 and simultaneously apply the closed-trace domination of Proposition 2.6 to every T3-like role. If T_0 were not T3-like, a maximal consecutive block of T3-like indices among $1, \dots, 5$ would have to match its roles bijectively to the same block of uncovered midpoints. The matching uses only adjacent indices, because a T3-like role misses its own midpoint and Vd0 roles cannot cover adjacent midpoints. Its first two roles therefore form a crossed pair forbidden by Lemma C.3. Hence T_0 is T3-like; reflect so that it supports r_1 .

The diagonal trace bounds from Section 3 imply that there are at most two T3-like roles: if there were $k \geq 3$, the center, T3-like, and nonsupercritical Vd0 diagonal caps would give

$$6 < \frac{3}{2} + \frac{k}{2} + (6 - k) \leq 6.$$

Midpoint locality then forces T_2, T_4, T_5 to be Vd0. Consequently r_1 has positive support only from T_C, T_0, T_1 , and r_5 only from T_C, T_5 .

The actual outgoing reaches of T_1, T_5 are bounded by the two safe maps in Lemma C.4; an extra adjacent-ray demand on a possible T3-like T_1 only shrinks its feasible set. Thus their sum is less than one. Rows T_2, T_3, T_4 must cover more than three units of the four intervening boundary edges, while their three nonsupercritical caps sum to at most three. This contradiction proves the proposition. $\square$

Lemma C.6 (Free strict-supercritical envelope). *For $0 \leq c < 1/2$,*

$$\sup\{b : \exists a, (a, b, c) \in \mathcal{A}, a + b > 1\} = \mathcal{B}(c) := \frac{c + \sqrt{c^2 - 8c + 4}}{2}.$$

The supremum is not attained. For $c \geq 1/2$ the strict feasible set is empty. The function $\mathcal{B}$ is strictly decreasing on $[0, 1/2]$.

Proof. Lemma 2.10 excludes $c \geq 1/2$. On the boundary $a = 1 - b$, a support rotation has strict slack precisely when

$$c < \frac{1 - b^2}{2 - b}.$$

The right side decreases from $1/2$ to zero. Solving equality gives the displayed larger root. Equality has no strict support slack, which explains nonattainment. Differentiation gives

$$\mathcal{B}'(c) = \frac{1}{2} \left(1 + \frac{c - 4}{\sqrt{c^2 - 8c + 4}} \right) < 0,$$

since $(4 - c)^2 - (c^2 - 8c + 4) = 12$. $\square$

C.3. Two reusable capped-loss inequalities.

Lemma C.7 (One-side capped loss). *Let $0 < R < 1$, $S = \sqrt{1 - R + R^2}$, and let $s, u, w > 0$ satisfy $s + u + w = 1$ and $0 < u < R$. Put*

$$\delta = \frac{R}{1 + S}, \quad \gamma = u - \delta - \frac{R}{1 - R}w,$$

and assume $\gamma > 0$. Set

$$c_1 = u/R, \quad c_5 = 1 - \gamma.$$

Then

$$(85) \quad F_{c_5}(s) + F_{c_1}(u) < 1.$$

Proof. Put $\eta = (1 - R)\delta = 1 - S$ and

$$\widehat{b}_5 = F_{c_5}(s), \quad \widehat{b}_1 = F_{c_1}(u).$$

The center identity may be written

$$(86) \quad c_5 = \frac{1 - u + \eta - Rs}{1 - R}.$$

The hypotheses imply $1/2 < c_1, c_5 < 1$: for example $u > \delta$ gives $c_1 > 1/(1 + S) > 1/2$, and $\gamma < u - \delta < R - \delta = RS/(1 + S) < 1/2$ gives $c_5 > 1/2$. Thus the four high-radial labels of Proposition A.3 are exhaustive.

Right low labels. If the right label is T_- , its exact frontier equation gives $(u + \widehat{b}_1)^2 = 1 - R + R^2 = S^2$, so $\widehat{b}_1 = S - u$. For a right L label write $\widehat{b}_1 = kc_1$, $0 \leq k \leq 1/2$, and $C = 1 - k + k^2 = c_1^2$. With $X_R = R + k$, the Cell- L selector factors as

$$q = c_1^2(X_R - 1)G_k(X_R), \quad G_k(X) = CX^3 + CX^2 - k(1 - k)X + k^2.$$

For $X \geq k$, $G_k(X) > 0$: when $k > 0$, $G'_k(X) \geq k(1 + 2k - k^2 + 3k^3) > 0$ and $G_k(k) = k^2(1 + k + k^3) > 0$. Hence $q \leq 0$ gives $X_R \leq 1$. If $h_R = 1 - X_R$, then

$$S^2 - (u + \widehat{b}_1)^2 = h_R(1 + 2k^2 + k(1 - k)h_R) \geq 0.$$

Consequently either right low label satisfies

$$(87) \quad \widehat{b}_1 \leq X := S - u.$$

A left low label against right high or Full. For left T_- put $z = s/c_5 = 1 - p$ and $h = \sqrt{1 - p + p^2}$. The component selector gives $0 \leq p \leq 1/2$, the frontier gives $s + \widehat{b}_5 = h$, and its selector gives $s \leq (1 - p)h$. Equation (86) becomes $c_5(1 - Rp) = 1 - u + \eta$, whence

$$u \geq \eta + (1 - h) + Rph.$$

Put $q_0 = ph$. If $q_0 > R$, then $p(1 - p) > R(1 - R)$ and the preceding lower bound would give $u > R$. Thus $q_0 \leq R$, and

$$\eta + 1 - h > \frac{R(1 - R) + p(1 - p)}{2} \geq (1 - R)q_0.$$

Substitution in the exact formula

$$w = \frac{(1 - R)(1 - u)p - \eta(1 - p)}{1 - Rp}$$

now gives $w < 1 - h$. A right high output has the form $H - u$, where $H = \sqrt{1 - \beta + \beta^2} \leq 1$; for Full take $H = 1$. Hence

$$\widehat{b}_5 + \widehat{b}_1 = h + H - (s + u) = 1 + w - (1 - h) - (1 - H) < 1.$$

For a left L label write

$$c_5 = h(x) := \sqrt{1 - x + x^2}, \quad \widehat{b}_5 = xh(x), \quad 0 \leq x \leq \frac{1}{2},$$

and, along the center line,

$$s(x) = \frac{R - u + \eta + (1 - R)(1 - h(x))}{R}.$$

If $y = s(x)/h(x)$, the exact selector factorization is

$$\frac{q}{h(x)^2} = (y - (1 - x))P_x(y),$$

where

$$P_x(y) = (1 - x + x^2)y^3 + (1 + 2x - 2x^2 + 3x^3)y^2 \\ + (x + 2x^2 - x^3 + 3x^4)y + (x^2 + x^3 + x^5).$$

Every coefficient is nonnegative on $0 \leq x \leq 1/2$, and the polynomial is positive at every nondegenerate point. Thus a selected L point has $s(x) \leq (1 - x)h(x)$. The two curves meet once in $(0, 1/2)$: at $x = 0$ the left side is smaller, while at $x = 1/2$ it is enough to use

$$u \leq \frac{R}{1+R} \quad (\text{Full}), \quad u \leq \min\{RS, S/2\} \quad (T_+^{\text{hi}}).$$

Writing $e_0 = 1 - \sqrt{3}/2$, the corresponding endpoint differences are

$$\frac{(1-R)(R^2 + 2Re_0 + 2e_0)}{2(1+R)} > 0$$

for Full, and respectively $[2e_0(1-R) - R^3]/2 \geq 0$ for $R \leq 1/2$ and $(1-R)(3R + 4e_0 - 2)/4 \geq 0$ for $R \geq 1/2$ in the high case. At the unique transition x_0 , the L frontier is the $q = 0$ endpoint of the preceding T_- branch. Since $xh(x)$ increases, every selected left L output is no larger than that transition output. The preceding strict loss therefore also covers (L, T_+^{hi}) and (L, Full) .

Full and high exclusions. Left Full forces $s < 1/2$ and $\gamma \geq s$. Right Full or right high gives $u \leq 1/2$, but

$$\gamma - s = 2u - 1 - \delta + \frac{1 - 2R}{1 - R}w < 0 :$$

for $R \geq 1/2$ this is immediate, while for $R < 1/2$ use $w < 1 - u$ to bound it by $(u - R)/(1 - R) - \delta < 0$. Thus left Full cannot pair with either label. If the left label is high and the right one Full, then $s < 1/2$ and $u \leq R/(1 + R)$, whereas $\gamma > 0$ and $w > 1/2 - u$ force $u > R/2 + \eta > R/(1 + R)$; the final inequality is equivalent to $2(1 + R) > 1 + S$. This pair is also impossible.

A right low label. Use (87). If $s > X$, the cap gives $\hat{b}_5 + \hat{b}_1 < 1$. Suppose $s \leq X$ and put

$$c_X = X + 2\delta, \quad f(c, z) = c^4 - c^2 + zc - z^2.$$

At the limiting value $u = R$, with $\kappa = \delta$, one has

$$X_0 = \frac{1 - 2\kappa}{1 + \kappa}, \quad c_0 = \frac{1 + 2\kappa^2}{1 + \kappa}.$$

Now $c_0 > \sqrt{3}/2$ because $16\kappa^4 + 13\kappa^2 - 6\kappa + 1 > 0$, and

$$f(c_0, X_0) = \frac{\kappa^2(1 - 2\kappa)^2(4\kappa^4 + 4\kappa^3 + 10\kappa^2 + 6\kappa + 5)}{(1 + \kappa)^4} > 0.$$

Along the u -line, $\frac{d}{du}f(c_X, X) = -4c_X^3 + c_X + X < 0$, so the same two signs hold before the limiting endpoint. Therefore, for $\ell(c) = c(1 - \sqrt{4c^2 - 3})/2$,

$$\ell(c_X) < X < c_X - \ell(c_X), \quad \ell(c_X) < 1 - X.$$

Write $c = h(z) = \sqrt{1 - z + z^2}$ and let z_X satisfy $h(z_X) = c_X$. The actual $c_5(s)$ corresponds to $0 < z \leq z_X < 1/2$, and

$$s(z) = s_0 + \frac{1 - R}{R}(1 - h(z)), \quad s_0 = \frac{R - u + \eta}{R} > 0.$$

The transition input $(1 - z)h(z)$ decreases, and the preceding root ordering gives $s(z_X) < (1 - z_X)h(z_X)$, hence the same strict inequality for all $z \leq z_X$. To check the order of the two boundary coordinates put

$$\phi(z) = s(z) - zh(z), \quad k_R = \frac{1 - R}{R}.$$

Its endpoint values are $s_0 > 0$ and $X - \ell(c_X) > 0$. Moreover

$$\phi'(z) = \frac{(k_R + z)(1 - 2z)}{2h(z)} - h(z).$$

For $k_R \leq 2$ this is nonpositive; for $k_R > 2$ its zero is described by the strictly increasing function $K_0(z) = (2 - 3z + 4z^2)/(1 - 2z)$, so the minimum is again at an endpoint. Thus $zh(z) < s(z)$. Applying the already displayed polynomial P_z gives the strict Cell- L selector, and $s + zh(z) \leq X + \ell(c_X) < 1$. Since $\partial_b F_L = c(1 - 2z) > 0$ and each feasible fiber is an interval, the capped maximum is

$$\widehat{b}_5 \leq zh(z) \leq z_X h(z_X) = \ell(c_X) < 1 - X.$$

Together with $\widehat{b}_1 \leq X$, this proves the loss for every right low label.

High-high. Let $c_L(z)$ be the unique root in $[\sqrt{3}/2, 1]$ of $f(c, z) = 0$. With $D_c = 4c^3 - 2c + z > 0$, implicit differentiation gives

$$c_L''(z) = \frac{2(c^2 - cz + z^2) + 16c^2 z(c - z)}{D_c^3} > 0.$$

A selected left high label requires $s < 1/2$ and $c_5(s) \leq c_L(s)$. Define

$$s_0 = \frac{R - u + \eta}{R};$$

then $0 < s_0 < s$ and $c_5(s_0) = 1$. The explicitly named function $g(z) = c_5(z) - c_L(z)$ is concave, with $g(s_0) > 0$. The right high bound $u \leq \min\{RS, S/2\}$ gives $c_5(1/2) > 7/8$. For $R \leq 1/2$ this follows after squaring from $17 - 10R + 9R^2 - 64R^3 - 64R^4 > 0$ (a decreasing polynomial with value $9/4$ at $1/2$); for $R \geq 1/2$ it follows from $(3 + R)^2 - 16S^2 = (1 - R)(15R - 7) > 0$. Finally

$$f(7/8, 1/2) = 33/4096 > 0$$

and f increases in c on this range, so $c_L(1/2) < 7/8$ and $g(1/2) > 0$. Concavity contradicts $g(s) \leq 0$. Thus high-high is impossible.

If both labels are low, the cap already gives $\widehat{b}_5 + \widehat{b}_1 \leq s + u = 1 - w < 1$. The preceding five paragraphs cover every other ordered pair of the four labels, proving (85). $\square$

Lemma C.8 (CE2 two-endpoint capped loss). *Let $[x, U]$ and $[y, v]$ be an exact CE2 pair as in Proposition A.4. Put*

$$p = 1 - U, \quad q = 1 - v, \quad r = \frac{x}{x + y}, \quad w = \frac{y}{x + y},$$

and reflect if necessary so that $0 < w \leq 1/2$ and $r = 1 - w$. Then

$$(88) \quad F_{p/w}(p) + F_{q/r}(q) < 1.$$

Proof. Put

$$S_0 = x + y, \quad K = \sqrt{1 - wr}, \quad z = \frac{w}{1 + K}, \quad \zeta = \frac{r}{1 + K}, \quad \alpha = \frac{p}{w}, \quad \beta = \frac{q}{r}.$$

Then $D = S_0 K$, $1 - K = wr/(1 + K)$, and the exact midpoint inequalities give $1/2 < \alpha, \beta < 1$. Dividing the coupling equation $(U + v)S_0 - xy = D$ by S_0 , and using successively $x < U$ and $y < v$, gives the strict endpoint inequalities

$$(89) \quad \beta + p > 1 + z, \quad \alpha + q > 1 + \zeta.$$

Define

$$\Phi_w(\alpha) = F_\alpha(w\alpha), \quad \Phi_r(\beta) = F_\beta(r\beta), \quad h(t) = \sqrt{1 - t + t^2}.$$

Thus the desired sum is $\Phi_w(\alpha) + \Phi_r(\beta)$.

Selected branch parametrizations. For a side of weight $\sigma \in \{w, r\}$, every L label has

$$(90) \quad c = h(k), \quad F_c(\sigma c) = kh(k), \quad 0 \leq k \leq w,$$

on either side. On the large side the range follows from

$$Q_{\text{cell}}/h(k)^2 = (k - w)G_k(r + k),$$

where $G_k(X) = h(k)^2 X^3 + h(k)^2 X^2 - k(1 - k)X + k^2 > 0$ for $X \geq k$. The remaining selected parametrizations are

$$(91) \quad \alpha = \frac{h(a)}{w + a}, \quad \Phi_w(\alpha) = a\alpha, \quad r \leq a < 1,$$

$$(92) \quad \beta = \frac{h(b)}{r + b}, \quad \Phi_r(\beta) = b\beta, \quad r \leq b < 1,$$

$$(93) \quad \beta = \frac{K}{r + t}, \quad \Phi_r(\beta) = t\beta = K - r\beta, \quad w \leq t \leq r.$$

For example, the small-high selector factors as

$$h(a)^2 a \left(\frac{w}{(w + a)^2} - (1 - a) \right), \quad w - (1 - a)(w + a)^2 = (a - r)(a^2 + wa - w),$$

which gives $a \geq r$; the other two factorizations are $(b - w)(b^2 + rb - r)$ and $(t - w)(r^2 - wt)$, respectively. These formulas specify the selected geometric components, not discarded algebraic roots.

The small-side T_- label is absent for $w < 1/2$. Large-side Full is incompatible with the first inequality in (89). Small-side Full can pair only with L : either high or T_- on the large side has $\beta \leq K$, which would force $p > (1 + r)z > w/(1 + w)$, contrary to Full. Thus, up to the symmetric $w = r = 1/2$ tie, the exact list is

$$(94) \quad (\text{Full}, L), (T_+^{\text{hi}}, L), (L, L), (L, T_-), (L, T_+^{\text{hi}}), (T_+^{\text{hi}}, T_-), (T_+^{\text{hi}}, T_+^{\text{hi}}).$$

Low-low sums are at most $K < 1$: an L output is at most wK , while (93) is at most rK . For Full- L , put

$$b_{\text{thr}} = 1 + z - \frac{w}{1 + w}, \quad \kappa = \frac{w}{1 + (1 + w)z}.$$

Full and (89) give $\beta > b_{\text{thr}}$ and $(1 + \kappa)b_{\text{thr}} = 1 + z$. Direct reduction yields

$$b_{\text{thr}}^2 - h(\kappa)^2 = \frac{w^2(A(w) + KC(w))}{D_0(w, K)} > 0,$$

where

$$\begin{aligned} A(w) &= 8 - 5w + 22w^2 + 6w^3 + 5w^4 + 5w^5 - w^6, \\ C(w) &= 8 - w + 14w^2 + 4w^3 - 2w^4 + w^5, \\ D_0(w, K) &= ((1+w)(1+K))^2(1+K+w+w^2)^2. \end{aligned}$$

Both numerator polynomials are positive on $[0, 1/2]$. Since h decreases and $(1+k)h(k)$ increases there, $\beta = h(k) > b_{\text{thr}} > h(\kappa)$ implies $(1+k)\beta < 1+z$; combined with (89), this gives $k\beta < p$ and hence the strict Full- L loss.

It remains to verify the four pairs containing a high label on the small side. (L, T_+^{hi}) . Use parameters $0 \leq k \leq w$ and $r \leq t < 1$:

$$\alpha = h(k), \quad \Phi_w = kh(k), \quad \beta = \frac{h(t)}{r+t}, \quad \Phi_r = \frac{th(t)}{r+t}.$$

The first output and the last output increase in their parameters, while $h(t)/(r+t)$ decreases. If $w \leq 2/5$, then

$$\Phi_w + \Phi_r < wK + \frac{1}{2-w} < 1;$$

after using $K \leq 1 - wr/2$, the positive cleared gap is $P(w) = w^4 - 3w^3 + 4w^2 - 6w + 2$, which decreases on $[0, 2/5]$ and has $P(2/5) = 46/625$. If $w \geq 2/5$, the first endpoint inequality gives $\beta > r + z \geq 3/4$. Since $h(3/4)/(r + 3/4) = \sqrt{13}/(7 - 4w) < 3/4$, monotonicity forces $t < 3/4$. Therefore

$$\Phi_w + \Phi_r < \frac{\sqrt{3}}{4} + \frac{3\sqrt{13}}{20} < 1.$$

(T_+^{hi}, T_-) . Use a, t from (91) and (93). Both demands are at most K . The first endpoint inequality gives

$$\alpha > \frac{1+z-K}{w} = \frac{1+r}{1+K} =: a_0.$$

Thus $K > 1 - w^2$, which forces $w > 1/3$, and comparison at $a = 2/3$ forces $a < 2/3$. The function

$$G(a) = \frac{(1+a)h(a)}{w+a}$$

decreases on $[r, 2/3]$, because its derivative numerator $2a^3 + (4w-1)a^2 + (1-w)a + w - 2$ is at most its value $-7/54$ at $(2/3, 1/2)$. Hence $G(a) \leq (1+r)K$. Using the second endpoint inequality and $\Phi_r = K - r\beta$ gives

$$\Phi_w + \Phi_r < (2+r)K - 1 - \zeta < 1.$$

After multiplication by $1+K$, the last positive gap is $r(3w - w^2 - K)$; its squared comparison is

$$(3w - w^2)^2 - K^2 = w^4 - 6w^3 + 8w^2 + w - 1 > 0$$

for $w > 1/3$ (value $1/81$ at $1/3$, positive derivative thereafter).

$(T_+^{\text{hi}}, T_+^{\text{hi}})$. Use parameters a, b from (91)–(92). The endpoint inequalities first force $w > 2/5$: for $w \leq 2/5$, the asserted contrary inequality $K(w + 1/(2r)) \leq 1 + z$ reduces, after multiplication by $2r(1+K)$, to

$$K(2w^2 - 4w + 1) + 2w^4 - 4w^3 + w^2 - w + 1 > 0,$$

which is immediate according as the coefficient of K is nonnegative or, using $K \leq 1$, from the decreasing polynomial $2w^4 - 4w^3 + 3w^2 - 5w + 2 \geq 172/625$.

For $F_c(t) = (1+t)h(t)/(c+t)$, the derivative numerator $2t^3 + (4c-1)t^2 + (1-c)t + c - 2$ is increasing on the present rectangle, so F_c has at most one interior minimum. Endpoint comparison gives

$$\Phi_w + \alpha \leq \frac{2}{1+w}, \quad \Phi_r + \beta \leq \frac{2}{1+r}.$$

The nontrivial endpoint signs are exactly

$$4 - (1+r)^2(1+w)^2K^2 = -w^2(w-1)^2(w^2-w-3) > 0$$

and

$$16r^2 - (1+r)^4h(r)^2 = (1-r)(r^5 + 4r^4 + 7r^3 + 9r^2 - 4r - 1) > 0;$$

the bracket is increasing on $[1/2, 1]$ and equals $13/32$ at $1/2$. Weighting the two strict endpoint inequalities by w/K^2 and r/K^2 gives

$$\alpha + \beta > \frac{2K+1}{K(1+K)}.$$

Consequently

$$\Phi_w + \Phi_r < \frac{2}{1+w} + \frac{2}{1+r} - \frac{2K+1}{K(1+K)} < 1.$$

The numerator of the last gap is $3Kwr - Q(w)$, where $Q(w) = w^4 - 2w^3 + 7w^2 - 6w + 2 > 0$. Its squared difference is

$$D(w) = -w^8 + 4w^7 - 9w^6 + 13w^5 - 41w^4 + 65w^3 - 55w^2 + 24w - 4.$$

Here $D'(w) = (1-2w)H(w)$ with

$$H(w) = 4w^6 - 12w^5 + 21w^4 - 22w^3 + 71w^2 - 62w + 24 > 0$$

on $[2/5, 1/2]$; use $71w^2 - 62w + 24 \geq 743/71$ and the remaining cubic block at least $-11/4$. Thus D increases and $D(2/5) = 4904/390625 > 0$, proving the claim.

(T_+^{hi}, L) . Use the fully specified parameters

$$\alpha = \frac{h(t)}{w+t}, \quad p = w\alpha, \quad b = t\alpha, \quad \frac{1}{2} \leq t < 1,$$

and

$$\beta = H = h(k), \quad \widehat{b}_L = kH, \quad 0 \leq k \leq w.$$

Since $p + b = h(t)$, the first endpoint inequality gives

$$(95) \quad 1 - (b + kH) > G := 2 + z - h(t) - (1+k)H.$$

Put

$$p_0 = 1 + z - K = \frac{w(2-w)}{1+K},$$

let $t_0 \in (1/2, 1)$ be the unique point with $p(t_0) = wh(t_0)/(w+t_0) = p_0$, and put $b_0 = t_0 p_0/w$. The exact endpoint lemma is

$$(96) \quad b_0 < 1 - wK.$$

To verify it, set $b_* = 1 - wK$ and $T_* = wb_*/p_0$. One has $0 < T_* < 1$, and

$$p_0^2(w + T_*)^2 - w^2h(T_*)^2 = -\frac{w^3(KE(w) + J(w))}{(1+K)^2(2-w)^2} > 0,$$

where

$$\begin{aligned} E(w) &= 2w^5 + 3w^3 - 20w^2 + 23w - 6, \\ J(w) &= w^6 + 8w^5 - 22w^4 + 12w^3 + 6w^2 + 2w - 6. \end{aligned}$$

Here $J \leq -111/64$; if $E > 0$, then $KE + J < E + J < 0$, because $E + J$ is increasing and equals $-139/64$ at $w = 1/2$. Thus $p(T_*) < p(t_0)$. Since $p(t)$ decreases, $t_0 < T_*$ and (96) follows.

Finally put $P = 1 + z - p$. If $P < K$, then $t < t_0$ and $k \leq w$, so

$$G > 2 + z - h(t_0) - (1 + w)K = 1 - b_0 - wK > 0.$$

If $P \geq K$, then $K \leq P \leq P_1 := 1 + z - w/(1 + w) \leq 1$ and there is $a \in [0, w]$ with $h(a) = P$. Since $P < H$, one has $k < a$ and $(1 + k)H < (1 + a)P = P + \ell(P)$, where $\ell(P) = P(1 - \sqrt{4P^2 - 3})/2$. Hence

$$G > \mathcal{D}(P) := 1 - b(P) - \ell(P).$$

This function is concave. Indeed $\ell''(P) > 0$, and the high output b is convex in $p = 1 + z - P$: with $A_t = 2 + w - (1 + 2w)t > 0$ and $Q_t = 8wt^2 + 4t^2 - 8wt - 13t - w + 4 < 0$,

$$\frac{d^2b}{dp^2} = -\frac{2h(t)(t + w)^3 Q_t}{w^2 A_t^3} > 0.$$

At $P = K$, (96) gives $\mathcal{D}(K) = 1 - b_0 - wK > 0$. At $P = P_1$, set $\kappa = w/(1 + (1 + w)z)$. The Full- L comparison already proved gives $P_1 > h(\kappa)$, hence $\ell(P_1) < \kappa P_1 = w/(1 + w)$, while $b(P_1) = 1/(1 + w)$. Thus $\mathcal{D}(P_1) > 0$, and concavity gives $G > 0$ throughout. Equation (95) proves the final hard pair.

The exact list (94) is now exhausted, proving (88). $\square$

C.4. Half-edge envelopes for adjacent CE2 placements. The adjacent CE2 placement in Section C uses a rational upper bound for a single admissible-set fiber. We isolate it here because the domain restriction is essential and because the same formula is false on the full nonsupercritical fiber domain.

Lemma C.9 (Half-edge rational radial envelope). *Suppose*

$$\frac{1}{2} \leq p < 1, \quad 0 \leq h \leq p, \quad p + h < 1.$$

Then

$$(97) \quad c_{\max}(p, h) \leq 1 - \frac{h}{3}.$$

The inequality is strict when the selected admissible-set component is $\mathcal{A}_T$.

Proof. Put

$$s = p + h, \quad d = p^2 + ph + h^2, \quad q = s^4 - s^2 + ph.$$

Because $s < 1$, one has $d = s^2 - ph < 1$, so only the two nonsupercritical cells of Proposition A.2 occur. Since $p \geq h$, in $\mathcal{A}_L$ the selected radial value is the unique root $C_L(h) \in [\sqrt{3}/2, 1]$ of

$$P_h(c) := c^4 - c^2 + hc - h^2 = 0,$$

with $C_L(0) = 1$. In $\mathcal{A}_T$ the selector $c \leq 2p$ keeps the smaller geometric root

$$C_T(p, h) = \frac{2p}{1 + \sqrt{4s^2 - 3}}.$$

The two expressions agree when $q = 0$.

First assume $q \leq 0$. If $h \geq 3/8$, then $p \geq 1/2$ and $s \geq 7/8$. On this region

$$\frac{\partial q}{\partial p} = 4s^3 - 2s + h \geq 4 \left(\frac{7}{8}\right)^3 - 2 \left(\frac{7}{8}\right) + \frac{3}{8} = \frac{167}{128} > 0.$$

Moreover

$$\frac{d}{dh} q \left(\frac{1}{2}, h \right) = h(4h^2 + 6h + 1) > 0.$$

Thus

$$q(p, h) \geq q \left(\frac{1}{2}, \frac{3}{8} \right) = \frac{33}{4096} > 0,$$

a contradiction. Hence $0 \leq h < 3/8$. At $c_0 = 1 - h/3$,

$$P_h(c_0) = \frac{h}{81} (h^3 - 12h^2 - 63h + 27).$$

The cubic in parentheses decreases on $[0, 3/8]$ and has value $891/512$ at $3/8$, so this expression is positive for $h > 0$. Also $c_0 > 7/8 > \sqrt{3}/2$, and

$$P'_h(c) = c(4c^2 - 2) + h > 0 \quad (c \geq \sqrt{3}/2).$$

Therefore the selected root lies below c_0 . The case $h = 0$ gives equality.

Now assume $q > 0$. Put

$$r = \sqrt{4s^2 - 3}, \quad \delta = \frac{p - h}{s}.$$

The exact identity

$$\frac{q}{s^2} = \frac{r^2 - \delta^2}{4}$$

gives $0 \leq \delta < r$. Thus $r > 0$ and we may write $\delta = wr$ with $0 \leq w < 1$. Then

$$p = \frac{s(1 + wr)}{2}, \quad h = \frac{s(1 - wr)}{2},$$

and

$$C_T(p, h) = \frac{s(1 + wr)}{1 + r}.$$

For fixed s, r , set

$$\Psi(w) = C_T(p, h) + \frac{h}{3}.$$

A direct differentiation gives

$$\Psi'(w) = \frac{sr(5 - r)}{6(1 + r)} > 0,$$

so

$$\Psi(w) < \Psi(1) = \frac{s(7 - r)}{6}.$$

The hypothesis $p \geq 1/2$ gives $s(1 + wr) = 2p \geq 1$, and hence $s(1 + r) \geq 1$. This forces $r > 1/8$: otherwise

$$4s^2(1 + r)^2 = (r^2 + 3)(1 + r)^2 \leq \frac{193}{64} \frac{81}{64} = \frac{15633}{4096} < 4.$$

Finally,

$$144 - (r^2 + 3)(7 - r)^2 = (1 - r)(r^3 - 13r^2 + 39r - 3).$$

On $[1/8, 1]$ the cubic factor is increasing because its derivative $3r^2 - 26r + 39$ is at least 16, and its value at $1/8$ is $857/512 > 0$. Since $1/8 < r < 1$, it follows that

$$\frac{s(7-r)}{6} < 1.$$

Consequently $C_T(p, h) + h/3 < 1$, which proves the strict $\mathcal{A}_T$ case. $\square$

Remark C.10 (Exact domain limitation). The hypothesis $p \geq 1/2$ cannot be removed. At

$$p = h = \frac{2}{5}$$

one has $d = 12/25 < 1$, $s = 4/5 < 1$, and $q = -44/625 < 0$, so the selected component is $\mathcal{A}_L$. At the proposed cap $c_0 = 13/15$,

$$P_{2/5}\left(\frac{13}{15}\right) = -\frac{14}{50625} < 0, \quad P_{2/5}(1) = \frac{6}{25} > 0.$$

Since $13/15 > \sqrt{3}/2$ and $P_{2/5}$ is strictly increasing on the selected root interval,

$$c_{\max}\left(\frac{2}{5}, \frac{2}{5}\right) > \frac{13}{15}.$$

Thus Lemma C.9 is a necessary relaxation only on its stated half-edge domain; it does not define a global replacement for the fibers B_c , F_c , or G_c .

Lemma C.11 (CE2 adjacent outer-ratio bound). *Let $0 < x, y, u, v < 1$, and put*

$$S = x + y, \quad D = \sqrt{x^2 + xy + y^2}.$$

Assume the exact CE2 relations

$$u + v = \frac{D + xy}{S}, \quad uS \geq x, \quad vS \geq y, \quad x^2 + xv + v^2 \leq 1.$$

Let $a_0, b_0 \geq 0$ satisfy

$$a_0 + b_0 > 1, \quad a_0^2 + a_0b_0 + b_0^2 \leq 1,$$

and define the two boundary tails by

$$A = \begin{cases} 1 - b_0, & b_0 < y, \\ \max\{0, 1 - \max\{b_0, v\}\}, & b_0 \geq y, \end{cases}$$

and

$$H = \begin{cases} 1 - a_0, & a_0 < x, \\ \max\{0, 1 - \max\{a_0, u\}\}, & a_0 \geq x. \end{cases}$$

If $H > 0$ and $A + H < 1/2$, then the center reach

$$d_2^C = \frac{D + xy - uS - y}{S} = v - \frac{y}{S}$$

satisfies

$$d_2^C < \frac{H}{3}.$$

Proof. Because $H > 0$, its definition leaves exactly two endpoint sources.

First source: $H = 1 - u$. Here $a_0 \geq x$ and $a_0 \leq u$. Define

$$K = x + 2y - xy - D.$$

Since $D + xy - y = S - K$, one has

$$d_2^C = 1 - u - \frac{K}{S} = H - \frac{K}{S}, \quad \frac{d_2^C}{H} = 1 - \frac{K}{SH}.$$

Put

$$b_o(x) = \frac{-x + \sqrt{4 - 3x^2}}{2}, \quad h(x) = b_o(x) - \frac{1}{2}.$$

We first claim

$$(98) \quad H \leq \min \left\{ \frac{y}{S}, h(x) \right\}.$$

The inequality $H \leq y/S$ follows from $H = 1 - u$ and $uS \geq x$. For the other inequality, $A + H < 1/2$ says that whichever of b_0 and v produces the $e_{0,1}$ tail is greater than $1/2 + H$. If it is b_0 , then $a_0 \geq x$ and the disk inequality give $b_0 \leq b_o(x)$. If it is v , the center-role diameter inequality $x^2 + xv + v^2 \leq 1$ gives $v \leq b_o(x)$. In either case $H < h(x)$, proving (98).

Set

$$m = \min \left\{ \frac{y}{S}, h(x) \right\}.$$

We prove $K/(Sm) > 2/3$. First suppose $y/S \leq h(x)$ and put $r = y/x$. Then

$$r \leq \frac{h(x)}{1 - h(x)}, \quad r \leq 1.$$

The exact inequality

$$\sqrt{1 + r + r^2} \leq 1 + \frac{r}{2} + \frac{r^2}{2}$$

follows because the difference of the squares is $r^2(r+1)^2/4$. Consequently

$$\frac{K}{y} \geq \frac{3}{2} - x - \frac{r}{2} \geq \frac{3}{2} - x - \frac{h(x)}{2(1 - h(x))}.$$

Write $b = h(x) + 1/2 = b_o(x)$. The diameter relation is symmetric, so $x = b_o(b)$. Define

$$F(b) = \frac{-b + \sqrt{4 - 3b^2}}{2} + \frac{b - 1/2}{3 - 2b}.$$

On $1/2 \leq b \leq 1$,

$$F'(b) = \frac{2}{(3 - 2b)^2} - \frac{1}{2} - \frac{3b}{2\sqrt{4 - 3b^2}} < 0.$$

For the last sign, put $z = 1 - b \in [0, 1/2]$. After moving the rational term and squaring the nonnegative sides, for $z > 0$ it is equivalent to

$$\begin{aligned} & 9(1 - z)^2(1 + 2z)^4 \\ & - (3 - 4z - 4z^2)^2(1 + 6z - 3z^2) > 0. \end{aligned}$$

The left side is $zP(z)$, where the Bernstein coefficients of the degree-5 polynomial P on $[0, 1/2]$ are

$$24, \quad 50, \quad \frac{364}{5}, \quad \frac{434}{5}, \quad \frac{432}{5}, \quad 72.$$

Thus F is decreasing, and

$$F(b) \leq F\left(\frac{1}{2}\right) = \frac{\sqrt{13}-1}{4}.$$

It follows that

$$\frac{K}{y} \geq \frac{7-\sqrt{13}}{4} > \frac{2}{3}.$$

Here $Sm = y$, so $K/(Sm) > 2/3$.

Now suppose $h(x) \leq y/S$, so $m = h(x)$. For fixed x , define

$$g(y) = \frac{x + 2y - xy - \sqrt{x^2 + xy + y^2}}{x + y}.$$

Then

$$g'(y) = \frac{x\{2(1-x)D + x - y\}}{2S^2D} > 0.$$

The sign is immediate for $y \leq x$. For $y > x$, the bound $D \geq y$ gives

$$2(1-x)D + x - y \geq x + (1-2x)y = x(1-y) + y(1-x) > 0.$$

Let y_0 satisfy $y_0/(x+y_0) = h(x)$. Since $y \geq y_0$,

$$\frac{K(x, y)}{Sh(x)} \geq \frac{K(x, y_0)}{(x+y_0)h(x)} = \frac{K(x, y_0)}{y_0} > \frac{2}{3},$$

where the last inequality is the boundary value from the preceding subcase. Hence $K/(Sm) > 2/3$ in all cases. Since $H \leq m$,

$$\frac{K}{SH} \geq \frac{K}{Sm} > \frac{2}{3},$$

and therefore $d_2^C/H < 1/3$.

Second source: $H = 1 - a_0$. Now $a_0 = 1 - H$. The $e_{0,1}$ tail cannot be $A = 1 - b_0$, since otherwise

$$A + H = 2 - a_0 - b_0 < \frac{1}{2}$$

would imply $a_0 + b_0 > 3/2$, contrary to $a_0 + b_0 \leq 2/\sqrt{3}$, which follows from the disk inequality. Thus $A = 1 - v$ and

$$v > \frac{1}{2} + H.$$

For v to determine the tail, T_0 reaches the start y of the CE2 interval, so $b_0 \geq y$. The disk inequality becomes

$$(1-H)^2 + (1-H)y + y^2 \leq 1,$$

or

$$H^2 - (2+y)H + y + y^2 \leq 0.$$

Put

$$L(y) = \frac{2+y-\sqrt{4-3y^2}}{2}.$$

Since $H < 1/2$, this gives $H \geq L(y)$. Moreover $uS \geq x$ and the CE2 coupling give

$$v \leq \frac{D+xy-x}{S},$$

and hence

$$(99) \quad \frac{D + xy - x}{S} > \frac{1}{2} + L(y).$$

We now prove the exact analytic implication needed here. Fix $0 < y < 1$, write $L = L(y)$, and suppose (99) holds. First,

$$\frac{D + xy - x}{S} < 1.$$

Indeed, both sides in $D < 2x + y - xy$ are positive, and

$$(2x + y - xy)^2 - D^2 = x\{y(3 - 2y) + x(1 - y)(3 - y)\} > 0.$$

Thus the premise forces $0 < L < 1/2$. Set

$$r = \frac{x}{y}, \quad R = \sqrt{r^2 + r + 1}, \quad r_0 = \frac{3 - 4L}{3 + 4L}.$$

Then

$$\frac{D + xy - x}{S} = \frac{R - r(1 - y)}{r + 1}, \quad \frac{D + xy - x - y}{S} = \frac{R - r(1 - y) - 1}{r + 1}.$$

Suppose first that $0 < r \leq r_0$. Using $R \leq 1 + r/2 + r^2/2$ gives

$$\frac{R - r(1 - y) - 1}{r + 1} \leq \frac{r(y - 1/2 + r/2)}{r + 1}.$$

The right side is convex in r , since its second derivative is $2(1 - y)/(1 + r)^3 > 0$, so its maximum on $[0, r_0]$ is at an endpoint. At $r = 0$ it is zero, while

$$\frac{L}{3} - \frac{r_0(y - 1/2 + r_0/2)}{r_0 + 1} = \frac{N(L)}{6(3 + 4L)},$$

where

$$N(L) = 16L^2y - 8L^2 + 18L - 9y, \quad y = \frac{L - 1 + \sqrt{1 + 6L - 3L^2}}{2}.$$

Write $s = \sqrt{1 + 6L - 3L^2}$. Then $N(0) = 0$ and

$$2sN'(L) = s(48L^2 - 64L + 27) - 144L^3 + 240L^2 + 59L - 27.$$

For $0 < L < 1/2$, one has $48L^2 - 64L + 27 > 0$ and

$$s \geq 1 + 3L - 6L^2,$$

because the right side is positive and the difference of the squares is $36L^3(1 - L)$. Therefore

$$2sN'(L) \geq 2L(-144L^3 + 192L^2 - 33L + 38) > 0.$$

The Bernstein coefficients of the last cubic on $[0, 1/2]$ are

$$38, \quad \frac{65}{2}, \quad 43, \quad \frac{103}{2},$$

so its positivity is exact. Thus $N(L) > 0$, proving

$$(100) \quad \frac{D + xy - x - y}{S} < \frac{L}{3}$$

when $r \leq r_0$.

It remains to exclude the premise when $r \geq r_0$. Put

$$J(r) = \frac{R - r(1 - y)}{r + 1}.$$

Its derivative has the sign of

$$\frac{r-1}{2R} - (1-y).$$

The function $(r-1)/(2R)$ is strictly increasing, since

$$\frac{d}{dr} \left(\frac{r-1}{2R} \right) = \frac{3(r+1)}{4R^3} > 0.$$

Hence J decreases and then increases, with at most one turning point, so its maximum on $[r_0, \infty)$ occurs at r_0 or in the limit. The limit is y , and

$$y < \frac{1}{2} + L$$

because $(3-y)^2 - (4-3y^2) = 4y^2 - 6y + 5 > 0$. At $r = r_0$, again using $R \leq 1 + r/2 + r^2/2$,

$$\frac{1}{2} + L - J(r_0) \geq \frac{N(L)}{(3+4L)^2} > 0.$$

Thus $J(r) < 1/2 + L$ for every $r \geq r_0$, contradicting the premise. This completes the proof of (100).

Finally,

$$d_2^C = v - \frac{y}{S} \leq \frac{D + xy - x - y}{S} < \frac{L(y)}{3} \leq \frac{H}{3}.$$

The two endpoint sources are exhaustive, so the lemma follows. $\square$

C.5. The exactly-one-Vd1/Vd2 placements. We now assume CE2, $N_+ = 1$, and exactly one row is Vd1 or Vd2. The corner normal form of Proposition 3.10 will be used in the following form: for some $t > 0$ and $d = \sqrt{t^2 + t + 1}$,

$$(101) \quad x - (t+1)y \leq a, \quad ty - (t+1)x \leq tb, \quad tx + y \leq d - a - tb.$$

In particular $a + b < 1/2$ for a positive-support row.

Lemma C.12 (Vd1 adjacent rescue). *Assume an original open-role skeleton cover in the CE2 exact- M_0 branch, with $T_C \cap \{M_0, \dots, M_5\} = \{M_0\}$. Suppose T_0 is the unique Vd1 row, $M_1 \in T_0$, T_1 is uniquely supercritical, and $T_2, \dots, T_5$ are nonsupercritical Vd0 rows. Suppose also that these are the only positive-adjacent-support roles. Then the roles do not cover the perimeter together with r_1 .*

Proof. From (101), the r_1 interval is

$$[c, u], \quad c = \frac{t(1-b)}{t+1}, \quad u = \frac{d-a-tb-1}{t}.$$

The midpoint inequalities and absence of positive r_5 support force $t \geq 1$. They also give

$$a \leq F(t, c) := d - 1 - \frac{3t}{2} + c(t+1) \leq L(c) := \sqrt{3} - \frac{5}{2} + 2c.$$

With $A(c) = 1 - \mathcal{B}(c)$, squaring positive sides gives

$$2L(c) \leq A(c),$$

because the difference is $4(20c^2 + (18\sqrt{3} - 52)c + 47 - 24\sqrt{3}) > 0$ on $[0, 1/2]$. Writing $\delta = 1 - u$, the same normal form yields

$$\frac{a}{a+\delta} \leq \frac{L(c)}{L(c)+1/2} \leq 2L(c) \leq A(c).$$

Thus $a \leq A(c)$. If the CE2 interval on $e_{5,0}$ hides the gap after T_0 , its side parameters X, Y, λ, ρ satisfy $Y \geq \lambda\delta$ and $S \leq a$, whence

$$T = X + \lambda \leq a + \lambda(u - a) \leq \frac{a}{a + \delta} \leq A(c).$$

In every other ordering its forced tail starts no later than a . Consequently the far-row reach is always $h \geq \mathcal{B}(c)$.

Coverage of r_1 gives the supercritical row a radial demand at least c . Therefore Lemma C.6 gives $b_1 < \mathcal{B}(c) \leq h$. The four remaining row caps would then have to satisfy

$$\sum_{i=2}^5 (a_i + b_i) \geq (1 - b_1) + 1 + 1 + 1 + h > 4,$$

a contradiction. $\square$

Lemma C.13 (Adjacent positive-support placement). *Assume an original open-role skeleton cover in the CE2 exact- M_0 branch, with $T_C \cap \{M_0, \dots, M_5\} = \{M_0\}$. Suppose T_0 is uniquely supercritical, T_1 is the unique Vd1/Vd2 row, and T_2, T_3, T_4, T_5 are nonsupercritical Vd0 rows; in particular, there is no additional positive-adjacent-support row. Then the perimeter together with r_1, r_2 is not covered. The reflected T_5 placement is also impossible.*

Proof. Use the CE2 variables $S = x + y$, $D = \sqrt{x^2 + xy + y^2}$ and $E = vS - y$. The center length on r_2 , measured from O , is $d_2^C = E/S$. Let A, H be the boundary demands forced on T_1 from $e_{0,1}$ and the propagated $e_{5,0}$ tail. The corner cap gives

$$(102) \quad A + H < \frac{1}{2}.$$

Apply Lemma C.9 with $p = 1/2 + A$ and $h = H$. Equation (102) gives

$$p \geq \frac{1}{2}, \quad 0 \leq h \leq p, \quad p + h < 1,$$

so every hypothesis of that lemma is present. It gives

$$(103) \quad c_{\max} \left(\frac{1}{2} + A, H \right) \leq 1 - \frac{H}{3}.$$

The exact CE2 coupling, the disk constraints for T_0 and the center role, and (102) are precisely the hypotheses of Lemma C.11. Therefore

$$(104) \quad d_2^C < \frac{H}{3}.$$

Now r_2 can be covered only by T_2 , the center, and a possible bridge from T_1 . The normal form makes a bridge imply $H < d_2^C$, contrary to (104). Without a bridge, boundary propagation gives T_2 inputs strictly exceeding $1/2 + A$ and H ; then (103) gives $c_2 \leq 1 - H/3 < 1 - d_2^C$, so T_2 cannot meet the center interval either. $\square$

Lemma C.14 (Nonadjacent positive-support placements). *Assume an original open-role skeleton cover in the CE2 exact- M_0 branch, with $T_C \cap \{M_0, \dots, M_5\} = \{M_0\}$. Suppose T_0 is uniquely supercritical, T_τ is the unique Vd1/Vd2 row for $\tau \in \{2, 3, 4\}$, and every other vertex row is nonsupercritical Vd0; in particular, there is no additional positive-adjacent-support row. Then the perimeter together with r_τ is not covered.*

Proof. Let A, H be the two boundary demands propagated to T_τ , and put $B = 1 - A$, $U = 1 - H$. The distance endpoint

$$\beta(z) = \frac{-z + \sqrt{4 - 3z^2}}{2}$$

gives $B \leq \beta(x)$ and $U \leq \beta(y)$, whether an extent comes from the supercritical row or from the corresponding CE2 endpoint. The exact CE2 comparison

$$(105) \quad \frac{D + xy}{S} + \beta(x) < 2$$

follows by squaring: with $b = \beta(x)$, the required quadratic in y is concave, positive at $y = 0$, and at $y = 1$ its numerator reduces to $4x^2(x^4 + 4x^3 + 9x^2 + 11x + 8)$. Reflection gives the y version. Since $u + v = (D + xy)/S$, it follows that

$$(106) \quad u + v - 1 < \min(A, H) =: m.$$

Put $P = vS - y$ and $Q = uS - x$. Then $P + Q = (u + v - 1)S$, while the center lengths on r_2, r_3, r_4 are

$$\frac{P}{S}, \quad \min \left\{ \frac{P}{y}, \frac{Q}{x} \right\}, \quad \frac{Q}{S}.$$

Equation (106) makes each strictly less than m . On the other hand (101) gives $c_\tau < 1 - m$. Hence the vertex-side demand $1 - d_\tau^C$ is strictly larger than c_τ , leaving r_τ uncovered. $\square$

Lemma C.15 (Vd1-supercritical replacement).

Assume an original open-role perimeter-and-radial-skeleton cover in the CE2 exact- M_0 branch and the complementary no-additional-positive-support case. Let T_i , $i \neq 0$, be uniquely supercritical, and let the unique Vd1 row, distinct from T_0 and adjacent to T_i , contain M_i . Assume every other vertex row is nonsupercritical Vd0 and the center has no boundary trace on the shared edge of this pair. Then the pair can be replaced by two open nonsupercritical Vd0 roles preserving all boundary and radial demands used by Proposition C.2.

Proof. Normalize the Vd1 row at V_0 and the supercritical row at V_1 . By the shared-edge hypothesis, no center trace intervenes on $e_{0,1}$; by the no-additional-positive-support hypothesis, no third vertex role has positive boundary or radial support in the handoffs used below. In (101), midpoint rescue and one-sided support imply $t \geq 1$ and, with

$$c = \frac{d - a - tb}{t + 1}, \quad \lambda = \frac{t(1 - b)}{t + 1}, \quad \mu = \frac{d - a - tb - 1}{t},$$

give the strict consequences

$$(107) \quad a + c < 1, \quad a < \lambda \leq \frac{1}{2}, \quad b < \frac{1}{2}, \quad \mu < 1 - a.$$

The shared edge forces the supercritical input $a_i \geq 1 - b > 1/2$, and radial coverage forces $c_i \geq \lambda$. A direct consequence of Proposition A.2 is

$$(x, y, z) \in \mathcal{A}, \quad x \geq 1/2, \quad z \leq 1/2 \implies y \leq 1 - z.$$

Indeed, substituting $x = 1/2, y = 1 - z$ in the selected supercritical polynomial gives $z^2(2z - 5)(2z - 1)/4 \geq 0$, and the polynomial is increasing in both relevant

coordinates. Hence $b_i \leq 1 - c_i \leq 1 - \lambda$ and $a + b_i < 1$. Open radial handoff further gives

$$c_i^{\text{req}} < \max\{c_i, \mu\} < \max\{1 - b_i, 1 - a\}.$$

Choose $a < p_1 < p_2 < 1 - b_i$ within these strict margins. In local coordinates use the unit triangles

$$\Delta_p^- = \text{conv}\{(0, 1 - p), (1, 1 - p), (0, -p)\}, \quad p \leq 1/2,$$

and

$$\Delta_p^+ = \text{conv}\{(p, 0), (p, 1), (p - 1, 0)\}, \quad p \geq 1/2.$$

Their incident reaches are $p, 1 - p$, their own-radial reach is respectively $1 - p, p$, and they have zero adjacent support. Small translations, chosen below every strict margin above, make the two replacements open Vd0 roles with boundary sum < 1 while preserving the shared boundary and both radial handoffs. Thus the datum reduces to Proposition C.2. $\square$

Proposition C.16 (Complementary exact Vd1/Vd2 placements). *Consider the CE2 exact- M_0 , $N_+ = 1$ branch with exactly one Vd1 or Vd2 role. Assume that no other vertex row has positive adjacent support and that the unique Vd2 role, if present, covers neither neighboring midpoint. Then the configuration is impossible.*

Proof. By the vertex classification, the no-additional-positive-support hypothesis makes every vertex row other than the unique Vd1/Vd2 row Vd0; all such rows other than the unique supercritical row are nonsupercritical. This is precisely the complementary branch in which Lemmas C.12–C.15 apply.

Let T_σ be uniquely supercritical and T_τ the unique Vd1/Vd2 row. They are distinct by the corner cap. If $\sigma = 0$, the adjacent placements $\tau = 1, 5$ are excluded by Lemma C.13, and $\tau = 2, 3, 4$ by Lemma C.14. If $\tau = 0$, midpoint forcing makes $\sigma = 1$ or 5 and T_0 cover that midpoint. Vd1 is excluded by Lemma C.12; Vd2 would cover the neighboring midpoint M_σ , contrary to the proposition's hypothesis.

If neither index is zero, midpoint forcing makes the two rows adjacent and the positive-support row rescue the supercritical midpoint. Vd2 is again excluded by the neighboring-midpoint hypothesis; Vd1 reduces by Lemma C.15 to the $N_+ = 0$ all-Vd0 obstruction. These alternatives exhaust the complementary exact-placement branch. $\square$

C.6. Exact-demand terminal routing.

Proposition C.17 (Exact-demand terminal branches). *The exact-demand method proves the following terminal exclusions:*

- (1) CE1 or CE2, $N_+ = 0$, all vertex roles Vd0;
- (2) CE1 or CE2, $N_+ = 0$, with at least one T3-like role and no Vd1/Vd2;
- (3) CE1 or CE2, $N_+ = 1$, all vertex roles Vd0, with at least one boundary gap;
- (4) CE1 or CE2, $N_+ = 1$, exactly one T3-like role and no Vd1/Vd2;
- (5) the complementary exact-placement subcases of CE2, $N_+ = 1$, with exactly one Vd1 or Vd2: no additional positive-support row and no Vd2 neighboring-midpoint rescue.

All gap alternatives include singleton gaps.

Proof. The five items are respectively Propositions C.2, C.5, B.3, C.1, and C.16. The strictness at open handoffs appears in the weak endpoint bounds, the nonattained supercritical envelope, and the strict replacement margins; no positive-length assumption is made about a V-gap itself. $\square$

APPENDIX D. THE DIRECT NINE-POINT OBSTRUCTION

This section supplies the exact frontier, witness, sign, and cap-overlap calculations for the direct nine-point obstruction.

This section proves a center-class-independent theorem. Its hypotheses are that all six vertex roles are Vd0, that they cover the full boundary, and that exactly one actual boundary row is supercritical. The theorem will therefore apply both to the CE0 branch and to the zero-gap subcases of CE1 and CE2. Six common radial points and three asymmetric points are excluded directly from all six actual vertex roles and are therefore forced into the center role. Only the unique supercritical row uses an AB -union; no nonsupercritical model core is introduced.

D.1. The finite caliper criterion. We first record the finite caliper certificate used below. It enters the proof through the exact statement and verification argument given here; neither a plot nor an unrecorded numerical convention is used.

Let R and J denote counterclockwise rotations through $2\pi/3$ and $\pi/2$, respectively. For a nonempty finite set $S \subset \mathbb{R}^2$ and a vector N , put

$$(108) \quad W_S(N) = \sum_{k=0}^2 \max_{z \in S} \langle z, R^k N \rangle.$$

Theorem D.1 (Finite caliper certificate). *If S has at least two distinct points, then S is contained in a closed unit equilateral triangle if and only if there are distinct $p, q \in S$ and $\varepsilon \in \{-1, 1\}$ for which*

$$(109) \quad W_S(\varepsilon J(q - p)) \leq \frac{\sqrt{3}}{2} \|p - q\|.$$

If S consists of one point, it is contained in equilateral triangles of arbitrarily small positive side length.

For $S_x = \{O, A, B, x\}$, every clause of (109) is a finite polynomial clause of degree at most four in the coordinates of A, B, x ; at most $12 \cdot 4^3$ clauses are required, and repeated point labels merely delete zero-length pairs.

Proof. For a unit vector n , write

$$G_S(n) = \sum_{k=0}^2 h_S(R^k n), \quad h_S(n) = \max_{z \in S} \langle z, n \rangle.$$

The three supporting half-planes with normals n, Rn, R^2n form an equilateral triangle of side $2G_S(n)/\sqrt{3}$. Hence S fits in a closed unit equilateral triangle precisely when $\min_{\|n\|=1} G_S(n) \leq \sqrt{3}/2$.

On an open normal-direction cell on which the three support points are fixed, $G_S(\theta) = c \cos \theta + d \sin \theta$ and $G_S'' = -G_S < 0$ unless all points coincide. Thus a minimum occurs at a cell boundary, where some support line contains two distinct points p, q . After a cyclic rotation of the three normals, its normal is $\varepsilon J(q - p)/\|p - q\|$. Homogeneity yields (109). A nondegenerate segment is checked directly with its two normals.

For the polynomial expansion fix (p, q, ε) , put $N = \varepsilon J(q - p)$ and $N_k = R^k N$, and choose support labels $z_0, z_1, z_2 \in S_x$. The support cell is

$$(110) \quad \langle z_k, N_k \rangle \geq \langle z, N_k \rangle \quad (z \in S_x, \quad k = 0, 1, 2),$$

with $\|p - q\|^2 > 0$. Since $W_S(N) \geq 0$, the remaining test is

$$(111) \quad 4 \left(\sum_{k=0}^2 \langle z_k, N_k \rangle \right)^2 \leq 3 \|p - q\|^2.$$

The points are affine-linear in the parameters, so (110)–(111) have degree at most four. Six unordered pairs, two signs, and 4^3 support-label triples give the asserted finite exhaustive list. $\square$

D.2. Corner-clipped AB -unions. Let e_A, e_B be unit vectors with angle 120° , put

$$O = 0, \quad A = ae_A, \quad B = be_B, \quad C = \{ue_A + ve_B : u, v \geq 0\},$$

and define

$$(112) \quad \mathcal{R}(a, b) = C \cap \bigcup \{T : T \text{ is a closed unit equilateral triangle and } O, A, B \in T\}.$$

The finite description in Theorem D.1 applies to the four-point set $\{O, A, B, x\}$ and makes membership in (112) an exact finite semialgebraic condition of degree at most four.

The strict supercritical branch admits a simpler closed form. In the next statement write $x = ue_A + ve_B$ and $\|x\|^2 = u^2 + v^2 - uv$.

Theorem D.2 (Strict supercritical AB -union). *Assume*

$$a + b > 1, \quad \rho = a^2 + ab + b^2 < 1,$$

put $h = \sqrt{3}/2$, $D = \sqrt{4\rho - 3}$, and define

$$\begin{aligned} \alpha &= \frac{h(a + 2b - aD)}{2\rho}, & \beta &= \frac{h(a - b + (a + b)D)}{2\rho}, \\ \gamma &= \frac{h(-a + b + (a + b)D)}{2\rho}, & \delta &= \frac{h(2a + b - bD)}{2\rho}. \end{aligned}$$

Then all four coefficients are positive and

$$(113) \quad \mathcal{R}(a, b) = (C \cap \mathcal{D}_A \cap \mathcal{H}_A) \cup (C \cap \mathcal{D}_B \cap \mathcal{H}_B),$$

where

$$\begin{aligned} \mathcal{D}_A &= \{(u, v) : (u - a)^2 + v^2 - (u - a)v \leq 1\}, \\ \mathcal{D}_B &= \{(u, v) : u^2 + (v - b)^2 - u(v - b) \leq 1\}, \\ \mathcal{H}_A &= \{(u, v) : \alpha(u - a) + \beta v \leq 0\}, \\ \mathcal{H}_B &= \{(u, v) : \gamma u + \delta(v - b) \leq 0\}. \end{aligned}$$

Its non-axis frontier consists, in counterclockwise order, of the A -circle, the A -line, the B -line, and the B -circle.

Proof. The identities

$$4\rho - 3(a+b)^2 = (a-b)^2, \quad (a+b)^2 D^2 - (a-b)^2 = 4\rho((a+b)^2 - 1)$$

give $0 < D < 1$ and positivity of the four coefficients.

We apply the caliper criterion of Theorem D.1. For a point $x = (u, v)$ in the relative interior of C but outside $\triangle OAB$, the four hull vertices occur in the order O, B, x, A . The two cone-axis hull edges cannot certify a unit triangle: their support sums, divided by h , are respectively

$$\max\{a, u\} + \max\{b, v - u\}, \quad \max\{b, v\} + \max\{a, u - v\},$$

and are at least $a + b > 1$. It remains to test the Ax and Bx calipers.

For the Ax caliper put $p = u - a$ and $r^2 = p^2 + v^2 - pv$. Exact evaluation of the three supports gives

(114)

$$\frac{G_A}{h} = \frac{\max\{r^2, -ap + (a+b)v\} + N_A}{r}, \quad N_A = \begin{cases} 0, & p \leq 0, \\ ap, & 0 \leq p \leq v, \\ (a+b)p - bv, & p \geq v. \end{cases}$$

No support label is omitted in these three sectors. If $p \leq 0$, $G_A \leq h$ is equivalent to

$$r \leq 1, \quad -ap + (a+b)v \leq r.$$

The second inequality is the half-plane condition in (113), because

$$r^2 - (-ap + (a+b)v)^2 = \frac{(cp + dv)(c'p + d'v)}{4\rho},$$

where

$c = a + 2b - aD$, $d = a - b + (a+b)D$, $c' = a + 2b + aD$, $d' = a - b - (a+b)D$, and $c, c', d > 0 > d'$. Thus the low sector is exactly $\mathcal{D}_A \cap \mathcal{H}_A$. In the middle sector $0 < p \leq v$ one has $r \leq v$, whereas the numerator in (114) is at least $(a+b)v > r$, so no fit is possible.

In the high sector $p \geq v$, the inequalities $G_A \leq h$ imply

$$(115) \quad r^2 + (a+b)p - bv \leq r, \quad bp + av \leq r.$$

Write $x - A = ry(\theta)$, $0 \leq \theta \leq 60^\circ$, with cone coordinates

$$y(\theta) = te_A + we_B, \quad t = \frac{2}{\sqrt{3}} \cos(\theta - 30^\circ), \quad w = \frac{2}{\sqrt{3}} \sin \theta.$$

Then (115) gives

$$r \leq f(\theta) = 1 - (a+b)t + bw, \quad S(\theta) = bt + aw \leq 1.$$

Let n_B be the unit vector with dual coordinates (γ, δ) . The identities

$$b\gamma + (a+b)\delta = h, \quad (a+b)\gamma + a\delta = \frac{h}{2}(1+D), \quad b\delta - a\gamma = \frac{h}{2}(1-D)$$

show that the unique angle θ_B of this normal satisfies $S(\theta_B) = 1$ and $f(\theta_B) = (1-D)/2$. The function S has at most one critical point, a maximum, so $S(\theta) \leq 1$ implies $\theta \leq \theta_B$. On the feasible part $f, f' \geq 0$; hence $f(\theta) \cos(\theta_B - \theta)$ is increasing. Consequently

$$\langle n_B, x - B \rangle \leq a\gamma - b\delta + \frac{h}{2}(1-D) = 0.$$

The diameter bound also gives $\|x - B\| \leq 1$. Thus every high sector fit belongs to $\mathcal{D}_B \cap \mathcal{H}_B$. Reflection exchanges a, u with b, v and supplies the corresponding result for the Bx caliper. This proves (113) in the cone interior; the axes follow by taking limits, since both sides are closed.

For completeness, the line–circle junctions are obtained by solving the two displayed line and circle equations. The line normals satisfy

$$\alpha^2 + \alpha\beta + \beta^2 = h^2, \quad \gamma^2 + \gamma\delta + \delta^2 = h^2,$$

and the line determinant is

$$\alpha\delta - \gamma\beta = \frac{3(1-D)(D+3)}{16\rho} > 0.$$

The successive cone slopes are

$$+\infty, \quad \frac{2}{1-D}, \quad \frac{a(1-D) + 2b}{2a + b(1-D)}, \quad \frac{1-D}{2}, \quad 0;$$

the two middle comparisons reduce to $a(4 - (1-D)^2) > 0$ and $b(4 - (1-D)^2) > 0$. This proves the asserted four-piece order. $\square$

D.3. Exact-one handoffs and common demands.

Lemma D.3 (Exact-one handoff chain). *Assume the six vertex roles cover ∂H and exactly one actual row is supercritical. Rotate its index to 4. The strict handoffs supplied by Theorem 2.9 satisfy*

$$x_3 < x_2 < x_1 < x_0 < x_5 < x_4.$$

With

$$a = 1 - x_3, \quad b = x_4, \quad p = 1 - b, \quad q = 1 - a,$$

one has

$$(116) \quad 0 < a, b < 1, \quad a + b > 1, \quad a^2 + ab + b^2 < 1,$$

and every selected row obeys

$$a_i \geq p, \quad b_i \geq q.$$

Proof. Every selected row other than 4 is strictly nonsupercritical, so $x_i < x_{i-1}$ for $i \neq 4$, which is precisely (D.3). In particular x_3 is the global minimum and x_4 the global maximum. The two anchors of row 4 lie in its open role; their mutual squared distance is $a^2 + ab + b^2$, strictly below the diameter squared of the closed unit triangle. This proves (116). Finally every $x_j \in [x_3, x_4]$, so $1 - x_{i-1} \geq 1 - x_4 = p$ and $x_i \geq x_3 = q$. $\square$

D.4. Six directly forced radial points. Put $h = \sqrt{3}/2$ and

$$s = p + q, \quad m = \min(p, q), \quad M = \max(p, q), \quad \chi = s^4 - s^2 + pq.$$

The strict domain (116) gives

$$(117) \quad pq > (1-s)(2-s), \quad m > 1-s, \quad s > 1-m.$$

Let $c_* = c_{\max}(p, q)$ be the exact radial envelope of Proposition A.2. On the present subcritical domain, equation (57) becomes

$$(118) \quad c_* = \begin{cases} C_L(m), & \chi \leq 0, \\ \frac{2M}{1 + \sqrt{4s^2 - 3}}, & \chi > 0, \end{cases}$$

where $C_L(m)$ is the unique root in $[\sqrt{3}/2, 1]$ of

$$z^4 - z^2 + mz - m^2 = 0.$$

At $\chi = 0$ the two expressions agree and equal s .

Lemma D.4 (Direct radial forcing). *One has*

$$(119) \quad 0 < c_* < 1, \quad c_* > 1 - m.$$

Assume that every U_i is Vd0 and that $U_C, U_0, \dots, U_5$ cover H . Then, for every i ,

$$D_i = (1 - c_*)V_i \in U_C.$$

Consequently U_C contains the centered closed disk

$$(120) \quad \mathcal{D}_\eta = \{x : \|x\| \leq \eta\}, \quad \eta = h(1 - c_*).$$

Proof. On the C_L branch, $C_L(m) < 1$ because the defining polynomial is positive at 1; if $s < \sqrt{3}/2$, then $c_* \geq \sqrt{3}/2 > s > 1 - m$, while if $s \geq \sqrt{3}/2$ its value at s is $\chi \leq 0$, so $c_* \geq s > 1 - m$. On the other branch put $E = \sqrt{4s^2 - 3}$. The selector gives $sE > M - m$, whence $c_* < s < 1$. Moreover, with

$$A_0 = \frac{2M}{1 - m} - 1,$$

one has

$$A_0^2 - E^2 = \frac{4(1 - s)\{(1 - m)(1 - 2m) + m(2 - m)(1 - s)\}}{(1 - m)^2} > 0.$$

Thus $A_0 > E$ and $c_* > 1 - m$. This proves (119).

Fix i . If $D_i \in U_i$, openness supplies $\varepsilon > 0$ with $c_* + \varepsilon < 1$ such that $(1 - c_* - \varepsilon)V_i \in U_i$. The same role contains its two selected handoff anchors. Since their incoming and outgoing demands dominate p and q , convexity and passage to the closure give a closed unit triangle realizing $(p, q, c_* + \varepsilon)$. This contradicts the definition of $c_* = c_{\max}(p, q)$. Hence $D_i \notin U_i$.

If D_i belonged to U_{i-1} or U_{i+1} , a small plane ball inside that open role would meet the adjacent radial arm r_i in a positive-length interval. This contradicts the Vd0 definition. For the remaining roles put $r = 1 - c_* > 0$. Directly,

$$\|rV_i - V_{i\pm 2}\|^2 = 1 + r + r^2 > 1, \quad \|rV_i - V_{i+3}\| = 1 + r > 1.$$

Since $V_j \in U_j$ and two points of an open unit triangle are at distance strictly below 1, these inequalities exclude D_i from the other three vertex roles. Thus no vertex role contains D_i . The point lies in $\text{int}(H)$ by (119), so the assumed cover forces $D_i \in U_C$.

The six points D_i form a regular hexagon of circumradius $1 - c_*$. Their convex hull contains its centered incircle of radius $h(1 - c_*)$, and convexity of U_C gives (120). $\square$

We next define the three asymmetric witnesses. At V_4 use local coordinates

$$P = V_4 + u(V_5 - V_4) + v(V_3 - V_4), \quad \mathbf{q}(u, v) = u^2 + v^2 - uv.$$

The outgoing demand is b and the incoming demand is a . Accordingly, set

$$\begin{aligned} \alpha &= \frac{h(b + 2a - bD)}{2\rho}, & \beta &= \frac{h(b - a + (a + b)D)}{2\rho}, \\ \gamma &= \frac{h(-b + a + (a + b)D)}{2\rho}, & \delta &= \frac{h(2b + a - aD)}{2\rho}, \end{aligned}$$

where now $\rho = a^2 + ab + b^2$ and $D = \sqrt{4\rho - 3}$. The two line pieces of Theorem D.2 are

$$L_-(\lambda) = (b - \beta\lambda, \alpha\lambda), \quad L_+(\mu) = (\delta\mu, a - \gamma\mu).$$

They meet at $\lambda_* = \mu_* = 8h\rho/(3(D+3))$ in

$$(121) \quad J = \left(\frac{2b + a - aD}{D+3}, \frac{b + 2a - bD}{D+3} \right).$$

Put

$$E_- = (1 - b, 2 - b), \quad E_+ = (2 - a, 1 - a).$$

The actual adjacent handoffs have local centers

$$C_5(t) = (1 + t, t), \quad 1 - a \leq t \leq b, \quad \text{and} \quad C_3(y) = (y, 1 + y), \quad 1 - b \leq y \leq a.$$

The restrictions of the two unit-circle equations are

$$(122) \quad \begin{aligned} g_-(\lambda) &= \frac{3}{4}\lambda^2 - 3(\alpha + b\beta)\lambda + 3b^2 - 3b + 2, \\ g_+(\mu) &= \frac{3}{4}\mu^2 - 3(\delta + a\gamma)\mu + 3a^2 - 3a + 2. \end{aligned}$$

Let λ_- and μ_+ be their first roots; explicitly,

$$\begin{aligned} \lambda_- &= 2(\alpha + b\beta) - \frac{2}{3}\sqrt{9(\alpha + b\beta)^2 - 9b^2 + 9b - 6}, \\ \mu_+ &= 2(\delta + a\gamma) - \frac{2}{3}\sqrt{9(\delta + a\gamma)^2 - 9a^2 + 9a - 6}. \end{aligned}$$

Lemma D.5 (Moving adjacent-role signs). *Throughout (116),*

$$(123) \quad \lambda_* < \lambda_- < h^{-1}, \quad \mu_* < \mu_+ < h^{-1}.$$

Moreover $J_u, J_v < 1/2$. The point $L_+(\mu_+)$ satisfies

$$(L_+(\mu_+))_u + (L_+(\mu_+))_v < 3 - 2a,$$

and the reflected bound holds for $L_-(\lambda_-)$. Writing $F(P, t) = \mathbf{q}(P - C_5(t)) - 1$, one has, for every $t \in [1 - a, b]$,

$$(124) \quad F(L_+(\mu_+), t) \geq 0, \quad F(J, t) > 0, \quad F(L_-(\lambda_-), t) > 0.$$

Under the reflection $(u, v, a, b) \leftrightarrow (v, u, b, a)$, the analogous signs hold for every $y \in [1 - b, a]$: the first-root point $L_-(\lambda_-)$ has nonnegative sign against $C_3(y)$, and the other two witnesses have strictly positive sign.

Proof. Put

$$U = a + b - 1, \quad z = a - b, \quad D^2 = z^2 + 6U + 3U^2.$$

Then $0 < U < 1/6$ and $z^2 < \Omega := 1 - 6U - 3U^2$. We give the fixed-line calculation for the circle centered at $E_+ = (2 - a, 1 - a)$; reflection gives the calculation for E_- . Put

$$F(P, t) = \mathbf{q}(P - (1 + t, t)) - 1, \quad t_0 = 1 - a,$$

so this circle is $F(P, t_0) = 0$. At the junction, exact expansion gives

$$(D + 3)^2 F(J, t_0) = 3D(z^2 + Uz + 1 - 3U) + P_J,$$

where

$$P_J = z^4 + 5z^2 + 6Uz^2 + 3U^2z^2 + 9Uz + 3U + 15U^2 > 0;$$

indeed $z^2 + Uz + 1 - 3U \geq 1 - 3U - U^2/4 > 0$, and $5z^2 + 9Uz + 15U^2$ is positive definite. Thus $F(J, t_0) > 0$. Also

$$J_u + J_v = \frac{(1+U)(3-D)}{3+D} < 1.$$

Indeed this inequality is $3U < D(2+U)$, which follows from $D^2 \geq 3U(2+U)$ and $(2+U)^3 > 3U$. Since

$$\frac{\partial F}{\partial t}(P, t) = 1 + 2t - P_u - P_v,$$

$F(J, t)$ is strictly increasing for $t \geq 0$, and therefore is positive throughout $[1-a, b]$.

On the opposite line L_- , the derivative at the junction, after multiplication by a positive factor, has at $t = 1-a$ the numerator

$$N_0 = D(9 + 3z^2 + 6Uz - 9U^2) + P_0,$$

where

$$P_0 = 18U^3 + 6U^2z^2 + 63U^2 + 18Uz^2 + 18Uz + 54U \\ + 2z^4 + 9z^2 + 9.$$

The coefficient of D is at least $9 - 6U - 9U^2 > 0$, while $P_0 \geq 9 + 54U - 18U > 0$. Hence $N_0 > 0$. At the other endpoint $t = b$, the corresponding numerator is

$$N_1 = D(9 + 3z^2 + 6U - 3U^2) + P_1,$$

where

$$P_1 = 9U^4 - 3U^3z + 45U^3 + 9U^2z^2 - 6U^2z + 72U^2 \\ - Uz^3 + 21Uz^2 + 9Uz + 45U + 2z^4 + 9z^2 + 9.$$

Its D -coefficient is positive and, using $|z| < 1$,

$$P_1 \geq 9 + 45U - 9U - U - 6U^2 - 3U^3 > 0.$$

The junction derivative is affine in t , so it is positive throughout $[1-a, b]$. Since the restriction of F to L_- is a convex quadratic, is positive at the junction, and is increasing there, it has no intersection on the opposite line side for any actual adjacent handoff.

On the correct line L_+ , the value at its outer endpoint h^{-1} has, after multiplication by a positive factor, numerator

$$A_0 = P_2 - 4DU(1 + U + z),$$

where

$$P_2 = 3U^4 + 6U^3z + 6U^3 + 4U^2z^2 + 12U^2z + 12U^2 \\ + 2Uz^3 + 6Uz^2 + 2Uz + 6U + z^4 + 2z^2 - 3.$$

For fixed U , replace the odd powers of z by $x = |z|$. The resulting polynomial $P_2^+(U, x)$ satisfies

$$\partial_x P_2^+ = 2(3U^3 + 4U^2x + 6U^2 + 3Ux^2 + 6Ux + U + 2x^3 + 2x) > 0,$$

and therefore its supremum for $z^2 < \Omega$ occurs at $x = \sqrt{\Omega}$. There

$$P_2^+(U, \sqrt{\Omega}) = 4U(U + \sqrt{\Omega} - 3) < 0.$$

As $1 + U + z = 2a > 0$, this gives $A_0 < 0$. The positive junction value and negative endpoint value force the first root strictly between them, proving the L_+ half of (123); reflection proves the other half.

For the coordinate bounds, the exact identities

$$D^2 - (3U - z)^2 = 6U(1 + z - U), \quad D^2 - (3U + z)^2 = 6U(1 - z - U)$$

give $J_u, J_v < 1/2$. It remains to control the coordinate sum along L_+ . For $E = L_+(h^{-1})$, the sign of $3 - 2a - E_u - E_v$ is the sign of

$$D(6U + 2z + 6) + B(U, z),$$

where

$$\begin{aligned} B(U, z) = & -9U^3 - 9U^2z - 9U^2 - 3Uz^2 - 18Uz + 3U \\ & - 3z^3 + 3z^2 - 3z + 3. \end{aligned}$$

Here

$$B_z = -3(3z^2 + 2Uz + 6U - 2z + 3U^2 + 1) < 0,$$

because the quadratic in parentheses has discriminant $-32U^2 - 80U - 8 < 0$. Hence its minimum on $z^2 < \Omega$ is at $z = \sqrt{\Omega}$, where $B(U, \sqrt{\Omega}) = 6(1 - 3U - \sqrt{\Omega}) > 0$ because $(1 - 3U)^2 - \Omega = 12U^2$. Thus $E_u + E_v < 3 - 2a$. Also $J_u + J_v < 1 < 3 - 2a$, and the coordinate sum is affine on the line segment, so the same strict bound holds at the first root. Now

$$\frac{\partial F}{\partial t}(P, t) = 1 + 2t - P_u - P_v.$$

Since $F(L_+(\mu_+), 1 - a) = 0$, the coordinate-sum bound makes its t -derivative strictly positive for $t \geq 1 - a$. Together with the no-opposite-line result, this proves (124). Reflection gives both the bound on L_- and the complete $C_3(y)$ statement. $\square$

Define the local and global witnesses by

$$(125) \quad Q_-^{\text{loc}} = L_-(\lambda_-), \quad Q_0^{\text{loc}} = J, \quad Q_+^{\text{loc}} = L_+(\mu_+),$$

and by the displayed conversion from (u, v) to the plane.

Lemma D.6 (Direct asymmetric forcing). *Assume that $U_C, U_0, \dots, U_5$ cover H . Then*

$$Q_-, Q_0, Q_+ \in \text{int}(H) \setminus \bigcup_{i=0}^5 U_i,$$

and consequently all three witnesses belong to U_C .

Proof. The root order and positivity of the line coordinates give $0 < u, v < 1$ for all three points. Each belongs to $\mathcal{R}(b, a)$ and hence lies in a closed unit triangle containing V_4 , so $\mathbf{q}(u, v) \leq 1$; hence $|u - v| < 1$ and all three points lie in $\text{int}(H)$.

The closure of U_4 is one of the triangles represented by $\mathcal{R}(b, a)$, since it contains V_4, X_3, X_4 . All three witnesses lie on its exact non-axis frontier. If one belonged to the open triangle U_4 , a sufficiently small plane ball about it would lie both in U_4 and in the interior of the corner cone, making the witness an interior point of $\mathcal{R}(b, a)$. This contradiction excludes U_4 .

For the roles U_0, U_1, U_2 , their contained vertices have local coordinates

$$\begin{array}{c|ccc} i & 0 & 1 & 2 \\ \hline V_i & (2, 1) & (2, 2) & (1, 2). \end{array}$$

For $P = (u, v)$,

$$\|P - (2, 1)\|^2 - 1 = \mathbf{q}(u, v) + 2 - 3u, \quad \|P - (1, 2)\|^2 - 1 = \mathbf{q}(u, v) + 2 - 3v.$$

The inequalities $J_u, J_v < 1/2$ and the line order give distance greater than one from V_0 for Q_-, Q_0 and from V_2 for Q_0, Q_+ . For the remaining pair, the first-root circle and the sum bound in Lemma D.5 give

$$\|Q_+ - V_0\|^2 - 1 = a(3 - a - (Q_+)_u - (Q_+)_v) > a^2,$$

and, by reflection, $\|Q_- - V_2\|^2 - 1 > b^2$. All three points are farther than one from $V_1 = (2, 2)$, because writing $u = 1 - \xi$, $v = 1 - \eta$ gives

$$\|P - V_1\|^2 = 1 + \xi + \eta + \xi^2 + \eta^2 - \xi\eta > 1.$$

Since two points in one open unit equilateral triangle have distance strictly less than one, these inequalities exclude U_0, U_1, U_2 .

The actual handoff $X_5 = C_5(x_5)$ belongs to U_5 , and $1 - a < x_5 < b$ by (D.3). The three signs in (124) therefore put every witness at distance at least one from X_5 , excluding U_5 . Similarly, if $y = 1 - x_2$, then $1 - b < y < a$, $X_2 = C_3(y) \in U_3$, and the reflected signs exclude U_3 . Thus no vertex role contains any witness. The assumed cover now forces all three interior witnesses into U_C . $\square$

The nine directly forced points are $D_0, \dots, D_5, Q_-, Q_0, Q_+$. Under a putative cover, Lemmas D.4 and D.6 therefore force the compact set

$$(126) \quad K_{\text{wit}}(a, b) = \mathcal{D}_\eta \cup \{Q_-, Q_0, Q_+\} \subset U_C.$$

D.5. Tangent reduction and exact mixed overlaps. For a compact set K , let $\Lambda(K)$ be the least side length of a closed equilateral triangle containing K . With R denoting rotation through $2\pi/3$,

$$(127) \quad \Lambda(K) = \frac{1}{h} \min_{\|n\|=1} \sum_{j=0}^2 h_K(R^j n).$$

For the nontrivial regime $c_* > 2/3$, we first remove the square radicals in the first-root witnesses. Normalize $\bar{\alpha} = \alpha/h$, and similarly for the other three coefficients, and put

$$\xi = \lambda/h, \quad \zeta = \mu/h, \quad \xi_* = \zeta_* = \frac{8\rho}{3(D+3)}.$$

Let g_-, g_+ be the rescaled forms of (122), and take one Newton step at the junction:

$$\widehat{\xi}_- = \xi_* - \frac{g_-(\xi_*)}{g'_-(\xi_*)}, \quad \widehat{\zeta}_+ = \zeta_* - \frac{g_+(\zeta_*)}{g'_+(\zeta_*)}.$$

Since each quadratic is strictly convex, positive and decreasing at the junction, its tangent zero lies strictly between the junction and the first root. Define, in local coordinates,

$$\begin{aligned} A &= \left(b - \frac{3}{4}\bar{\beta}\widehat{\xi}_-, \frac{3}{4}\bar{\alpha}\widehat{\xi}_- \right), \\ B &= J, \\ C &= \left(\frac{3}{4}\bar{\delta}\widehat{\zeta}_+, a - \frac{3}{4}\bar{\gamma}\widehat{\zeta}_+ \right), \end{aligned}$$

and convert them to global vectors. Then

$$A \in (Q_0, Q_-), \quad C \in (Q_0, Q_+).$$

Consequently, for

$$\widehat{K} = \text{conv}(\mathcal{D}_\eta \cup \{A, B, C\}),$$

we have $\widehat{K} \subseteq \text{conv}(K_{\text{wit}})$ and

$$(128) \quad \Lambda(K_{\text{wit}}) \geq \Lambda(\widehat{K}).$$

Put $S = \widehat{K} + R\widehat{K} + R^2\widehat{K}$. Then

$$h_S(n) = \sum_{j=0}^2 h_{\widehat{K}}(R^j n),$$

and S contains the full R -orbits of the four disks

$$(129) \quad \mathbb{D}(A, 2\eta), \quad \mathbb{D}(B, 2\eta), \quad \mathbb{D}(C, 2\eta), \quad \mathbb{D}(W, \eta), \quad W = C + RA.$$

For a disk $\mathbb{D}(X, r)$ let

$$I(X, r) = \{n : \|n\| = 1, \langle X, n \rangle + r \geq h\}.$$

D.6. The exact global certificate for the mixed overlaps. This subsection proves the two mixed-overlap signs used in Theorem D.9. It first replaces the exact radial value by rational upper envelopes, then reduces the mixed residuals to eight integer-polynomial signs. Three fixed analytic charts and twenty global Bernstein expansions prove those signs. No interval arithmetic, box subdivision, pruning, or branch-and-bound is used.

Retain the notation of Section D. Thus

$$p = 1 - b, \quad q = 1 - a, \quad s = p + q, \quad m = \min\{p, q\}, \quad M = \max\{p, q\},$$

$$\chi = s^4 - s^2 + mM, \quad h = \frac{\sqrt{3}}{2}, \quad c_* = c_{\max}(p, q).$$

The strict domain gives

$$(130) \quad mM > (1 - s)(2 - s), \quad m > 1 - s, \quad s > \frac{2}{3}.$$

Put

$$F_m(x) = x^4 - x^2 + mx - m^2, \quad \kappa = F'_m(s) = 4s^3 - 2s + m.$$

Since $m > 1 - s$,

$$(131) \quad \kappa > 4s^3 - 3s + 1 = (s + 1)(2s - 1)^2 > 0.$$

D.6.1. Rational radial upper envelopes. Define

$$(132) \quad \bar{c}_L = s - \frac{\chi}{\kappa} \quad (\chi \leq 0), \quad \bar{c}_T = s - \frac{\chi}{1 - m} \quad (\chi \geq 0).$$

Lemma D.7 (Branchwise radial upper bounds). *On the corresponding radial cells,*

$$c_* \leq \bar{c}_L < 1, \quad c_* \leq \bar{c}_T < 1.$$

At $\chi = 0$ both upper bounds equal $s = c_*$.

Proof. On the $C_L(m)$ cell, $F_m(s) = \chi \leq 0$ and $F_m(c_*) = 0$. For $x \geq s > 2/3$ one has $F''_m(x) = 12x^2 - 2 > 0$, so convexity and (131) give

$$0 = F_m(c_*) \geq F_m(s) + F'_m(s)(c_* - s),$$

whence $c_* \leq s - \chi/\kappa = \bar{c}_L$. Let $U = 1 - s$. From (130), $mM > U(1 + U)$, and therefore

$$\begin{aligned} \chi + U\kappa &> U\{m + 3s^3 - s^2 - 3s + 2\} \\ &= U\{m + 1 - 4U + 8U^2 - 3U^3\} > 0. \end{aligned}$$

Indeed $0 < U < 1/3$ and $1 - 4U + 8U^2 - 3U^3 > 1 - 4U + 7U^2 > 0$. Thus $\bar{c}_L < 1$.

On the other cell put

$$E = \sqrt{4s^2 - 3}, \quad m_0 = \frac{s(1-E)}{2}, \quad M_0 = \frac{s(1+E)}{2}.$$

Then

$$\chi = (m - m_0)(M_0 - m), \quad s - c_* = \frac{2(m - m_0)}{1 + E}.$$

Hence

$$\frac{\chi}{s - c_*} = \frac{1 + E}{2}(M_0 - m) \leq 1 - m,$$

because $0 \leq E \leq 1$ and $M_0 \leq 1$. This proves $c_* \leq \bar{c}_T$, while $\chi > 0$ gives $\bar{c}_T < s < 1$. $\square$

Reflect so that $z = a - b \geq 0$ and put $U = a + b - 1$. Then

$$s = 1 - U, \quad m = \frac{1 - U - z}{2}, \quad D^2 = z^2 + 6U + 3U^2 =: K(U, z),$$

$$\Xi(U, z) := 4\chi = 1 - 10U + 21U^2 - 16U^3 + 4U^4 - z^2,$$

$$G(U, z) := 2\kappa = 5 - 21U + 24U^2 - 8U^3 - z.$$

Consequently

$$(133) \quad \bar{c}_L = 1 - U - \frac{\Xi(U, z)}{2G(U, z)}, \quad \bar{c}_T = 1 - U - \frac{\Xi(U, z)}{2(1 + U + z)}.$$

Neither expression contains the quartic root or the radial radical $\sqrt{4s^2 - 3}$.

D.6.2. The Gram residuals and exact core reduction. On the active cell write $\bar{c}$ for the corresponding expression in (133), and put

$$\bar{\eta} = h(1 - \bar{c}), \quad e = 1 - \bar{c}, \quad t = 2\bar{c} - 1.$$

Lemma D.7 gives $0 < \bar{\eta} \leq \eta = h(1 - c_*)$ when $c_* > 2/3$. Thus the three disks obtained from $\mathbb{D}(C, 2\eta)$, $\mathbb{D}(W, \eta)$, and $\mathbb{D}(RA, 2\eta)$ by replacing η with $\bar{\eta}$ are contained in the disks already present in the Minkowski sum.

Represent a global vector as (x, hy) and put $C' = R^{-1}C$,

$$N_A = \|A\|^2, \quad N_C = \|C\|^2, \quad \nu = \langle A, C' \rangle, \quad \Delta_0 = x_A y_{C'} - y_A x_{C'}.$$

The Gram identity is

$$h^2 \Delta_0^2 = N_A N_C - \nu^2.$$

The two mixed square residuals for the smaller disks factor as

$$(134) \quad \bar{P}_A = h^2 N_A Q_A, \quad Q_A = \Delta_0^2 - \|tA - eC'\|^2,$$

$$(135) \quad \bar{P}_C = h^2 N_C Q_C, \quad Q_C = \Delta_0^2 - \|tC' - eA\|^2.$$

All witness coordinates are rational in U, z, D . Substituting (133) into (134)–(135) and reducing modulo $D^2 - K(U, z)$ gives, for $B \in \{L, T\}$ and $X \in \{A, C\}$,

$$(136) \quad \text{num}(Q_{B,X}) = 32(K + 3)^2 \{A_{B,X}(U, z) + DB_{B,X}(U, z)\},$$

where $A_{B,X}, B_{B,X} \in \mathbb{Z}[U, z]$. Every omitted denominator is strictly positive: it is a positive constant times $(D + 3)^4$, the square of the relevant upper-envelope

denominator, and squares of the two nonzero Newton-derivative numerators. The term counts are

(B, X)	$\#A_{B,X}$	$\#B_{B,X}$
(L, A)	347	298
(L, C)	347	298
(T, A)	342	294
(T, C)	341	294.

Since $D \geq 0$, it remains to prove all eight integer polynomials nonnegative.

D.6.3. *Three fixed charts and global positive bases.* Set

$$\mathcal{G}(U) = 1 - 6U - 3U^2, \quad H(U) = 1 - 10U + 21U^2 - 16U^3 + 4U^4,$$

$$U_H = 1 - \frac{\sqrt{3}}{2}, \quad U_{\max} = \frac{2}{\sqrt{3}} - 1.$$

The identities

$$4\chi = H(U) - z^2, \quad H(U) = (U - 1)^2(4U^2 - 8U + 1)$$

and

$$\mathcal{G}(U) - H(U) = 4U(1 - 6U + 4U^2 - U^3) \geq 0 \quad (0 \leq U \leq U_H)$$

show that the radial cells are exactly covered by

$$(137) \quad \begin{aligned} T : & \quad 0 \leq U \leq U_H, \quad 0 \leq z^2 \leq H(U), \\ L_0 : & \quad 0 \leq U \leq U_H, \quad H(U) \leq z^2 \leq \mathcal{G}(U), \\ L_1 : & \quad U_H \leq U \leq U_{\max}, \quad 0 \leq z^2 \leq \mathcal{G}(U). \end{aligned}$$

We use the elementary positive-basis fact that a polynomial whose coefficients in the tensor Bernstein basis on $[0, 1]^2$ are all nonnegative is nonnegative on that square. This is one global polynomial identity, not a subdivision procedure. All coefficients below lie in $\mathbb{Q}(\sqrt{3})$, and their signs are decided exactly by rational comparisons.

For the T cell use

$$(138) \quad U = \frac{(x-1)(3-x)}{4x}, \quad z = y \frac{9-x^4}{8x^2}, \quad 1 \leq x \leq \sqrt{3}, \quad 0 \leq y \leq 1.$$

Here

$$H\left(\frac{(x-1)(3-x)}{4x}\right) = \left(\frac{9-x^4}{8x^2}\right)^2,$$

so the chart covers the whole T cell. Clearing the positive powers of 2 and x , then putting $x = 1 + (\sqrt{3} - 1)r$, gives four global Bernstein expansions on $[0, 1]^2$:

	bidegree	coefficients	zero coefficients
$A_{T,A}$	(88, 22)	2047	2
$B_{T,A}$	(80, 20)	1701	0
$A_{T,C}$	(88, 22)	2047	2
$B_{T,C}$	(80, 20)	1701	0.

Every nonzero coefficient is strictly positive.

For an L -cell polynomial $F(U, z)$, write

$$F(U, z) = E_F(U, w) + zO_F(U, w), \quad w = z^2,$$

and put

$$R_F(U, w) = E_F(U, w)^2 - wO_F(U, w)^2.$$

The implications $E_F \geq 0$ and $R_F \geq 0$ give $E_F \geq \sqrt{w}|O_F|$, hence $F \geq 0$. Use the two fixed charts

$$(139) \quad U = U_H r, \quad w = H(U) + s\{\mathcal{G}(U) - H(U)\}$$

for L_0 , and

$$(140) \quad U = U_H + (U_{\max} - U_H)r, \quad w = s\mathcal{G}(U)$$

for L_1 , with $0 \leq r, s \leq 1$. The global Bernstein expansions of E_F and R_F have the following bidegrees and coefficient counts:

F	$L_0 : E_F$	$L_0 : R_F$	$L_1 : E_F$	$L_1 : R_F$
$A_{L,A}$	(46, 11); 564	(92, 22); 2139	(26, 11); 324	(52, 22); 1219
$B_{L,A}$	(42, 10); 473	(84, 20); 1785	(24, 10); 275	(48, 20); 1029
$A_{L,C}$	(46, 11); 564	(92, 22); 2139	(26, 11); 324	(52, 22); 1219
$B_{L,C}$	(42, 10); 473	(84, 20); 1785	(24, 10); 275	(48, 20); 1029

Every coefficient in these sixteen expansions is strictly positive. Therefore all eight polynomials in (136) are nonnegative on the corresponding radial cells.

Lemma D.8 (Exact mixed-cap overlaps). *In the regime $c_* > 2/3$, the tangent witnesses defined above satisfy*

$$I(C, 2\eta) \cap I(W, \eta) \neq \emptyset, \quad I(W, \eta) \cap I(RA, 2\eta) \neq \emptyset.$$

Proof. Since $D \geq 0$, the eight signs prove $Q_A, Q_C \geq 0$ for $z \geq 0$ by (136); reflection supplies $z \leq 0$. Equations (134)–(135) give the two mixed cap overlaps for the smaller disks, hence for the original disks. $\square$

D.7. The enclosure inequality.

Theorem D.9 (Witness enclosure). *For every (a, b) in (116),*

$$\Lambda(K_{\text{wit}}(a, b)) \geq 1.$$

Proof. The disk alone gives $\Lambda(K_{\text{wit}}) \geq 3\eta/h = 3(1 - c_*)$, so the assertion is automatic when $c_* \leq 2/3$. Assume henceforth $c_* > 2/3$.

The tangent construction above gives (128), the Minkowski sum S , and the four disks in (129). If the caps of the disks in (129) cover the unit circle, then $h_S(n) \geq h$ for every n and (127) proves $\Lambda(\hat{K}) \geq 1$.

Put $\tau = h - 2\eta = h(2c_* - 1)$. The rays of

$$(141) \quad A, B, C, W, RA$$

occur in this cyclic order from A to RA . Indeed the two line distances give

$$\frac{\text{cross}(A, B)}{\|B - A\|} = \alpha(1 - b) + \beta > 0, \quad \frac{\text{cross}(B, C)}{\|C - B\|} = \gamma + \delta(1 - a) > 0.$$

If $A = (-X, -Y)$ and $C = (-P, -Q)$ in the centered cone basis, then $0 < X, Y, P, Q < 1$, $X, Q > 1/2$, and

$$\frac{\text{cross}(C, RA)}{h} = PX + (Q - P)Y > 0.$$

The last sign is immediate unless $Q < P$ and $X < Y$, in which case it is larger than $Q - P/2 > 0$. The ray of $W = C + RA$ lies between those of C and RA . The same coordinates give

$$(142) \quad \|A\|, \|C\| > h/2 > \eta.$$

For the two equal-radius overlaps it is enough to prove

$$(143) \quad \frac{\text{cross}(A, B)}{\|B - A\|} \geq \tau, \quad \frac{\text{cross}(B, C)}{\|C - B\|} \geq \tau.$$

We record the exact analytic verification. By reflection take $a \leq b$ and set $m = 1 - b$, $M = 1 - a$, $U = a + b - 1$, $s = m + M$, and $w = M - m$. The smaller of the two normalized left sides in (143) is

$$(144) \quad d = \frac{\mathcal{A} + U(2m - 1) + D(\mathcal{A} + U)}{2\rho}, \quad \mathcal{A} = 1 - m + m^2.$$

On the C_L sheet, write $F_L(z) = z^4 - z^2 + mz - m^2$. This defining polynomial evaluated at $r = 1 - m/2 + m^2/2$ is

$$F_L(r) = \frac{m^2(m-1)}{16}(m^5 - 3m^4 + 11m^3 - 17m^2 + 28m - 12) > 0.$$

Indeed the polynomial in parentheses is increasing on $(0, 1/2)$ and has value $-33/32$ at $1/2$; its derivative is $28 - 34m + m^2(5m^2 - 12m + 33) > 0$. Since $F_L(h) = -(m - h/2)^2 \leq 0$ and the selected root is unique, $2c_* - 1 \leq \mathcal{A}$. Put

$$X_0 = \mathcal{A} + U, \quad Y_0 = 2\rho\mathcal{A} - \mathcal{A} - U(2m - 1).$$

Then

$$d - \mathcal{A} = \frac{D(\mathcal{A} + U) - Y_0}{2\rho},$$

and exact expansion gives

$$\begin{aligned} Y_0 &= 2(m^2 - m + 1)U^2 + (2m^3 - 2m + 3)U \\ &\quad + (m^2 - m + 1)(2m^2 - 2m + 1) > 0. \end{aligned}$$

Furthermore

$$D^2X_0^2 - Y_0^2 = 4m\rho\Phi(U, m),$$

where

$$\begin{aligned} \Phi(U, m) &= (1 - m)(m^2 - m + 2)U^2 \\ &\quad + (2 - m^2)(m^2 - m + 1)U \\ &\quad - m(1 - m)^2(m^2 - m + 1). \end{aligned}$$

In these variables

$$\chi = U^4 - 4U^3 + 5U^2 - (m + 2)U + m(1 - m).$$

Since $U^2(U^2 - 4U + 5) > 0$, the selector $\chi \leq 0$ implies $U > m(1 - m)/(m + 2)$; Φ is increasing in U and at that lower endpoint is

$$\frac{m^2(1 - m)(m^4 - 2m^3 + 6m^2 - 6m + 4)}{(m + 2)^2} > 0.$$

The last quartic equals $(m^2 - m)^2 + 5m^2 - 6m + 4 > 0$. Thus $d > \mathcal{A} \geq 2c_* - 1$.

On the other radial sheet let $E = \sqrt{4s^2 - 3}$. Then $0 \leq w < sE$ and $c_* = (s + w)/(1 + E)$. At fixed U , differentiation of (144) can be checked without an implicit estimate. Namely,

$$d = \frac{1 + D}{2} - UB(D, w), \quad B(D, w) = \frac{(1 + U)(3 + D) + w(1 - D)}{D^2 + 3}.$$

Here $D^2 = w^2 + 6U + 3U^2$, so $0 \leq D' = w/D \leq 1$, and

$$B_w = \frac{1 - D}{D^2 + 3} \geq 0.$$

Writing $N = (1 + U)(3 + D) + w(1 - D)$, direct differentiation gives

$$B_D = \frac{(1 + U - w)(D^2 + 3) - 2DN}{(D^2 + 3)^2} > -\frac{34}{27}.$$

Indeed the first term in the numerator is positive, while $1 + U - w = 2a > 0$, $N < (7/6)4 + 1 = 17/3$, $D < 1$, and $D^2 + 3 \geq 3$. If $B_D < 0$, then $B_D D' \geq B_D$; otherwise $B_D D' \geq 0$. Hence $B' = B_w + B_D D' > -34/27$, and $U < 1/6$ yields

$$d' < \frac{1}{2} + \frac{1}{6} \frac{34}{27} = \frac{115}{162} < 1,$$

whereas $(2c_* - 1)' = 2/(1 + E) \geq 1$. Hence their difference decreases with w . At the boundary $w = sE$, where $\chi = 0$ and $c_* = s$, the preceding sheet applies: this is an admissible point because $h < s < 1$ and $D^2 = 4s^4 - 12s + 9 < 1$. Indeed $s^4 - 3s + 2 = (s - 1)(s^3 + s^2 + s - 2) < 0$ on this interval; the second factor is increasing and already equals $(7h - 5)/4 > 0$ at $s = h$. The preceding sheet gives $d \geq 2s - 1$. Therefore (143) holds on both sheets, and reflection supplies the other line.

It remains to check the two mixed overlaps. On the two radial cells use the exact rational envelopes and global Bernstein identities established in Subsection D.6. Lemma D.8 gives precisely the overlaps for $\mathbb{D}(C, 2\eta)$ with $\mathbb{D}(W, \eta)$ and for $\mathbb{D}(W, \eta)$ with $\mathbb{D}(RA, 2\eta)$.

Thus the four consecutive pairs of caps centered on the rays (141) overlap. Every cap is a single arc of half-width less than $\pi/2$, and the positive crosses put each overlap in the intervening short sector. They cover the sector from A to RA ; rotation through $2\pi/3$ covers the circle. Hence $h_S \geq h$, so $\Lambda(\widehat{K}) \geq 1$. Equation (128) completes the proof. $\square$

D.8. The center-independent obstruction.

Theorem D.10 (Center-independent direct nine-point obstruction). *There is no cover of H by open unit equilateral triangles $U_C, U_0, \dots, U_5$ such that*

- (1) *every U_i is a Vd0 role at V_i ;*
- (2) *$U_0, \dots, U_5$ cover ∂H ; and*
- (3) *exactly one actual maximal boundary row is supercritical.*

No center class is assumed for U_C .

Proof. Lemma D.3 gives a parameter pair in (116). Direct radial forcing and direct asymmetric forcing give the compact containment $K_{\text{wit}}(a, b) \subset U_C$. Theorem D.9 says $\Lambda(K_{\text{wit}}) \geq 1$.

On the other hand, a compact subset of an open unit equilateral triangle has positive clearance from its three sides. Moving all three sides inward by a sufficiently small equal amount gives a closed equilateral triangle of side strictly less than one still containing the compact set. Thus $K_{\text{wit}} \subset U_C$ would imply $\Lambda(K_{\text{wit}}) < 1$, a contradiction. $\square$

Remark D.11 (Why exact-one is essential). The exact-one handoff theorem identifies the unique selected supercritical row and makes every other selected row a

strict descent. Thus (D.3) simultaneously provides the global minimum and maximum used in the common radial demands, puts the two actual adjacent handoffs in the intervals of Lemma D.5, and leaves only one strict AB -frontier to analyze. With two or more ascents this single chain need not exist. The $N_+ \geq 2$ branch is therefore handled by Strategy 3 or Strategy 1.

Proposition D.12 (Branches closed by direct nine-point forcing). *Theorem D.10 closes precisely the following rows of the final routing table:*

- (1) CE0, $N_+ = 1$, all six vertex roles Vd0;
- (2) CE1 or CE2, $N_+ = 1$, all six vertex roles Vd0, and zero boundary gaps.

Proof. In the first row, a missed boundary point would have to lie in the open CE0 center role, giving a positive-length center trace on an incident edge. Hence the six vertex roles cover ∂H . In the second row, zero gaps means by definition that the two adjacent open vertex traces cover every boundary edge, including every endpoint. Thus in both rows all three hypotheses of Theorem D.10 hold. $\square$

APPENDIX E. TABLE OF SYMBOLS

This table lists the notation used across more than one proof mechanism. Local auxiliary variables are defined where they occur.

symbol	meaning	first formal use
H, H_L	The regular hexagons of side lengths 1 and L .	Section 1.2
O, V_i	The center and cyclically indexed vertices of H .	Section 1.2
$e_{i,i+1}, r_i, M_i$	A boundary edge, a radial arm, and the midpoint $V_i/2$.	Section 1.2
$D, S, S_{1/2}$	The radial target, full skeleton, and half-skeleton.	Section 2
U_C, U_i	The original open center role and vertex roles in a hypothetical cover.	Section 1.2
T_C, T_i	The closures of U_C and U_i .	Section 1.2
CE0, CE1, CE2	The exhaustive center-role boundary-trace classes.	Section 1.2
Vd0, Vd1, Vd2, T3-like	The exhaustive vertex-role classes.	Section 1.2
(A_i, B_i, C_i)	Actual maximal incoming boundary, outgoing boundary, and radial reaches of U_i .	Section 1.2
(a_i, b_i, c_i)	Selected lower bounds on incoming, outgoing, and radial reaches, rather than the actual maxima defining N_+ .	Section 2
N_+	The number of actual rows satisfying $A_i + B_i > 1$.	Section 1.2
B_c, F_c, G_c	The outgoing feasible envelope, its nonsupercritical cap, and the induced demand map.	Section 4
$e(d)$	The smaller high-radial Cell- L root $\ell(1 - d)$ used as a demand threshold.	Section 4

symbol	meaning	first formal use
$\psi(x)$	The universal selected- T_+ increment $1 - \sqrt{1 - x + x^2}$.	Proposition 4.1
z	The optional rational parameter for the universal selected- T_+ curve.	Equation (25)
$f(a, b), G(a, b)$	Normalized maximal retained area and the corresponding local area loss.	Section 5
$c_*, \mathcal{D}_\eta$	The common radial envelope in the nine-point argument and its forced centered disk.	Section 6
Q_-, Q_0, Q_+	The three exact asymmetric witnesses on the strict supercritical frontier.	Section 6
A, B, C	The Newton inner points used in the rational four-cap reduction.	Section 6
$\hat{\xi}_-, \hat{\xi}_+$	The one-step Newton parameters on the two frontier lines.	Section 6
K_{wit}	The compact set consisting of the forced disk and three asymmetric witnesses.	Section 6
$\Lambda(K)$	The least side length of a closed equilateral triangle containing K .	Section 6
$I(X, r)$	The level- $\sqrt{3}/2$ support cap of the disk $\mathbb{D}(X, r)$.	Lemma 6.1

APPENDIX F. EXACT CERTIFICATE RECORD

The mathematical reduction and global positive-basis proof of the two mixed overlaps appear in Section D.6. This appendix records the associated exact data and replay interface; it introduces no additional hypothesis or numerical evidence into the proof.

The canonical data live with the proof package in
`proof/3XXX_CE0/31XX_Nplus1/310X_all_Vd0/
3105X_self_contained_direct_Vd0_nine_point/
3105X_computation/`

The module `mixed_overlap_core_polynomials.py` assembles the six numbered `mixed_overlap_core_data_*.py` shards. The two exact checks are

```
verify_mixed_overlap_core_derivation.py
verify_global_core_positivity.py
```

The first derives the eight core polynomials from the witness formulas and checks equality with the stored transcript. The second checks the twenty global Bernstein identities and the exact signs of their coefficients in $\mathbb{Q}(\sqrt{3})$. Neither check uses interval arithmetic, adaptive subdivision, pruning, or branch-and-bound.

Both checks authenticate the canonical core-polynomial data with SHA-256 digest

```
dc46aaf263655d5159ecd3a81db72ee82477951d06172f4743b248df37209485
```

and return

```
adaptive_subdivision: false
branch_and_bound: false
result: PASS
```

using exact integer, rational, and $\mathbb{Q}(\sqrt{3})$ arithmetic.